

## Nonlinear Equations

## 1. Bisection-Algorithm.

- (a) Improve the function `Bisekt`. Your `[x,y]=Bisection(f,a,b,tol)` should also compute a zero for functions with  $f(a) > 0$  and  $f(b) < 0$  to a given tolerance `tol`. Be careful to stop the iteration in case the user asks for a too small tolerance! If by the bisection process we arrive at an interval  $(a,b)$  which does not contain a machine number anymore then it is high time to stop the iteration.

*Solution:*

```
function [x,y]=Bisection(f,a,b,tol)
% BISECTION computes a root of a scalar equation
% [x,y]=Bisection(f,a,b,tol) finds a root x of the scalar function
% f in the interval [a,b] up to a tolerance tol. y is the
% function value at the solution

fa=f(a); v=1; if fa>0, v=-1; end;
if fa*f(b)>0
    error('f(a) and f(b) have the same sign')
end
if ( nargin<4), tol=0; end;
x=(a+b)/2;
while (b-a>tol) & ((a < x) & (x<b))
    if v*f(x)>0, b=x; else a=x; end;
    x=(a+b)/2;
end
if nargin==2, y=f(x); end;
```

- (b) Solve with bisection the equations

$$a) \quad x^x = 50 \quad b) \quad \ln(x) = \cos(x) \quad c) \quad x + e^x = 0.$$

Hint: a starting interval is easy to find by sketching the functions involved.

*Solution:*

a) The function  $x^x$  is monotonically increasing. Since  $1^1 = 1$  and  $4^4 = 256$  the values  $a = 1$  and  $b = 4$  can be used for the bisection. The solution becomes

```
>> [x,f]=Bisection(@(x) x^x-50,1,4)
x =
    3.287262195355581
f =
    7.105427357601002e-15
```

b) Drawing the functions  $\ln(x)$  and  $\cos(x)$  we see that their cutting point is in the interval  $(0, \pi/2)$ , thus

```
>> [x,f]=Bisection(@(x) log(x)-cos(x),0,pi)
x =
    1.302964001216012
f =
   -2.220446049250313e-16
```

c) We write the equation  $e^x = -x$  and from the graph of the two functions we get the interval  $(-1, 0)$  for the solution, so

```
>> [x,f]=Bisection(@(x) exp(x)+x,-1,0)
x =
   -0.567143290409784
f =
   -1.110223024625157e-16
```

2. Find  $x$  such that

$$f(x) = \int_0^x e^{-t^2} dt - 0.5 = 0.$$

**Hint:** the integral cannot be evaluated analytically, so expand it in a series and integrate. Write a function `f(x)` to evaluate the series.

Since a function evaluation is expensive (summation of the Taylor series) but the derivatives are cheap to compute, a higher order method is appropriate. Solve this equation with Newton's or Halley's method.

*Solution:*

Take the series for  $e^x$ , substitute  $x = -t^2$  and integrate to get the expansion

$$\int_0^x e^{-t^2} dt = x - \frac{x^3}{1! \cdot 3} + \frac{x^5}{2! \cdot 5} - \frac{x^7}{3! \cdot 7} + \frac{x^9}{4! \cdot 9} \mp \dots \quad (1)$$

For evaluating the series we introduce the expressions

$$ta := (-1)^{i-1} \frac{x^{2i-1}}{(i-1)!} \quad t := (-1)^i \frac{x^{2i+1}}{i!}$$

then  $t = -ta * x^2 / i$  and the partial sum is updated by  $s_{\text{new}} = s_{\text{old}} + t / (2 * i + 1)$ . We will stop the summation when  $s_{\text{new}} = s_{\text{old}}$ . Thus we get

```
function y = ff(x);
% is used in IntegralExp.m
t = x; snew = x; sold=0; i=0;
while sold ~= snew
    i = i+1;
    sold = snew;
```

```

    t = -t*x^2/i;
    snew = sold+t/(2*i+1);
end
y = snew;

% Solve \int_{0}^{x} e^{-t^2}dt - 0.5 = 0 with Newton and Halley
% use ff.m to compute Taylor series
%
format compact
format long

disp(' Newton')
x = 1; xa=2;
while abs(xa-x)>1e-10
    xa=x;
    y = ff(x)-0.5; ys = exp(-x^2);
    x = x - y/ys
end

disp('Halley')
x = 1; xa=2;
while abs(xa-x)>1e-10
    xa=x;
    y = ff(x)-0.5; ys = exp(-x^2); yss = -2*x*ys;
    t = y*yss/ys^2;
    x = x - y/ys/(1-0.5*t)
end

>> IntegralExp
Newton
x =
    0.329062444950818
x =
    0.532365165339031
x =
    0.550852862865461
x =
    0.551039408434969
x =
    0.551039427609027
x =
    0.551039427609027
Halley
x =
    0.598466410057177
x =
    0.551087168834467
x =
    0.551039427609074

```

$\mathbf{x} =$   
0.551039427609027

3. Compute the intersection points of an ellipsoid with a sphere and a plane. The ellipsoid has the equation

$$\left(\frac{x_1}{3}\right)^2 + \left(\frac{x_2}{4}\right)^2 + \left(\frac{x_3}{5}\right)^2 = 3.$$

The plane is given by  $x_1 - 2x_2 + x_3 = 0$  and the sphere has the equation  $x_1^2 + x_2^2 + x_3^2 = 49$

- How many solutions do you expect for this problem?
- Solve the problem with the `solve` and `fsolve` commands from MAPLE.
- Write a MATLAB script to solve the three equations with Newton's method. Vary the initial values so that you get all the solutions.

*Solution:*

- The intersection of the plane with the sphere is a circle and with the ellipsoid we get an ellipse. The intersection points are therefore the intersections of a circle with an ellipse. We expect thus in general 4 different solutions. Looking at the equations we notice that with a solution  $\mathbf{x}$  also  $-\mathbf{x}$  is a solution. Thus if we have found two different solutions then the two remaining solutions are given by changing the signs.
- With the MAPLE commands

```
eqs:={ (x1/3)^2+(x2/4)^2+(x3/5)^2=3, x1-2*x2+x3=0, x1^2+x2^2+x3^2=49};  
solve(eqs, {x1, x2, x3});
```

we obtain

$$\begin{aligned}x1 &= -\frac{31634}{247975} \left( \text{RootOf} \left( -3621220 \_Z^2 + 63268 \_Z^4 + 50197225 \right) \right)^3 \\ &\quad + \frac{180746}{49595} \text{RootOf} \left( -3621220 \_Z^2 + 63268 \_Z^4 + 50197225 \right), \\x2 &= -\frac{15817}{247975} \left( \text{RootOf} \left( -3621220 \_Z^2 + 63268 \_Z^4 + 50197225 \right) \right)^3 \\ &\quad + \frac{230341}{99190} \text{RootOf} \left( -3621220 \_Z^2 + 63268 \_Z^4 + 50197225 \right), \\x3 &= \text{RootOf} \left( -3621220 \_Z^2 + 63268 \_Z^4 + 50197225 \right)\end{aligned}$$

Using the numerical solver MAPLE delivers only one solution

```
fsolve(eqs, {x1, x2, x3});
```

$$\{x1 = -3.101446015, x2 = -3.977629342, x3 = -4.853812670\}$$

- (c) Programming in MATLAB Newton's algorithm we obtain      With the starting vector  $\mathbf{x} = [1, 1, 1]$  we get convergence in 8 steps:

```
x =
    1
    1
    1
ans =
 1.414125519548956e+01
ans =
 6.353644140517445e+00
ans =
 2.186622967262920e+00
ans =
 3.581210720347850e-01
ans =
 1.070558121458718e-02
ans =
 9.718696508742296e-06
ans =
 8.014862832469677e-12
ans =
 1.199111836538749e-15
k =
    8
x =
 3.101446014850100e+00
 3.977629342271496e+00
 4.853812669692894e+00
```

The quadratic convergence is visible from the printout of the norm of the correction vector  $\mathbf{h}$ . Using the starting vector  $\mathbf{x} = [-4, 1, 6]$  we converge to the second solution in 4 steps

```
x =
   -4
    1
    6
ans =
 2.872708652671924e-01
ans =
 7.067873458438775e-03
ans =
 5.582183986704225e-06
ans =
 4.059746603817636e-12
k =
    4
x =
-3.781770586037488e+00
```

```
1.010696349931241e+00
5.803163285899970e+00
```

4. Modify the fractal program by replacing  $f(z) = z^3 - 1$  with the function

$$f(z) = z^5 - 1.$$

- (a) Compute the 5 zeros of  $f$  using the command `roots`.
- (b) In order to distinguish the 5 different numbers, study the imaginary parts of the 5 zeros. Invent a transformation such that the zeros are replaced by 5 different positive integer numbers.

*Solution:* We first compute the zeros of  $z^5 - 1$ . The coefficients of the polynomial  $z^5 - 1$  are

```
p=[1 0 0 0 0 -1]
```

With the function `roots` we can compute the zeros

```
>> W=roots(p)
W =
-0.809016994374948 + 0.587785252292473i
-0.809016994374948 - 0.587785252292473i
 0.309016994374947 + 0.951056516295152i
 0.309016994374947 - 0.951056516295152i
 1.000000000000000 + 0.000000000000000i
```

If we multiply the imaginary part by 2 we get

```
>> 2*imag(W)
ans =
 1.175570504584946
-1.175570504584946
 1.902113032590305
-1.902113032590305
 0
```

Now we can add 3 and round the result to get

```
>> round(2*imag(W)+3)
ans =
 4
 2
 5
 1
 3
```

```

n=1000; m=30;
x=-1:2/n:1;
[X,Y]=meshgrid(x,x);
Z=X+1i*Y; % define grid for picture
for i=1:m % perform m iterations in parallel
    Z=Z-(Z.^5-1)./(5*Z.^4); % for all million points
end; % each element of Z contains one root
a=10;
image(round(2*imag(Z)+3)*a);

```

