


Diverzita savců

Asi 5 500 recentních druhů

Nové objevy, popisy a nálezy
New discoveries, descriptions and finds




Podle:

Jan Hošek, 2007: Saola aneb
největší zoologické objevy
posledních let.

Scientia, Praha 2007, 215 str.

Počty žijících druhů obratlovců podle přehledů uveřejněných v posledních třech letech a jejich ohrožení podle Červeného seznamu IUCN.

<i>skupina</i>	<i>odhad počtu známých druhů</i>	<i>počet vymřelých druhů</i>	<i>počet ohrožených druhů</i>
savci (Mammalia)	5416	70	1093
ptáci (Aves)	9934	135	1206
plazi („Reptilia“)	8240	22	341
obojživelníci (Amphibia)	5918	34	1811
svaloploutví (Sarcopterygii)	8	0	1
paprskoploutvé ryby (Actinopterygii)	26848	80	1058
paryby (Chondrichthyes)	970	0	110
kruhoústí (Cyclostomata)	108	0	2
celkem	57442	342	5621



1901

Okapia johnstoni

-
-
-



- g. horská


nda – pohoří Virunga

rageni – prase pralesní



ká, J-Etiopie






„ZLATÝ VĚK OBJEVŮ“

Kopytníci od roku 1900 (Ungulates)

- 1901 okapi pruhovaná (*Okapia johnstoni*) – Kongo, Uganda (Giraffidae)
- mazama yucatánský (*Mazama pandora*) – Mexiko (
- chocholotka Weynsova (*Cephalophus weynsi*) – Afri
- chocholotka ruwensorská (*Cephalophus rubidus*)
- 1903 antilopka zakrská (*Neotragus batesi*) – rovníková A
- 1904 prase pralesní (*Hylochoerus meinertzhageni*) –
- 1908 mazama venezuelský (*Mazama bricenii*) – J Amerik
- 1910 nyala horská (*Tragelaphus buxtoni*) - J Etiopie (mazama
- anoa horský (*Bubalus quarlesi*) – Celebes (Sulawesi) (Bovidae)
- 1911 dikdik somálský (*Madoqua piacentinii*) – Somálsko (Bovidae)
- 1914 goral červený (*Nemorhaedus baileyi*) – Tibet, Barma (B
- 1918 kl (Cephalophus adersi) – V Afric
- 1929 ka (schus berezovskii) – J Čína, S
- (M
- 1930 p (onius objev
- ta)
- 1932 m (ntiacu – J Asie (Cervidae)
- 1935 gazela arabská (*Gazella sau* ý polc
- 1937 kuprej (*Bos sauveli*) – Zadní Ir
- 1959 mazama zakrslý (*Mazama chun* vidae)
- 1963 nahur Schaeferův (*Pseudois sch* ng-c'-t



- 1981 kabar čínský (*Moschus fuscus*) – Čína (Bovidae)
- 1982 muntžak žlutý (*Muntiacus atherodes*) – Borneo (Cervidae)
- 1985 gazela jemenská (*Gazella bilkis*) – Jemen (Bovidae)
- 1987 prase floreské (*Sus heurenii*) – J Asie (Suidae)
- 1990 muntžak gongšanský (*Muntiacus gongshanensis*) – J Asie (Tibet, Čína), (Cervidae)
- 1993 saola (*Pseudoryx nghetinhensis*)** - Vietnam-Laos (Bovidae)
- 1994 muntžak obrovský (*Megamuntiacus vuquangensis*)** – Vietnam-Laos (Cervidae)
- lyrorožec - ling (*Pseudonovibos spiralis*)** – Vietnam, Kambodža (Bovidae)
- 1996 mazama bororo (*Mazama bororo*) – JV Brazílie (Cervidae)
- 1997 muntžak černý (*Muntiacus truongsonensis*) – stř. Vietnam (Cervidae)
- 1999 muntžak listový (*Muntiacus putaoensis*) – Myanmar (Barma) (Cervidae)
- 2003 buvolec zambijský (*Damaliscus superstes*) – Zambie, Kongo (Bovidae)
- 2005 kančil cejlonský (*Moschiola kathygre*) – Sri Lanka (Tragulidae – kančilovití)
- voduška konžská (*Cobus anelli*) – Kongo (Bovidae)
- 2008 jelínek mazamu (*Mazama ochroleuca*) – Amazónie




kančil

1937

Bos sauveli

kuprej





letters to nature

Nature 363, 443 - 445 (03 June 1993); doi:10.1038/363443a0

THIS ARTICLE IS

[Available as PDF](#)[Full Article](#)[Full Text](#)[Request reprints](#)[Send to a friend](#)[More articles like this](#)[Table of Contents](#)[Previous issue](#)

A new species of living bovid from Vietnam

VU QUANG UNG, PHAM HONG QUOC, NGUYEN HUU CHINH (DOT LOC), PETER ARCTANDORA, JOHN MACKINNON

Forest Inventory and Planning Institute, Ministry of Forestry, Hanoi, Vietnam

Asia In Bureau for Conservation, 18/E Capital Building, 1757 19 1 Lockhart Road, Wanchai, Hong Kong
To whom correspondence should be addressed.

IN May 1992 a joint survey by the Ministry of Forestry and World Wide Fund for Nature of the Vu Quang Nature Reserve, Ha Tinh province, found three sets of long straight horns of a new bovid (Mammalia, Artiodactyla) in hunters' houses¹. None of the specimens had dentition. On four follow-up visits by Vietnamese scientists new specimens were discovered and surveys of forests in neighbouring Nghe an province revealed more localities and some partial specimens. In all, we have examined more than 20 specimens. Three have complete upper skulls and dentitions, two have lower jaws and dentitions. Three complete skins have been collected. The specimens are distinct in appearance, morphology and DNA sequence and cannot be ascribed to any known genus. Only two bovid genera are known from this part of Asia, *Bos* and *Naemorhedus* = *Capricornis*^{2,3}. A new genus and species are therefore




Muntiacus vuquangensis – muntžak obrovský 1994



Muntiacus truongsonensis – muntžak černý 1997



Muntiacus putaoensis – muntžak listový 1999



Pseudonovibos spiralis – ling 1994




Figure 1. Phylogenetic tree reconstructed from the combination of cytochrome b and lactoferrin sequences. Shown is a bootstrap 50 % majority-rule consensus tree performed with the maximum parsimony (MP) method [26]. *Antilocapra americana* family Antilocapridae, *Cervus elaphus* and *Odocoileus hemionus* (family Cervidae) were used to root the bovid tree. The analyses were conducted with either equal weighting, or differential weighting of the nucleotide transformations (see section 5.2). Bootstrap proportions were computed after 1 000 replicates. These are indicated below branches, to the left of the slash for the weighted MP analysis and to the right for the unweighted MP analysis. The type and position of the molecular signatures characteristic of the different groups of species (i.e. exclusive synapomorphies), are indicated on the branches as shown in the key.



Moschiola kathygre – kančil cejlonský 2005

Kytovci od roku 1900 (Cetacean, Whale)

- 1908 vorvaňovec australský (*Mesoplodon bowdoini*)
- 1912 sviňucha jižní (*Australophocaena dioptrica*)
- 1913 vorvaňovec tmavý (*Mesoplodon mirus*)
- 1918 delfínovec čínský (*Lipotes vexillifer*)
- 1926 vorvaňovec Longmanův (*Indocetus pacificus*)
- 1934 delfín Graffmanův (*Stenella graffmani*)
- 1937 vorvaňovec Shepherdův (*Tasmacetus shepherdi*)
- 1956 plískavice saravacká (*Lagenodelphis hosei*)
- 1958 vorvaňovec japonský (*Mesoplodon ginkkodens*)
sviňucha kalifornská (*Phocoena sinus*)
- 1963 vorvaňovec kalifornský (*Mesoplodon carlhubbsi*)
- 1991 vorvaňovec peruánský (*Mesoplodon peruvianus*)
- 2002 vorvaňovec Perrinův (*Mesoplodon perrini*) – DNA
- 2003 plejtvák Omurův (*Balaenoptera amurai*) – DNA
- 2005 orcela tupoploutvá (*Orcaella heinsohni*)

Letouni od roku 2000


2000	3 druhy
2001	6 druhů, v Evropě <i>Myotis alcaethoe</i> – netopýr alcaethoe (menší, nymfin)
2002	8 druhů
2003	3 druhy
2004	10 druhů, <i>Plecotus gaisleri</i> – S Afrika, <i>Pipistrellus hanaki</i> - Libye
2005	13 druhů
2006	9 druhů
2007	1 druh, <i>Styloctenium mindorensis</i>
2011	1 druh, <i>Niumbaha superba</i> – J Sudán

Reeder DM, Helgen KM, Vodzak ME, Lunde DP, Ejotre I., 2013. A new genus for a rare African vespertilionid bat: insights from South Sudan. *ZooKeys*, 285: 89-115. doi: 10.3897/zookeys.285.4892



Ostatní savci po roce 1990

- 1995 prase vietnamské (*Sus bucculentus*)– Laos, popis 1892, Lazarus taxon
králík Timminsův (*Nesolagus timminsi*) – Laos
- 1996 khanyou (*Laonastes aenigmaemus*) – stř. Laos – skalní krysa
(Laonastidae), Lazarus taxon, bazální linie hystricognátních hlodavců
- 1997 langur duk (*Pygathrix nemaeus*) – Vietnam (Cercopithecidae)
cibetka tainguenská (*Viverra zibetha*) – Vietnam, Annamity
- 2001 slon pralesní (*Loxodonta cyclotis*) – kryptický druh, DNA, dnes nepřijímán



- BIODIVERSITÀ





Figure 2. Global distribution of new mammals described since 1992. The distribution is overlaid on currently recognized regions of high threat and irreplaceability. Variable levels of shading indicate the number of global biodiversity conservation templates that prioritize the region (Brooks et al. 2005).

SAVCI – distribuce nových druhů objevených po 1992



Titi - 23 druhy

Jižní Amerika

Primáti

1998 *Callithrix humilis* – kosman (*Callithrichidae*, drápkaté opice)

2000 *Callithrix manicorensis*

Callithrix acariensis

2003 *Callicebus bernhardi* – titi (*Cebidae* – malpovití)

Calicebus stephennashi

? *Callicebus sp.nov.* 1

Callicebus sp.nov. 2

Pithecia sp.nov. 1 - chvostan

Pithecia sp.nov. 2

Saguinus sp.nov. 1 - tamarín

Saguinus sp.nov. 2

Ateles sp.nov. 1 - chápan

Ateles sp.nov. 2

Cacajao sp.nov. - uakari

Lagothrix sp.nov. 1 - chápan

Lagothrix sp.nov. 2

Callithrix sp.nov. - kosman

Callicebus sp. nov. - titi

Saimiri sp.nov. - kotul

Ostatní

Agouti sp. nov. – paka (Agoutidae)

Eira sp.nov. – hyrare, brazilská kuna (Mustelidae)



Nasua sp.nov. – nosál (Procyonidae)

Myrmecophaga sp.nov. – mravenečník (Myrmecophagidae)

Pteronura sp.nov. – vydra (Mustelidae)

Panthera sp.nov. – jaguár (Felidae)





Marc van Roosmalen

24.7. 1947, Tilburg, Holanďan žijící v brazilském Manau, primatolog

<http://www.marcvanroosmalen.org/news.htm>



marcvanroosmalen.org

ENTER

NAMED NEW SPECIES

Black-crowned dwarf marmoset

Callibella humilis – kosman černošlý

Rio Acarí Amazonian marmoset

Mico (Callithrix) acariensis – kosman akarský

Satarè Amazonian marmoset

Mico (Callithrix) saterei - kosman brazilský

Rio Manicoré Amazonian marmoset

Mico (Callithrix) manicorensis - kosman

Stephen Nash's titi monkey

Callicebus stephennashi - titi

HRH Prince Bernhard's titi monkey

Callicebus bernhardi- titi

Giant collared peccary

Pecari maximus – pekari obrovský

Van Roosmalen's dwarf porcupine

Sphiggurus roosmalenorum - kuandu

Dwarf manatee

Trichechus pygmaeus sp. nov. - kapustňák

Lecythidaceae (Brazilnut Family)

Lecythis oldemani sp. nov. - rostlina

Black dwarf lowland tapir

Tapirus pygmaeus sp. nov. - tapír

Van Tienhoven's fair brocket deer

Mazama tienhoveni sp. nov. - jelínek

NEW SPECIES SEARCH

Arboreal giant anteater - *Myrmecophaga sp. nov.*

White-throated black jaguar - *Panthera sp. nov.* - jaguár

Black giant otter - *Pteronura sp. nov.* - vydra

Orange coati-mundè - *Nasua sp. nov.* - nosál

Orange tayra - *Eira sp. nov.* - kuna

Black woolly monkey - *Lagothrix sp. nov.* – chápan vlnatý

Cruz Lima's saddleback tamarin monkey

Saguinus (fuscicollis) cruzlimai sp. nov. - tamarin

Rio Pauini white bald-headed uacari

Cacajao (calvus) sp. nov. - uakari

Rio Aripuanã green-backed squirrel monkey

Saimiri (ustus) sp. nov. - kotul

Rio Mampurú titi monkey

Callicebus (moloch) sp. nov. - titi

Upper Xingú Amazonian marmoset monkey -titi

Mico (Callithrix) sp. nov. - kosman



sp. nov. - chápan

teles sp. nov. - chápan

sp. nov. - chápan

sp. nov. - tamarin

bus (torquatus) sp. nov.

bus (moloch) sp. nov.

Kotul - http://www.youtube.com/watch?v=KJe_7d1oRwg&noredirect=1

A NEW SPECIES OF LIVING PECCARY (MAMMALIA: TAYASSUIDAE)
FROM THE BRAZILIAN AMAZON

Marc G. M. VAN ROOSMALEN, Lothar FRENZ, Pim VAN HOOFT, Hans H. DE IONGH & Herwig LEIRS



Pecari maximus 2007 – největší pekari

A NEW SPECIES OF LIVING MANATEE FROM THE AMAZON

Shallow clear-water adapted dwarf manatee is already on the verge of extinction

Marc G.M. van Roosmalen¹, Pim van Hooft² & Hans H. de Iongh³

¹ AAPA Manaus-Amazonas, Brazil

² Wageningen University and Research Centre, Resource Ecology Group, Bornsesteeg 69, 6708PD
Wageningen, The Netherlands

³ Leiden University, Institute of Environmental Sciences, PO Box 9518, 2300RA Leiden, The Netherlands



(LEFT) The Rio Arauazinho harbors dwarf manatee and numerous other new mega-fauna species. Marc proposes this be the cornerstone for a new Brazilian National Park.



Trichechus bernhardi 2007

- nejmenší kapustňák

A NEW SPECIES OF LIVING BROCKET DEER
(MAMMALIA: CERVIDAE) FROM THE BRAZILIAN AMAZON

Marc G. M. VAN ROOSMALEN & Pim VAN HOOFT



(LEFT) *Mazama ochroleuca* sp. nov. redrawn from plate depicting *Mazama (gouazoupira) nemorivaga* (Eisenberg, 1989).

(ABOVE) Two spikes of *Mazama americana* above, one of *Mazama ochroleuca* sp. nov. below.

Here we report on the existence of a new species of even-toed ungulate in the Brazilian Amazon, which we name *Mazama ochroleuca* sp. nov., the fair brocket deer. It is intermediate in size between the two known species of brocket deer, *Mazama americana* and *Mazama (gouazoupira) nemorivaga*, and occurs in sympatry with both. Preliminary mitochondrial partial cytochrome b sequences of fair brocket deer compared with that of the sympatric and morphologically most related grey brocket deer {*Mazama (gouazoupira) nemorivaga*} revealed a sequence difference of 3.7%. Divergence time is therefore estimated at 1.0 million years before present. As in other brocket deer, fair brocket deer seem to live solitary or in pairs. In view of recent developments in the Rio Aripuanã basin where it lives and due to its limited distribution, we consider the fair brocket deer highly endangered.

KEY WORDS

New species, Artiodactyla, Cervidae, *Mazama ochroleuca* sp. nov., fair brocket deer, Brazilian Amazon

Mazama ochroleuca 2008

A NEW SPECIES OF LIVING LOWLAND TAPIR
(MAMMALIA: TAPIRIDAE) FROM THE BRAZILIAN AMAZON

Marc G.M. van Roosmalen / AAPN Manaus-Amazonas, Brazil



(LEFT) Adapted rendering of a black dwarf lowland tapir.
(ABOVE) On the left a skull of Brazilian lowland tapir, on the right the skull of black dwarf lowland tapir.

Here we report on the existence of a new species of odd-toed ungulate in the Brazilian Amazon, which we name *Tapirus pygmaeus sp. nov.*, the black dwarf lowland tapir. It is much smaller than the since long known Brazilian lowland tapir, *Tapirus terrestris*. As in other tapirs, dwarf tapirs seem to live solitary or in pairs. In view of recent developments in the interfluves where it lives and due to its limited distribution and apparent rariness, we consider the dwarf tapir highly endangered.

KEY WORDS

New species, Perissodactyla, Tapiridae, *Tapirus pygmaeus sp. nov.*, black dwarf lowland tapir, Brazilian Amazon

Tapirus pygmaeus 2008 – nejmenší tapír

Současné tempo popisování


- RYBY cca 150 druhů ročně
- OBOJŽIVELNÍCI cca 70 druhů ročně
- PTÁCI cca 5 - 10 druhů ročně
- SAVCI cca 40 – 50 druhů ročně

1992 – 2005


ptáci vs. savci

83 : 341

Log of Birds (+) and Mammals (o) described per year



Log of Birds (-) and Mammals (- -) described per year



REEDER ET AL.—NEW MAMMALS CONTINUE TO BE DISCOVERED






Figure 1. Cumulative and decadal descriptions of taxonomically valid extant mammal species.

Patterson 2001




Patterson 2001




Chiroptera


Patterson 2001



Patterson 2001




Patterson 2001





Rodentia

Patterson 2001



Patterson 2001







Andy

Cuscomys ashaninka

– činčilák ašaninka (Abrocomidae – činčilákovití) – 1999 L.H. Emmonsová



Abrocoma oblativa Eaton 1916
Cuscomys oblativus

Cuscomys ashaninka

1992 – 2005

341 nový druh savců

1992 – 2005

Rodentia 155 druhů



Mallomys sp. nov. – obří krysa (Murinae)

Papua – Nová Guinea

1992 – 2005

Chiroptera 78 druhů



Styloctenium mindorensis Esselstyn 2007 – kaloň, nížinné lesy Filipín

1995 – 2007

36 nových druhů primátů

2006

Tarsius lariang – nártoun (nártounovití – Tarsiidae)
Microcebus jollyae – maki (makiovití - Cheirogaleidae)
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamiratra
Lepilemur aeeclis – lemur (lemurovití noční – Megaladapidae)
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmasoni
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi – avahi (indriovití – Indridae)
Cebus queirozi – malpa (malpovití – Cebidae)

2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mambiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mampiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mambiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrightii
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mamiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrightii
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mambiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahalazensis
Lepilemur ahmansoni
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mamiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrightii
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mamiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mamiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang _____
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mambiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrightii
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsii
Microcebus mamiratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcockii
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsonii
Lepilemur wrighti
Avahi peyrierasi
Cebus flavius




Lophocebus kipunji *Rungwecebus*



Tanzánie 2005

mangabej kipundži – kočkodanovití (Cercopithecidae)





because of the terrain, thick secondary forest, and the animal's cryptic nature, sightings were infrequent and poor. It was not until December 2003, during work in the contiguous Livingstone Forest, that the monkey was clearly observed and recognized as a new species of mangabey.

Ndundulu population. The Udzungwa Mountains (Udzungwas: 10,000 km², 07°40'S to 08°40'S and 35°10'E to 36°50'E) lie 350 km to the northeast of Rungwe-Livingstone (4, 5). Supporting circa (ca.) 1017 km² of fragmented forest (6), the Udzungwas receive a maximum annual rainfall of roughly 2200 mm and were previously thought to hold 10 primate species (7), including the endemic Sanje mangabey, *Cercocebus sanjei*, discovered in 1979 (8).

Two populations of the Sanje mangabey are known from the Udzungwas (7, 9). During visits from 1991 to 2000, ornithologists working in the Ndundulu Forest Reserve (Fig. 1) reported a third population of the Sanje mangabey (10). Subsequent surveys failed to confirm the presence of this species in Ndundulu (7, 9, 11) and led to our intensified surveys in July and September 2004. During these surveys, Sanje mangabeys were not encountered or heard. However, on 7 July 2004, the new species of mangabey was discovered. It now seems certain that the ornithologists had misidentified the new species of mangabey as the Sanje mangabey.

The researchers working on each of these two new populations of mangabeys did not become aware that a second population was known until October 2004.

Lophocebus kipunji Ehardt, Butynski, Jones, and Davenport sp. nov.



Fig. 2. Holotype: adult male highland mangabey *Lophocebus kipunji* in the type locality, Rungwe-Livingstone, Tanzania. [Photograph by T.R.B. Davenport]

live individual should be collected at this time to serve as the holotype. The Rungwe-Livingstone population is designated the source population for physical specimens in support of the holotype.

Paratype. Adult in photograph (Fig. 3). Sex not known. Photograph taken in Ndundulu Forest Reserve (07°48'45"S 36°31'05"E), Udzungwa Mountains, Tanzania.



Fig. 3. Paratype: adult highland mangabey *Lophocebus kipunji*, Ndundulu Forest Reserve, Tanzania. [Photograph by T. Jones]

tail off-white. Crown with very long, broad, erect crest of hair. Eyelids black, not contrasting with color of face. Adults emit a distinctive, loud, low-pitched "honk-bark" (Fig. 4). Arboreal. Found only at high altitudes (1300 up to 2450 m asl) and low-temperature tolerant; temperatures in Rungwe-Livingstone drop to at least -3°C.


Description. A primarily brown, medium-sized, long-tailed, arboreal monkey. Muzzle elongated. Facial skin, including eyelids, black. Suborbital fossae "tear line" pronounced. Eyes brown. Pelage light to rufous brown except as follows: center of ventrum and distal half of tail, white to off-white; hands and feet, black; lower forelimbs, dark brown




Fig. 2. Holotype: adult male highland mangabey *Lophocebus kipunji* in the type locality, Rungwe-Livingstone, Tanzania. [Photograph by T.R.B. Davenport]

Holotype. Adult male in [REDACTED] (Fig. 2). Photograph taken in the type locality at $9^{\circ}07'S$ $33^{\circ}44'E$ (12). The number of individuals in each of the two populations of this species is undoubtedly very small; no live individual should be collected at this time to serve as the holotype. The Rungwe-Livingstone population is designated the source population for physical specimens in support of the holotype.

Lophocebus kipunji



Rungwecebus kipunji






Galagoides udzungwensis – komba

1996

Congosorex phillipsorum - bělozubka

2005





Rhynchocyon udzungwensis Rathbun 2008

Ital Rovero – 2005 fotopasti, Dr. Rathbun – Kalifornská akademie věd - 2008
Afrotheria – Macroscelidea – bércoun = elephant shrew, 1 kg, 30 cm






Dr. Rathbun







Národní přírodovědné muzeum Paříž

Genneta burloni 2003 – ženetka Bourlonova
coll. 1959 – podle kůže, stř. Z Afrika



...na trhu







Neofelis nebulosa diardi – nový druh, pův. poddruh levharta obláčkového
Borneo 2006, 1/3 terestrických savců na Borneu - endemiti









1995

...fotopasti



Nesolagus netscheri – králík krátkouchý (Leporidae),
Sumatra, od 1929 neznámý



1998: *Nesolagus netscheri*



Nesolagus netscheri



Nesolagus timminsi Averianov, Abramov, & Tikhonov,
2000 – králík Timminsův

Trh v Laosu 1995 - Timmins, 2000 - popsán Rusy, endemit Anamitského pohoří na hranici Laosu a Vietnamu



Nesolagus timminsi 2000



Laonastes aenigmamus 2005


- khanyou (Laonastidae = Diatomyidae), skalní krysa
(Laos=kámen, skála –řecky, tajuplná skalní myš z Laosu), objevena 1996.



Extinct relatives
Diatomys shantungensis fossil (20 mya)
from Shandong, China

Laonastes aenigmamus (extant)
Khammouan Province, Laos

Huchon et.al. 2007



Pectinator spekei – gundi somálský

Ctenodactylus gundi – gundi saharský (Ctenodactylidae) *Massoutiera mzabi* – g. východní



Nové čeledi na základě nálezů nových druhů

1905



Calomyscidae — křečci myší (také jako podčeleď – Calomyscinae)
– blízký a střední východ

1918



Lipotes vexillifer – delfínovec čínský (1918, ex. 2007 ?),

Baiji, vymřelá čeleď Lipotidae (Inniidae nebo Platanistidae) - delfínovcovití

1974



Craseonycteridae

- ...fotopasti



1998

Pseudoryx nghetinhensis



2007

Muntiacus vuquangensis

Bornejská kočkoliška (2003)



1. puchol hnědý *Diplogale hosei* (Viveridae) – netypické zbarvení
2. nový druh cibetky (2006)
3. poletucha Thomasova *Aeromys thomasi* (Sciuridae)
Meijaard et al., Mammal Review 2006




















Lepus thomasi vliegende eekhoorn
Evan Koortwijk '06



Aeromys thomasi



Sciurus thomasi vliegende eekhoorn
Evan Koortwijk '06





IUCN Small Carnivore Specialist Group - Mohd. Azlan J.

2003 *Diplogale hosei* – puchol hnědý (Borneo)



2006 *Bdeogale jacksoni* – mangusta Jacksonova (Herpestidae)



2007 *Dicerorhinus sumatrensis* – nosorožec sumaterský



2008 *Hexaprotodon liberiensis* – hrošík liberijský

© Ben Collen/ ZSL



Stealth Cam 02-11-2008 08:02:38

Hexaprotodon liberiensis




Hexaprotodon liberiensis



2003 *Catopuma badia* –
kočka bornejská



Thylacinus cynocephalus
1936 - ?






Figure 1 Species richness of mammals across Europe. This illustrative map was prepared using Worldmap software and shows species richness (number of species per grid square) with one level of smoothing. The entire atlas dataset has been used, including introduced species. There are some differences between the *Atlas Flora Europaea* UTM grid used by this software and the one used in the Atlas, particularly around the south-eastern borders of Europe and also in the treatment of some island groups.

