



Research centre
for toxic compounds
in the environment

Introduction to Ecotoxicology

Ludek Blaha + ecotox colleagues

cecoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE




OP Research and
Development for Innovation



Global anthropogenic threats ?

A safe operating space for humanity & the nine planetary boundaries

Rockstrom et al. 2009
(*Ecology and Society* 14(2): 32; *Nature* 461, 472-475)



Environmental pollution

Any examples ???

**CHEMICAL
ENTERS THE
ENVIRONMENT**



**LEVELS, FATE,
PROCESSES**




**Bioavailable
fraction**



"EXPOSURE"

acute

chronic



**CHEMICAL
ENTERS THE
ORGANISM**

biomonitoring




Toxicokinetics

*biotransformation
bioactivation
excretion / sequestration*

Target site

"EFFECT"




Assessment of chemical hazards

...to...

Humans
(**TOXICOLOGY**)



Other organisms
(**ECO**toxicology)



Chemicals in the environment


*Do you believe that **chemicals in products** sold to consumers have been proven **safe**?*

Think again

Most chemicals in modern use have simply not been tested for their impacts on human, even very basic effects.

... what about the effects in nature, then ?

Chemicals in the environment



- **Rats exposed in the womb to a single low dose of a widespread brominated flame retardant become hyperactive and have decreased sperm counts...**
- *Experiments with dioxin and similar compounds provide support for the assumption that cancer risks mediated by the aryl hydrocarbon receptor are additive. Previously untested for cancer, this assumption underpins a standard way of estimating exposure risks to these compounds. The results reinforce the need to focus health standards on mixtures rather than single compounds.*
- *At exposure levels within the range experienced by the general public, the phthalate **DBP** reduces expression of genes necessary for testosterone synthesis in fetal rats...*
- **Eutrophication of frog ponds** is linked to epidemics of frog deformities, because it creates conditions that lead to **higher rates of parasitic infections of tadpoles**. The parasitic infections in turn disrupt normal development of the tadpoles' limb buds during metamorphosis.

Published online: 21 October 2005; | doi:10.1038/news051017-16

Pollution makes for more girls

The stress of dirty air skews sex ratios in Sao Paulo.

[Erika Check](#)

Toxic fumes favour the fairer sex, a group of researchers in Brazil has found.

Jorge Hallak and his team at the University of Sao Paulo turned up the surprising result by studying babies born in their city. They divided the metropolis of 17 million people into areas of low, medium and high air pollution, using test results from air-quality monitoring stations. They then studied birth registries of children born from 2001 to 2003.

The team found that 48.3% of babies were female in the least polluted areas, but 49.3% were female in the dirtiest parts of town. After measuring the ratio of boys to girls born in all the areas, they calculated that 1,180 more babies would have been boys in the polluted areas if they had the same sex ratios as the cleaner areas. The team reported their findings on 17 October at the American



Babies born in highly polluted areas are more likely to be girls.

© Alamy

Major anthropogenic threats – example: waters

Direct



Indirect



Major impacts

- **Loss of biodiversity**



Changes in biodiversity

NATURE (2012) 482: 20



ATTACK OF THE BLOBS

Blooms of giant Nomura's jellyfish (*Nemopilema nomurai*) have troubled Japanese fishing crews.

increase in the global population of jellyfish — a catch-all term that covers some 2,000 species of true cnidarian jellyfish, ctenophores (or comb jellies) and other floating creatures called tunicates. But many marine biologists are now questioning the idea that jellyfish have started to overrun the oceans.

This week, a group of researchers published preliminary results from what will be the most comprehensive review of jellyfish population data¹. They say that there is not yet enough evi-

NYO/DU/NEWS.COM



Research centre
for toxic compounds
in the environment

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



OP Research and
Development for Innovation

Major impacts

Loss of biodiversity



• Impairment of ecosystem services

– Unbalanced water cycles

- Water scarcity
- Draughts/floods

– Impaired water quality

- Drinking waters
- Bathing waters
- Toxicants in food chain


– Shrinking of food supplies

- Direct → lowering fish amounts
- Indirect → crop yield



Impacts on fish → decreased crop yields

NATURE (2005) 437: 880



Research centre
for toxic compounds
in the environment

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



OP Research and
Development for Innovation

Impacts on biota → global effects

Mixing oceans

→ cooling the atmosphere

[Nature 447, p.522, May 31, 2007]



Marine life supplies up to 50% of the mechanical energy required worldwide to mix waters from the surface to deeper cool layers

[Dewar, Marine Res 64:541 (2006)]

[Katija a Dabiri, Nature 460:624 (2009)]



Research centre
for toxic compounds
in the environment

cetocoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



Ecotoxicology - from molecules to ecosystems ... and backwards







Figure 3.1 Biological levels of organization. The dimensions of time and space are less important for the investigation up to the levels of populations and biocoenoses.

1962



© Patuxent Wildlife Refuge, MA, USA

"DDT is good for me-e-e!"

The great expectations held for DDT have been realized. During 1946, exhaustive scientific tests have shown that, when properly used, DDT kills a host of destructive insect pests, and is a benefactor of all humanity.

Pennsalt produces DDT and its products in all standard forms and is now one of the country's largest producers of this amazing insecticide. Today, everyone can enjoy added comfort, health and safety through the insect-killing powers of Pennsalt DDT products . . . and DDT is only one of Pennsalt's many chemical products which benefit industry, farm and home.

GOOD FOR STEERS—Beef grows meatier nowadays . . . for it's a scientific fact that—compared to untreated cattle—beef steers gain up to 50 pounds extra when protected from horn flies and many other pests with DDT insecticides.

GOOD FOR THE HOME—helps to make healthier, more comfortable homes . . . protects your family from dangerous insect pests. Use Knox-Out DDT Powders and Sprays as directed . . . then watch the bugs "bite the dust"!

GOOD FOR DAIRIES—Up to 20% more milk . . . more butter . . . more cheese . . . tests prove greater milk production when dairy cows are protected from the annoyance of many insects with DDT insecticides like Knox-Out Stock and Barn Spray.

GOOD FOR FRUITS—Bigger apples, juicier fruits that are free from unsightly worms . . . all benefits resulting from DDT dusts and sprays.

GOOD FOR ROW CROPS—25 more barrels of potatoes per acre . . . actual DDT tests have shown crop increases like that! DDT dusts and sprays help truck farmers pass these gains along to you.

KNOX FOR INDUSTRY—Food processing plants, laundries, dry cleaning plants, hotels . . . dozens of industries gain effective bug control, more pleasant work conditions with Pennsalt DDT products.


PENN SALT
CHEMICALS
87 Years' Service to Industry • Farm • Home
PENNSYLVANIA SALT MANUFACTURING COMPANY
WIDENER BUILDING, PHILADELPHIA 7, PA.

Bitman et al. *Science* 1970, 168(3931): 594



Biochemistry


bird carbonate dehydratase



In vivo: shell thinning



In situ: bioaccumulation
-> bird population decline



ECOTOXICOLOGY by definition

- **Aim:** to maintain the natural structure and function of ecosystems
- **Definitions:**
 - ecotoxicology is concerned with the **toxic effects** of chemical and physical agents on living organisms, especially on populations and communities within defined ecosystems; **it includes the transfer pathways** and their interactions with the environment
 - science of contaminants in the biosphere and their effect on constituents of the biosphere, including humans' (Newman & Unger, 2002)
 - science that provides critical information on effects of toxic compounds on living organisms which SERVE various practical aims (environmental protection)

Ecotoxic effects





Figure 1 The effective concentration of a pollutant in an organism (e.g. fish, daphnia, algae) or at the target site inside the organism is the link between the environmental fate of a pollutant and its toxic effect.

Escher, B. I., Behra, R., Eggen, R. I. L., Fent, K. (1997), "Molecular mechanisms in ecotoxicology: an interplay between environmental chemistry and biology", *Chimia*, **51**, 915-921.

1) From molecules to individuals


MECHANISMS OF TOXICITY



2) From molecules to individuals - AOPs

ADVERSE OUTCOME PATHWAYS

Mechanistic effect models for ecotoxicology




→ Arrows indicate a causal relationship

See also: Ashauer & Escher *JEM* (2010), Rubach *et al. IEAM* (2011), Jager *et al. ES&T* (2011), Ashauer *et al. ET&C* (2011)

AOP Example: ethinylestradiol


Ethinylestradiol



**Binds to
ESTROGEN
RECEPTOR**

Target genes

- Proliferation/Apoptosis (sexual organs)
- Synthesis of egg yolk (fish, amphibia)





Effects

- Females: reproduction regulation
- Males: feminization
(+ e.g. cancer promotion, development, immunomodulation)

Kidd, K.A. et al. 2007. **Collapse of a fish population** following exposure to **a synthetic estrogen**. *Proceedings of the National Academy of Sciences* 104(21):8897-8901




5 ng/L (!)
7 years



Controls

+ Ethinylestradiol



Effects at different levels


- **Molecular**

- **Nonspecific effects**

- **Hydrophobic interactions** with phospholipid membranes (baseline = narcotic toxicity)
- Direct **reactivity**: electrophilic compounds → nucleophilic organism (e.g. oxidation of PROTEINS, lipids (membranes), DNA ...)

- **Specific effects**

- Activation of ER, AR and other „**nuclear receptors**“
- Inhibition of **enzymes** (e.g. CN- inhibits hemes in mitochondria/hemoglobin, insecticides ...)



Effects at different levels

- **Cellular**

- Effects on structure
- Effects on metabolism (maintenance)
- Effects on regulation


→ Changes in functions (e.g. Ethinylestradiol)

→ Repair, survival, growth

→ Death (apoptosis or necrosis)

→ Proliferation

→ Differentiation



Energy
hv
food



Losses
heat
faeces



Life
(maintenance)



Metabolism



Control,
Interactions
with environment



Defence
against pathogens
predators ...



Defence against
toxicants



**Chemical
stress**

Growth
to sexual
maturity



Reproduction



Research centre
for toxic compounds
in the environment

cercoen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

OP Research and
Development for Innovation

Chemical stress

→ energy re-allocation
→ „insufficient“ resources elsewhere

Energy
hv
food



Losses
heat
faeces



Life
(maintenance)



Metabolism



Control,
Interactions
with environment



Defence
against pathogens
predators ...



Defence against
toxicants



Chemical
stress

Growth
to sexual
maturity



Reproduction



Research centre
for toxic compounds
in the environment

ce/coen



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

OP Research and
Development for Innovation

Chemical stress

**+ ... another stress
(food scarcity)**

Energy
hv
food



Losses
heat
faeces



**Life
(maintenance)**



Metabolism



Control,
Interactions
with environment



Defence
against pathogens
predators ...



Defence against
toxicants



**Growth
to sexual
maturity**



Reproduction



**Chemical
stress**

Effects at different levels

- **Organism**

- Effects on structure
- Effects on metabolism (maintenance)
- Effects on regulation


→ Changes in functions (e.g. Ethinylestradiol)

→ Repair, survival, **growth**

→ **Death**

→ Proliferation = **Reproduction**

3 key apical endpoints
(reflected e.g. in regulations)



Effects at different levels

- **Population**

(... all the organisms that both belong to the same group or species (i.e. can sexually reproduce) and live in the same time within the same geographical area)

- Effects on **structure**
 - elderly vs. young, males vs. females
- Effects on **maintenance & growth**
 - Natality, mortality, reproduction fitness



Effects at different levels

- **Community & Ecosystem**

(... a group of interacting living organisms sharing a populated environment)

- Effects on **structure**
 - Loss of species, loss of biodiversity
- Effects on **functioning**
 - (including „ecosystem functions“)




Figure #31: Simplified Food Web (Source Down) similar to warm water lower end of river before entry into Mississippi River System or impoundment. The Flathead acts as a super predator when present as large specimens, and many predators such as walleyes and Gars compete for minnows and shad. Channel Catfish also appear and prey upon mussels and other invertebrates.

(Eco)toxicology – science of „doses“

Paracelsus (1493 - 1541)



*‘What is there which
is not a poison?’*

„Cause-effect paradigm“


- *All things are poison and nothing without poison.*
- *Solely **the dose determines** that a thing is not a **poison**.*

Ecotoxicology – ultimate goal ?

To identify (or predict)
safe vs hazardous
levels



Ecotoxicology: problems and approaches




Testing ecotoxicity – basics

Bioassays

- single / multiple species
- acute / chronic effects
- standardized (practical)
vs. experimental (research)

Simulation of the ecosystem



- major **trophic levels**
 - producers
 - consumers
 - decomposers



centre
compounds

In the environment

Ecotoxicology methods 1. - standardized assays




Cu addition



Concentration:

0.0 $\mu\text{g/L}$ 13 $\mu\text{g/L}$ 25 $\mu\text{g/L}$ 50 $\mu\text{g/L}$ 100 $\mu\text{g/L}$ 200 $\mu\text{g/L}$



Control 1 2 3 4 5


96-hour LC50 = 50 $\mu\text{g/L}$

Effect concentrations expressed in total/dissolved Cu




??? Safe concentrations ???

Laboratory ecotoxicology – data and results




Threshold:
**No Observed Effect
Concentration (NOEC)**



Ecotoxicology – methods 2. Micro & Mesocosms

Expensive & time consuming (e.g. *Pesticide testing*)
Variable results (natural variability ...)
Higher ecological relevancy



Research centre
for toxic compounds
in the environment

cetocoen






Fig. 5.2 Components of a standardized aquatic microcosm.

Ecotoxicology – methods 3. Field assessment / biomonitoring

- complex issue (geology, climate, chemistry, biology ..)
- Ecotoxicology mixes with Ecology
- comparing „contaminated“ with „control“ sites



Notes on practical testing

- Testing chemicals
 - Traditional / bioassays developed to assess **individual chemicals**
 - **Advantages**: Standardized approaches
 - **Disadvantage**: Limited ecological relevance
 - often **acute** tests only
 - „too standardized...“ (? Less representative ?)
 - does not assess/consider bioavailability
 - no consideration of mixture effects
 - no consideration of specific modes of action
 - no consideration of ecological situation
- Example: Acute (96h) fish toxicity assay with ethanol
 - No deaths (but fish are passive – slow swimming) → OK ?
 - Real life: easy prey → population decline


Notes on practical testing

- Testing toxicity of natural **contaminated matrices**
 - Rather new in ecotoxicology – many open challenges
 - Whole effluent toxicity testing (WET)
 - Contact soil toxicity assays
 - More complex and more complicated
 - „cause-effects“ often not clear
 - Natural variability in matrices
 - Algal tests - nutrients (Nitrogen, Phosphorus) >> Toxic compounds

Ecotoxicology in current practice

- Most legislations on chemicals) (e.g. REACH, Pharmaceuticals, Pesticides) have very simple (basic) requirements
 - EC50 from acute toxicity
 - Of 3 basic assays
 - Algae
 - Daphnia
 - Fish

Ecotox database:
www.epa.gov/ecotox



Ecotoxicology in current practice

- How to extrapolate 3 (or few more) EC50 values to get legally binding safe concentration, which is protecting virtually all organisms?




PNEC

(Predicted No Effect Concentration)
“value recommended by scientists”



EQS

(Environmental Quality Standard)
“value that occurs in legislation”






Cause – effect → Risk assessment

Exposure

(resulting from load)




Predicted Environmental
Concentration (**PEC**)




Effects

(what exposures cause effects)



Laboratory (and field) studies
Ecotoxicity tests

effective concentrations
(**PNEC**)



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



Extrapolation approaches

Ecotoxicological data

Assessment / Extrapolation factors

Data	Assessment factor
L(E)C50 short-term toxicity tests	1000
NOEC for 1 long-term toxicity test	100
NOEC for additional long-term toxicity tests of 2 trophic levels	50
NOEC for additional long-term toxicity tests of 3 species of 3 trophic levels	10

PNEC


Extrapolation approaches

Ecotoxicological data

Assessment / Extrapolation factors


Data	Assessment factor
L(E)C50 short-term toxicity tests	1000
NOEC for 1 long-term toxicity test	100
NOEC for additional long-term toxicity tests of 2 trophic levels	50
NOEC for additional long-term toxicity tests of 3 species of 3 trophic levels	10

Species sensitivity distribution (SSD)



PNEC

Risk assessment & management




Results of ecotoxicology

WHAT IS IT GOOD FOR ?

SOLVING PRACTICAL PROBLEMS

EQS in reality – example EU Water Framework Directive



List of priority compounds EU WFD (selection/examples)

Most recent (2015)

44 priority compounds (table here)
+ additional “watch list” → see further

AA: annual average;

MAC: maximum allowable concentration.

Unit: [µg/l]

(1)	(2)	(3)	(4)	(5)	(6)	(7)
No	Name of substance	CAS number ⁽¹⁾	AA-EQS ⁽²⁾ Inland surface waters ⁽³⁾	AA-EQS ⁽²⁾ Other surface waters	MAC-EQS ⁽⁴⁾ Inland surface waters ⁽³⁾	MAC-EQS ⁽⁴⁾ Other surface waters
(1)	Alachlor	15972-60-8	0,3	0,3	0,7	0,7
(2)	Anthracene	120-12-7	0,1	0,1	0,4	0,4
(3)	Atrazine	1912-24-9	0,6	0,6	2,0	2,0
(4)	Benzene	71-43-2	10	8	50	50
(5)	Brominated diphenylether ⁽⁵⁾	32534-81-9	0,0005	0,0002	not applicable	not applicable
(6)	Cadmium and its compounds (depending on water hardness classes) ⁽⁶⁾	7440-43-9	≤ 0,08 (Class 1) 0,08 (Class 2) 0,09 (Class 3) 0,15 (Class 4) 0,25 (Class 5)	0,2	≤ 0,45 (Class 1) 0,45 (Class 2) 0,6 (Class 3) 0,9 (Class 4) 1,5 (Class 5)	≤ 0,45 (Class 1) 0,45 (Class 2) 0,6 (Class 3) 0,9 (Class 4) 1,5 (Class 5)
(6a)	Carbon-tetrachloride ⁽⁷⁾	56-23-5	12	12	not applicable	not applicable

Watch list of substances for Union-wide monitoring as set out in Article 8b of Directive 2008/105/EC

Name of substance/group of substances	CAS number ⁽¹⁾	EU number ⁽²⁾	Indicative analytical method ⁽³⁾ ⁽⁴⁾ ⁽⁵⁾	Maximum acceptable method detection limit (ng/l)
17-Alpha-ethinylestradiol (EE2)	57-63-6	200-342-2	Large-volume SPE — LC-MS-MS	0,035
17-Beta-estradiol (E2), Estrone (E1)	50-28-2, 53-16-7	200-023-8	SPE — LC-MS-MS	0,4
Diclofenac	15307-86-5	239-348-5	SPE — LC-MS-MS	10
2,6-Ditert-butyl-4-methylphenol	128-37-0	204-881-4	SPE — GC-MS	3 160
2-Ethylhexyl 4-methoxycinnamate	5466-77-3	226-775-7	SPE — LC-MS-MS or GC-MS	6 000
Macrolide antibiotics ⁽⁶⁾			SPE — LC-MS-MS	90
Methiocarb	2032-65-7	217-991-2	SPE — LC-MS-MS or GC-MS	10
Neonicotinoids ⁽⁷⁾			SPE — LC-MS-MS	9
Oxadiazon	19666-30-9	243-215-7	LLE/SPE — GC-MS	88
Tri-allate	2303-17-5	218-962-7	LLE/SPE — GC-MS or LC-MS-MS	670

Another example where
ecotoxicology results are used

European strategy how to deal with chemicals



- **± 40 Directives** or Regulations concerning the evaluation and management of the dangers/risks associated with chemical substances
 - Regulation EEC 793/93 – **Existing substances**
 - Dir. 67/548/EEC – **New substances**
 - Dir. 98/8/EC – Biocides / Plant Protection Products
 - Further Directives – E.R.A. of new pharmaceuticals

Existing substances

- > 95,000,000 known chemicals
(...and counting <http://www.cas.org/>)
- 100,000 substances in EINECS (i.e. commercial use)
- 2747 HPVCs (High Production Volume Chemicals)
 - 14% minimum data-set (base-set)
 - 65% less than base-set
 - 21% no toxicity data
- Various priority lists
 - Aquatic hazard (EU Water framework directive)
 - Endocrine disruptors
 -

REACH

Registration, Evaluation and Authorisation of Chemicals

- 27-2-2001: White Paper on the Strategy for Future Chemicals Policy
- 23-10-2003: Commission's proposal REACH
- December 2008: Pre-registration mandatory (all chemicals in EU must be registered at ECHA)

 An agency of the European Union [Document library](#) | [News and Ever](#)

ECHA
EUROPEAN CHEMICALS AGENCY

[About Us](#) | [Regulations](#) | [Addressing Chemicals of Concern](#) | [Information on Chemicals](#)


ECHA > Homepage

European Chemicals
Agency
(<http://echa.europa.eu>)

15/06/2015 - Press release


Two new substances of very high concern (SVHCs) added to the Candidate List

ECHA took the decision to include two substances on the Candidate List based on proposals by Sweden and the Netherlands respectively, following the SVHC identification process with involvement of the Member State Committee. The Candidate List now contains 163 substances. Of those, 31 have subsequently been included in the Authorisation List.





REACH legislation in EU

Registration, Evaluation and Authorisation and Restriction of Chemicals



REACH





- **Major goals**

- Protection of man and the environment
- Increase competitiveness of EU chemical industry
- Increase transparency
- Avoid fragmentation of market
- Integration with international policies
- Reduction use of test animals

- **Approach**

- Industry is responsible – provides data

- **30000 existing substances**

- 0-3 year (2010): all HPVC and CMR substances (~ 3000)
- 4-6 year (2013): all 100-1000 t/y substances
- 7-11 year (2018'): all 10-100 and 1-10 t/y substances



- **Physico-chemical properties, e.g.:**
 - Vapour pressure, boiling point, Kow,...
- **Human toxicology, e.g.:**
 - Acute and chronic toxicity, skin irritation, carcinogenicity,...
- **Environment/ **Ecotoxicological** information, e.g.:**
 - Acute and/or chronic toxicity for aquatic organisms, biodegradation, ...

REACH: testing



Classification categories	Test requirements in REACH			
	>1t		>10t	>100t
		New or prioritised substance		
Reproductive toxicity (a generation test)	no	no	no	no
Chronic toxicity and cancer	no	no	no	(yes)
90-day study	no	no	no	(yes)
28-day study	no	no	(yes)	yes
Acute toxicity (a second route of exposure)	no	no	yes	yes
Acute toxicity	no	yes	yes	yes
Skin allergy	no	yes	yes	yes
Skin and eye irritation	no	yes	yes	yes
Mutagenicity (in vitro)	no	yes	yes	yes
Further ecotoxicity studies (incl long term tests)	no	no	no	yes
Acute toxicity: fish	no	no	yes	yes
Acute toxicity: algae	no	yes	yes	yes
Acute toxicity: Daphnia	no	yes	yes	yes
Biotic degradation	no	yes	yes	yes

REACH: implications




- Total: 2,8 to 5,6 billion €
- Industry pays
- Test costs (50-60% of total cost):
 - 86% for HH tests
 - 14% for environment tests
 - 0% for analyses
- Manpower and expertise?
 - Tests
 - Risk assessments
 - Evaluations
- Financial and time pressure:
danger for 'hazard-based' instead of 'risk-based' approach

Risks of chemicals: a balancing act


between perception, uncertainties,
science and pragmatism?

Final considerations

Effects of chemicals



Risks vs. Benefits





- All scientific questions answered ..., and
- We have standard tools and good legislations ...

Do we still need ecotoxicologists?

Problems of ecotoxicology – pharmaceuticals

Unexpected effects at “non-target” organisms

- **nephrotoxicity** at vultures
- relevant also in EU (ESP, EL, CY)



Problems of ecotoxicology – pharmaceuticals


Avermectin antiparasitics

Moxidectin –
small animals: spot-on



Ivermectin – antiparasitics for large herds

- Used **2-times per season** per sheep/cow
- **Kills 100% parasites** in sheep
- Released in dung - **kills 80-90% larvae of dung flies**
- High concentrations in dung (released 2 days post application)
- **Persistent in the soil** (half-life 30 days)
- Can be washed into adjacent streams (highly toxic to water insects)



ECONOMIC importance of BATS ...

Science AAAS.ORG | FEEDBACK | HELP | LIBRARIANS All Science Jour
MASARYK I
AAAS NEWS SCIENCE JOURNALS CAREERS MULTIMEDIA COLLECTION
Science The World's Leading Journal of Original Scientific Research, Global News, and Comme
Science Home Current Issue Previous Issues Science Express Science Products My Science Abou

<http://www.sciencemag.org/content/332/6025/41.figures-only>

Home > Science Magazine > 1 April 2011 > Boyles et al., 332 (6025): 41-42

- Article Views
- Summary
 - Full Text
 - Full Text (PDF)
 - Figures Only
 - Supporting Online Material



Science 1 April 2011:
Vol. 332 no. 6025 pp. 41-42
DOI: 10.1126/science.1201366

POLICY FORUM

CONSERVATION

Economic Importance of Bats in Agriculture

Justin G. Boyles^{1,*}, Paul M. Cryan², Gary F. McCracken³, Thomas H. Kunz⁴



Society is a balancing act ...



Scientists

Closing remarks



- Ecotoxicology is exciting **science!**
- **Interface:** science and society
- Many **opportunities**
- Sometimes **hard work**
10% inspiration and 90% „perspiration“

- Be **creative:** move frontiers
- **Keep the purpose** in mind
- **Be critical:** do not accept perceptions as facts
- **Speak up:** you have something to say!

Introduction to ecotoxicology

Ludek Blaha

blaha@recetox.muni.cz

<http://www.recetox.cz>