



Research centre
for toxic compounds
in the environment

Ecotoxicological bioassays



Jakub Hofman & Klára Hilscherová



Research centre
for toxic compounds
in the environment

Introduction

Protection of environment / nature

- Is and must be primary aim of **sustainably developing society**
- why? – discussion (ecosystem services, human, ethics ...)

How to protect ?


- Policy
- Legislation
- Research
- Education

Ecotoxicology – offers knowledge and tools useful for the effective and reasonable environmental protection

tools = ecotoxicological bioassays




Ecotoxicology



Discipline on the border of ecology and toxicology studying and evaluating direct and indirect effects of man-made or natural harmful chemicals or other stressors on animals (except human), plants and microorganisms at all levels of biological organization

Ecotoxicity bioassay, ecotoxicity test

- **a tool (method, procedure ...)** for ecotoxicological research and praxis – for environmental legislation and protection
- biota (tissue, organism, population, ecosystem ...) is exposed to chemicals (and/or other factors), in the lab (controlled conditions) or in the field (less controlled) and effects are evaluated and related to exposure
- **WHY?** To understand the cause-effects relationships (causality, dose-response ...)




Why bioassays?

Chemical analyses only are not able to identify risks properly because:

- 1) Real exposure is different – **bioavailability** in particular situation
- 2) Pollutant **mixture** - always in real ecosystems
- 3) **Matrix** itself has effects or interacts with effects of contaminants
- 4) Analytical methods are **limited** vs. wide spectrum of possibly toxic chemicals


Bioassays useful for:

- prospective ecological risk assessment
 - using bioassays for chemical compounds, pesticides
 - using bioassays for materials, mixtures
 - before they enter the environment



Bioassays useful for:

- retrospective ecological risk assessment
 - using bioassays for real environmental samples
 - searching the causalities between pollution and effects



Bioassays useful for:

Time: NOW !

RISK ASSESSMENT

PROSPECTIVE

RISK ASSESSMENT

RETROSPECTIVE

DISASTERS

PREDICTIONS for future

Bioassessment
Field assessment
Monitoring

Bioassessment
Field assessment
Monitoring

MIXTURES OF CHEMICALS
CONTAMINATED ENVIRONMENT

Lab studies




Lab studies

MIXTURES OF
TOXICANTS

INDIVIDUAL
TOXICANTS

Simulated small
ecosystems

Bioassays useful for:




Bioassay development

- **old bioassays** – acute, ecologically irrelevant, testing pure chemicals, pesticides
- **new bioassays** – sublethal endpoints, ecological relevancy, chemical mixtures, miniaturization, simple to measure endpoints




Differentiation of bioassays

- major trophic levels
 - producers
 - consumers
 - destruents
- aquatic / soil
- single / multiple species
- acute / chronic effects
- contact bioassays / eluate bioassays / TIE
- legislative / standardized (practical) / experimental (research)
- toxicity / bioaccumulation / biodegradation tests




Differentiation of bioassays




size, complexity
 ecological relevancy
 controlling the conditions
 old / fresh contamination
 no. of species

...




Differentiation of bioassays



Differentiation of bioassays

- limit test / comparison test
- concentration – response tests – preliminary, final




Use of bioassays

- Testing toxicity of natural **contaminated matrices**
 - Rather new in ecotoxicology – many open challenges
 - Whole effluent toxicity testing (WET)
 - Contact soil toxicity assays
 - More complex and more complicated
 - „cause-effects“ often not clear
 - Natural variability in matrices
 - Algal tests - nutrients (Nitrogen, Phosphorus) >> Toxic compounds



eluate 1:10



Control
soil

Mixture at
different
ratios



Aquatic organisms

| | |
|---|--------------|
| Test No. 201: Alga, Growth Inhibition Test | 11 July 2006 |
| Test No. 221: Lemna sp. Growth Inhibition Test | 11 July 2006 |
| Test No. 202: Daphnia sp. Acute Immobilisation Test | 23 Nov 2004 |
| Test No. 211: Daphnia magna Reproduction Test | 16 Oct 2008 |
| Test No. 203: Fish, Acute Toxicity Test | 17 July 1992 |
| Test No. 204: Fish, Prolonged Toxicity Test: 14-Day Study | 04 Apr 1984 |
| Test No. 210: Fish, Early-Life Stage Toxicity Test | 17 July 1992 |
| Test No. 212: Fish, Short-term Toxicity Test on Embryo and Sac-Fry Stages | 21 Sep 1998 |
| Test No. 215: Fish, Juvenile Growth Test | 21 Jan 2000 |
| Test No. 229: Fish Short Term Reproduction Assay | 08 Sep 2009 |
| Test No. 230: 21-day Fish Assay | 08 Sep 2009 |
| Test No. 231: Amphibian Metamorphosis Assay | 08 Sep 2009 |

Sediment organisms

| | |
|--|--------------|
| Test No. 218: Sediment-Water Chironomid Toxicity Using Spiked Sediment | 23 Nov 2004 |
| Test No. 219: Sediment-Water Chironomid Toxicity Using Spiked Water | 23 Nov 2004 |
| Test No. 233: Sediment-Water Chironomid Life-Cycle Toxicity Test Using Spiked Water or Spiked Sediment | 23 July 2010 |
| Test No. 225: Sediment-Water Lumbriculus Toxicity Test Using Spiked Sediment | 15 Oct 2007 |

Soil organisms


| | |
|--|-------------|
| Test No. 208: Terrestrial Plant Test: Seedling Emergence and Seedling Growth Test | 17 Aug 2006 |
| Test No. 227: Terrestrial Plant Test: Vegetative Vigour Test | 17 Aug 2006 |
| Test No. 207: Earthworm, Acute Toxicity Tests | 04 Apr 1984 |
| Test No. 220: Enchytraeid Reproduction Test | 23 Nov 2004 |
| Test No. 222: Earthworm Reproduction Test (<i>Eisenia fetida</i> / <i>Eisenia andrei</i>) | 23 Nov 2004 |
| Test No. 228: Determination of Developmental Toxicity of a Test Chemical to Dipteran Dung Flies(<i>Scathophaga stercoraria</i> L. (<i>Scathophagidae</i>), <i>Musca autumnalis</i> De Geer (<i>Muscidae</i>)) | 16 Oct 2008 |
| Test No. 232: Collembolan Reproduction Test in Soil | 08 Sep 2009 |
| Test No. 226: Predatory mite (<i>Hypoaspis</i> (<i>Geolaelaps</i>) <i>aculeifer</i>) reproduction test in soil | 16 Oct 2008 |
| Test No. 216: Soil Microorganisms: Nitrogen Transformation Test | 21 Jan 2000 |
| Test No. 217: Soil Microorganisms: Carbon Transformation Test | 21 Jan 2000 |

Other tests

| | |
|--|--------------|
| Test No. 213: Honeybees, Acute Oral Toxicity Test | 21 Sep 1998 |
| Test No. 214: Honeybees, Acute Contact Toxicity Test | 21 Sep 1998 |
| Test No. 205: Avian Dietary Toxicity Test | 04 Apr 1984 |
| Test No. 206: Avian Reproduction Test | 04 Apr 1984 |
| Test No. 223: Avian Acute Oral Toxicity Test | 23 July 2010 |

Aquatic microorganisms

| | |
|------------------|---|
| ISO 10712:1995 | Water quality -- <u>Pseudomonas putida growth inhibition test</u> (Pseudomonas cell multiplication inhibition test) |
| ISO 11348-1:2007 | Water quality -- Determination of the inhibitory effect of water samples on the <u>light emission of Vibrio fischeri</u> (Luminescent bacteria test) -- Part 1: Method using freshly prepared bacteria |
| ISO 11348-2:2007 | Water quality -- Determination of the inhibitory effect of water samples on the light emission of Vibrio fischeri (Luminescent bacteria test) -- Part 2: Method using liquid-dried bacteria |
| ISO 11348-3:2007 | Water quality -- Determination of the inhibitory effect of water samples on the light emission of Vibrio fischeri (Luminescent bacteria test) -- Part 3: Method using freeze-dried bacteria |
| ISO 13641-1:2003 | Water quality -- Determination of inhibition of gas production of anaerobic bacteria -- Part 1: General test |
| ISO 13641-2:2003 | Water quality -- Determination of inhibition of gas production of anaerobic bacteria -- Part 2: Test for low biomass concentrations |
| ISO 13829:2000 | Water quality -- Determination of the genotoxicity of water and waste water using the <u>umu-test</u> |
| ISO 16240:2005 | Water quality -- Determination of the genotoxicity of water and waste water -- Salmonella/microsome test (<u>Ames test</u>) |
| ISO/DIS 11350 | Water quality -- Determination of the genotoxicity of water and waste water -- Salmonella/microsome fluctuation test (Ames fluctuation test) |
| ISO 15522:1999 | Water quality -- Determination of the inhibitory effect of water constituents on the growth of activated sludge microorganisms |
| ISO 21338:2010 | Water quality -- Kinetic determination of the inhibitory effects of sediment, other solids and coloured samples on the light emission of Vibrio fischeri (<u>kinetic luminescent bacteria test</u>) |
| ISO 8192:2007 | Water quality -- Test for inhibition of oxygen consumption by activated sludge for carbonaceous and ammonium oxidation |
| ISO 9509:2006 | Water quality -- Toxicity test for assessing the inhibition of nitrification of activated sludge microorganisms |



Aquatic plants

| | |
|-------------------|--|
| ISO 20079:2005 | Water quality -- Determination of the toxic effect of water constituents and waste water on duckweed (<i>Lemna minor</i>) -- Duckweed growth inhibition test |
| ISO 8692:2004 | Water quality -- Freshwater algal growth inhibition test with unicellular green algae |
| ISO/CD 16191 | Water quality - Determination of the toxic effect of sediment and soil on the growth behaviour of <i>Myriophyllum aquaticum</i> - Myriophyllum test |
| ISO 10253:2006 | Water quality -- Marine algal growth inhibition test with <i>Skeletonema costatum</i> and <i>Phaeodactylum tricoratum</i> |
| ISO 10710:2010 | Water quality -- Growth inhibition test with the marine and brackish water macroalga <i>Ceramium tenuicorne</i> |
| ISO 14442:2006 | Water quality -- Guidelines for algal growth inhibition tests with poorly soluble materials, volatile compounds, metals and waste water |
| ISO/DIS 13308 | Water quality -- Toxicity test based on reproduction inhibition of the green macroalga <i>Ulva pertusa</i> |
| ISO/TR 11044:2008 | Water quality -- Scientific and technical aspects of batch algae growth inhibition tests |



Aquatic invertebrates

| | |
|-----------------|--|
| ISO 6341:1996 | Water quality -- Determination of the inhibition of the mobility of <u>Daphnia magna</u> Straus (Cladocera, Crustacea) -- <u>Acute toxicity test</u> |
| ISO 10706:2000 | Water quality -- Determination of <u>long term toxicity of substances to Daphnia magna</u> Straus (Cladocera, Crustacea) |
| ISO/DIS 14380 | Water quality -- Determination of the <u>acute toxicity to Thamnocephalus platyurus</u> (Crustacea, Anostraca) |
| ISO/CD 16303 | Water quality -- Determination of toxicity of <u>fresh water sediments using Hyalella azteca</u> |
| ISO 10872:2010 | Water quality -- Determination of the toxic effect of sediment and soil samples on growth, fertility and <u>reproduction of Caenorhabditis elegans</u> (Nematoda) |
| ISO 16712:2005 | Water quality -- Determination of acute toxicity of marine or estuarine sediment to amphipods |
| ISO 20665:2008 | Water quality -- Determination of chronic toxicity to Ceriodaphnia dubia |
| ISO 20666:2008 | Water quality -- Determination of the chronic toxicity to Brachionus calyciflorus in 48 h |
| ISO 14669:1999 | Water quality -- Determination of acute lethal toxicity to marine copepods (Copepoda, Crustacea) |
| ISO/DIS 14371 | Water quality -- Determination of freshwater-sediment subchronic toxicity to Heterocypris incongruens (Crustacea, Ostracoda) |
| ISO 7828:1985 | Water quality -- Methods of biological sampling -- Guidance on handnet sampling of aquatic benthic macro-invertebrates |
| ISO 8265:1988 | Water quality -- Design and use of quantitative samplers for benthic macro-invertebrates on stony substrata in shallow freshwaters |
| ISO 8689-1:2000 | Water quality -- Biological classification of rivers -- Part 1: Guidance on the interpretation of biological quality data from surveys of benthic macroinvertebrates |
| ISO 8689-2:2000 | Water quality -- Biological classification of rivers -- Part 2: Guidance on the presentation of biological quality data from surveys of benthic macroinvertebrates |
| ISO/DIS 10870 | Water quality -- Guidelines for the selection of sampling methods and devices for benthic macroinvertebrates in fresh waters |
| ISO/WD 16778 | Water quality -- Calanoid copepod development test with Acartia tonsa |



Aquatic vertebrates

| | |
|---------------------|---|
| ISO 15088:2007 | Water quality -- Determination of the acute toxicity of <u>waste water to zebrafish eggs</u> (Danio rerio) |
| ISO 7346-1:1996 | Water quality -- Determination of the <u>acute lethal toxicity of substances to a freshwater fish</u> [Brachydanio rerio Hamilton-Buchanan (Teleostei, Cyprinidae)] -- Part 1: Static method |
| ISO 7346-2:1996 | Water quality -- Determination of the acute lethal toxicity of substances to a freshwater fish [Brachydanio rerio Hamilton-Buchanan (Teleostei, Cyprinidae)] -- Part 2: Semi-static method |
| ISO 7346-3:1996 | Water quality -- Determination of the acute lethal toxicity of substances to a freshwater fish [Brachydanio rerio Hamilton-Buchanan (Teleostei, Cyprinidae)] -- Part 3: Flow-through method |
| ISO 10229:1994 | Water quality -- Determination of the prolonged toxicity of substances to freshwater fish -- Method for evaluating the effects of substances on the <u>growth rate of rainbow trout</u> (Oncorhynchus mykiss Walbaum (Teleostei, Salmonidae)) |
| ISO 12890:1999 | Water quality -- Determination of toxicity to embryos and larvae of freshwater fish -- Semi-static method |
| ISO 21427-1:2006 | Water quality -- Evaluation of genotoxicity by measurement of the induction of micronuclei -- Part 1: Evaluation of genotoxicity using amphibian larvae |
| ISO 21427-2:2006 | Water quality -- Evaluation of genotoxicity by measurement of the induction of micronuclei -- Part 2: Mixed population method using the cell line V79 |
| ISO 23893-1:2007 | Water quality -- Biochemical and physiological measurements on fish -- Part 1: Sampling of fish, handling and preservation of samples |
| ISO/TS 23893-2:2007 | Water quality -- Biochemical and physiological measurements on fish -- Part 2: Determination of ethoxyresorufin-O-deethylase (EROD) |
| ISO/CD 23893-3 | Water quality -- Biochemical and physiological measurements on fish -- Part 3: Determination of vitellogenin |





Soil microorganisms

| | |
|---------------------|--|
| ISO 10381-6:2009 | Soil quality -- Sampling -- Part 6: Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory |
| ISO 14240-1:1997 | Soil quality -- Determination of soil microbial biomass -- Part 1: Substrate-induced respiration method |
| ISO 14240-2:1997 | Soil quality -- Determination of soil microbial biomass -- Part 2: Fumigation-extraction method |
| ISO 16072:2002 | Soil quality -- Laboratory methods for determination of microbial soil respiration |
| ISO 17155:2002 | Soil quality -- Determination of abundance and activity of soil microflora using respiration curves |
| ISO 15685:2004 | Soil quality -- Determination of potential nitrification and inhibition of nitrification -- Rapid test by ammonium oxidation |
| ISO 14238:1997 | Soil quality -- Biological methods -- Determination of nitrogen mineralization and nitrification in soils and the influence of chemicals on these processes |
| ISO 23753-1:2005 | Soil quality -- Determination of dehydrogenase activity in soils -- Part 1: Method using triphenyltetrazolium chloride (TTC) |
| ISO 23753-2:2005 | Soil quality -- Determination of dehydrogenase activity in soils -- Part 2: Method using iodotetrazolium chloride (INT) |
| ISO/DIS 11063 | Soil quality -- Method to directly extract DNA from soil samples |
| ISO/TS 29843-1:2010 | Soil quality -- Determination of soil microbial diversity -- Part 1: Method by phospholipid fatty acid analysis (PLFA) and phospholipid ether lipids (PLEL) analysis |
| ISO/PRF TS 29843-2 | Soil quality -- Determination of soil microbial diversity -- Part 2: Method by phospholipid fatty acid analysis (PLFA) using the simple PLFA extraction method |
| ISO/TS 10832:2009 | Soil quality -- Effects of pollutants on mycorrhizal fungi -- Spore germination test |
| ISO/TS 22939:2010 | Soil quality -- Measurement of enzyme activity patterns in soil samples using fluorogenic substrates in micro-well plates |
| ISO 11266:1994 | Soil quality -- Guidance on laboratory testing for biodegradation of organic chemicals in soil under aerobic conditions |
| ISO 15473:2002 | Soil quality -- Guidance on laboratory testing for biodegradation of organic chemicals in soil under anaerobic conditions |
| ISO 14239:1997 | Soil quality -- Laboratory incubation systems for measuring the mineralization of organic chemicals in soil under aerobic conditions |

Soil invertebrates

| | |
|------------------|---|
| ISO 11268-1:1993 | Soil quality -- Effects of pollutants on earthworms (<i>Eisenia fetida</i>) -- Part 1: Determination of acute toxicity using artificial soil substrate |
| ISO 11268-2:1998 | Soil quality -- Effects of pollutants on earthworms (<i>Eisenia fetida</i>) -- Part 2: Determination of effects on reproduction |
| ISO 11268-3:1999 | Soil quality -- Effects of pollutants on earthworms -- Part 3: Guidance on the determination of effects in field situations |
| ISO 11267:1999 | Soil quality -- Inhibition of reproduction of <i>Collembola (Folsomia candida)</i> by soil pollutants |
| ISO 16387:2004 | Soil quality -- Effects of pollutants on Enchytraeidae (<i>Enchytraeus sp.</i>) -- Determination of effects on reproduction and survival |
| ISO 15952:2006 | Soil quality -- Effects of pollutants on juvenile land snails (Helicidae) -- Determination of the effects on growth by soil contamination |
| ISO 20963:2005 | Soil quality -- Effects of pollutants on insect larvae (<i>Oxythyrea funesta</i>) -- Determination of acute toxicity |
| ISO 17512-1:2008 | Soil quality -- Avoidance test for determining the quality of soils and effects of chemicals on behaviour -- Part 1: Test with earthworms (<i>Eisenia fetida</i> and <i>Eisenia andrei</i>) |
| ISO/DIS 17512-2 | Soil quality -- Avoidance test for determining the quality of soils and effects of chemicals on behaviour -- Part 2: Test with collembolans (<i>Folsomia candida</i>) |
| ISO 23611-1:2006 | Soil quality -- Sampling of soil invertebrates -- Part 1: Hand-sorting and formalin extraction of earthworms |
| ISO 23611-2:2006 | Soil quality -- Sampling of soil invertebrates -- Part 2: Sampling and extraction of micro-arthropods (<i>Collembola</i> and <i>Acarina</i>) |
| ISO 23611-3:2007 | Soil quality -- Sampling of soil invertebrates -- Part 3: Sampling and soil extraction of enchytraeids |
| ISO 23611-4:2007 | Soil quality -- Sampling of soil invertebrates -- Part 4: Sampling, extraction and identification of soil-inhabiting nematodes |
| ISO/DIS 23611-5 | Soil quality -- Sampling of soil invertebrates -- Part 5: Sampling and extraction of soil macro-invertebrates |
| ISO/DIS 23611-6 | Soil quality -- Sampling of soil invertebrates -- Part 6: Guidance for the design of sampling programmes with soil invertebrates |



Plants

| | |
|------------------|---|
| ISO 11269-1:1993 | Soil quality -- Determination of the effects of pollutants on soil flora -- Part 1: Method for the measurement of <u>inhibition of root growth</u> |
| ISO 11269-2:2005 | Soil quality -- Determination of the effects of pollutants on soil flora -- Part 2: Effects of chemicals on the emergence and growth of higher plants |
| ISO 17126:2005 | Soil quality -- Determination of the effects of pollutants on soil flora -- Screening test for emergence of lettuce seedlings (<i>Lactuca sativa</i> L.) |
| ISO 22030:2005 | Soil quality -- Biological methods -- <u>Chronic toxicity in higher plants</u> |
| ISO/CD 29200 | Soil quality -- Assessment of genotoxic effects on higher plants -- Micronucleus test on <i>Vicia faba</i> |



Aquatic

- [850.1010 - Aquatic Invertebrate Acute Toxicity, Test, Freshwater Daphnids \(PDF\) \(10 pp, 36K\)](#)
- [850.1020 - Gammarid Acute Toxicity Test \(PDF\) \(11 pp, 36K\)](#)
- [850.1025 - Oyster Acute Toxicity Test \(Shell Deposition\) \(PDF\) \(9 pp, 32K\)](#)
- [850.1035 - Mysid Acute Toxicity Test \(PDF\) \(10 pp, 34K\)](#)
- [850.1045 - Penaeid Acute Toxicity Test \(PDF\) \(9 pp, 32K\)](#)
- [850.1055 - Bivalve Acute Toxicity Test \(Embryo Larval\) \(PDF\) \(7 pp, 27K\)](#)
- [850.1075 - Fish Acute Toxicity Test, Freshwater And Marine \(PDF\) \(13 pp, 45K\)](#)
- [850.1085 - Fish Acute Toxicity Mitigated By Humic Acid \(PDF\) \(10 pp, 35K\)](#)
- [850.1300 - Daphnid Chronic Toxicity Test \(PDF\) \(12 pp, 42K\)](#)
- [850.1350 - Mysid Chronic Toxicity Test \(PDF\) \(10 pp, 36K\)](#)
- [850.1400 - Fish Early-Life Stage Toxicity Test \(PDF\) \(15 pp, 66K\)](#)
- [850.1500 - Fish Life Cycle Toxicity \(PDF\) \(4 pp, 16K\)](#)
- [850.1710 - Oyster BCF \(PDF\) \(14 pp, 50K\)](#)
- [850.1730 - Fish BCF \(PDF\) \(25 pp, 74K\)](#)
- [850.1735 - Whole Sediment Acute Toxicity Invertebrates, Freshwater \(PDF\) \(19 pp, 65K\)](#)
- [850.1740 - Whole Sediment Acute Toxicity Invertebrates, Marine \(PDF\) \(14 pp, 50K\)](#)
- [850.1790 - Chironomid Sediment Toxicity Test \(PDF\) \(16 pp, 57K\)](#)
- [850.1800 - Tadpole/Sediment Subchronic Toxicity Test \(PDF\) \(15 pp, 49K\)](#)
- [850.1850 - Aquatic Food Chain Transfer \(PDF\) \(4 pp, 16K\)](#)
- [850.1900 - Generic Freshwater Microcosm Test, Laboratory \(PDF\) \(28 pp, 76K\)](#)
- [850.1925 - Site-Specific Aquatic Microcosm Test, Laboratory \(PDF\) \(21 pp, 91K\)](#)
- [850.1950 - Field Testing For Aquatic Organisms \(PDF\) \(7 pp, 21K\)](#)
- [850.4400 - Aquatic Plant Toxicity Test Using Lemna Spp., Tiers I and II \(PDF\) \(10 pp, 36K\)](#)
- [850.4450 - Aquatic Plants Field Study, Tier III \(PDF\) \(9 pp, 30K\)](#)
- [850.5400 - Algal Toxicity, Tiers I and II \(PDF\) \(11 pp, 42K\)](#)
- [850.6800 - Modified Activated Sludge, Respiration Inhibition Test for Sparingly Soluble Chemicals \(PDF\) \(9 pp, 37K\)](#)



Soil

- [850.2450 - Terrestrial \(Soil-Core\) Microcosm Test \(PDF\) \(19 pp, 123K\)](#)
- [850.4000 - Background-Nontarget Plant Testing \(PDF\) \(15 pp, 50K\)](#)
- [850.4025 - Target Area Phytotoxicity \(PDF\) \(15 pp, 51K\)](#)
- [850.4100 - Terrestrial Plant Toxicity, Tier I \(Seedling Emergence\) \(PDF\) \(8 pp, 29K\)](#)
- [850.4150 - Terrestrial Plant Toxicity, Tier I \(Vegetative Vigor\) \(PDF\) \(8 pp, 28K\)](#)
- [850.4200 - Seed Germination/Root Elongation Toxicity Test \(PDF\) \(8 pp, 29K\)](#)
- [850.4225 - Seedling Emergence, Tier II \(PDF\) \(10 pp, 36K\)](#)
- [850.4230 - Early Seedling Growth Toxicity Test \(PDF\) \(9 pp, 33K\)](#)
- [850.4250 - Vegetative Vigor, Tier II \(PDF\) \(10 pp, 35K\)](#)
- [850.4300 - Terrestrial Plants Field Study, Tier III \(PDF\) \(8 pp, 27K\)](#)
- [850.4600 - Rhizobium-Legume Toxicity \(PDF\) \(14 pp, 73K\)](#)
- [850.4800 - Plant Uptake and Translocation Test \(PDF\) \(13 pp, 35K\)](#)
- [850.5100 - Soil Microbial Community Toxicity Test \(PDF\) \(11 pp, 46K\)](#)
- [850.6200 - Earthworm Subchronic Toxicity Test \(PDF\) \(13 pp, 43K\)](#)

Other

- [850.2100 - Avian Acute Oral Toxicity Test \(PDF\) \(11 pp, 38K\)](#)
- [850.2200 - Avian Dietary Toxicity Test \(PDF\) \(12 pp, 42K\)](#)
- [850.2300 - Avian Reproduction Test \(PDF\) \(16 pp, 53K\)](#)
- [850.2400 - Wild Mammal Acute Toxicity \(PDF\) \(5 pp, 18K\)](#)
- [850.2500 - Field Testing For Terrestrial Wildlife \(PDF\) \(43 pp, 115K\)](#)
- [850.3020 - Honey Bee Acute Contact Toxicity \(PDF\) \(8 pp, 27K\)](#)
- [850.3030 - Honey Bee Toxicity of Residues on Foliage \(PDF\) \(6 pp, 23K\)](#)
- [850.3040 - Field Testing for Pollinators \(PDF\) \(5 pp, 18K\)](#)



Testing strategy


- **Battery of assays**
 - Fast screening tests (Vibrio fisheri bioluminescence, 30 min toxicity)
 - Standardized acute toxicity tests
 - Further studies with chronic assays
 - **Combine trophic levels! Combine exposure routes!**
- **Various purposes -> guidelines and recommendations**
 - REACH (EU - Registration, Evaluation and Authorisation of Chemicals)
 - Plant protection products + biocides
 - Veterinary and human pharmaceuticals
 - Waste materials ...
- **The most common set ups**
 - algae / D. magna / fish for aquatic environment
 - earthworm (enchytraeid/springtail) / plant for soil environment

Testing strategy


ISO 15799 (2003): Guidance on the ecotoxicological characterization of soils and soil materials

ISO 17616 (2008): Guidance on the choice and evaluation of bioassays for ecotoxicological characterization of soils and soil materials

Retention function – Biotests with eluates



Habitat function - Biotests with solids



General scheme of bioassay

1) Prepare the organism

Culture media, standardized numbers, age, etc.

2) Prepare the sample

Dilution series

water/culture media – direct organism exposure

Include BLANK (medium only)

solvent for organic compounds – minimum to be added

Include SOLVENT CONTROL

3) Expose organisms


... for appropriate time, number of repetitions,
under specified conditions

4) Evaluate and report results

measure the endpoint / count organisms

validity criteria

statistical evaluation (means, ANOVA, dose-response ...)






Research centre
for toxic compounds
in the environment

Aquatic ecotoxicology bioassays

Algal growth inhibition test (ISO 8692)



Organisms

Freshwater species: *Selenastrum capricornutum*,
Scenedesmus subspicatus, *Chlorella vulgaris*,
Microcystis aeruginosa, *Anabaena flos-aquae*,
Navicula pelliculosa;

Saltwater species: *Skeletonema costatum*,
Thalassiosira pseudonana, and *Dunaliella tertiolecta*

Number of organisms
per chamber ($\pm 10\%$)

Selenastrum capricornutum and other freshwater green algae 2×10^4 cells/ml

Navicula pelliculosa 2×10^4 cells/ml

Microcystis aeruginosa 5×10^4 cells/ml

Anabaena flos-aquae 2×10^4 cells/ml

Saltwater species 2×10^4 cells/ml

Experimental design

Test vessel type and size

Test solution volume

Sterile Erlenmeyer flasks of borosilicate glass, any size
Not to exceed 50% of the flask volume for tests
conducted on a shaker, and not more than 20% of the
flask volume for tests not conducted on a shaker

Number of replicate chambers
per sample

2 or more

Test duration

96 h

Physical and chemical parameters

Water temperature

$24 \pm 2^\circ\text{C}$ for freshwater green and blue-green algae
 $20 \pm 2^\circ\text{C}$ for *Navicula pelliculosa* and other saltwater
algae

Light quality

Light intensity

Continuous "cool-white" fluorescent

Should not vary by more than $\pm 15\%$:

$60 \mu\text{E m}^{-2}/\text{s}^{-1}$ ($4300 \text{ lm}/\text{m}^2$) for freshwater diatoms and
green algae

$30 \mu\text{E m}^{-2}/\text{s}^{-1}$ ($2150 \text{ lm}/\text{m}^2$) for freshwater blue-green
algae

$82\text{--}90 \mu\text{E m}^{-2}/\text{s}^{-1}$ (5900 to $6500 \text{ lm}/\text{m}^2$) for *Thalassiosira*

$60 \mu\text{E m}^{-2}/\text{s}^{-1}$ ($4300 \text{ lm}/\text{m}^2$) for *Skeletonema*

14 h light/10 h dark for *Skeletonema*

Photoperiod

Test solution pH

7.5 ± 0.1 for freshwater


8.0 ± 0.1 for saltwater

Endpoint

Biomass, cell number, area underneath the growth curve

Algal growth inhibition test (ISO 8692)


- Miniaturization




- Alternative: Algaltokit

ALGALTOXKIT F™ MICROBIOTESTS

Cost-effective, culture/maintenance free* bioassays with the micro-algae *Selenastrum capricornutum* (renamed *Raphidocoelis subcapitata*/*Pseudokirchneriella subcapitata*)



Algal beads (2 mm)
> 1 million algal cells per bead



Algal cells

The micro-algae are included in the kits in “algal beads” from which they can be set free “on demand”

Each Algaltokit contains all the materials to perform two 72h growth inhibition tests



Duckweed bioassay (ISO 20079)

- *Lemna minor*
- 10 leaves per 1 beaker
- pH 6.5; 10 000 lx; 24°C
- 96 hours
- growth, biomass, no. of leaves
- image analysis possible
- validity:
 - 8x increase in control
 - IC₅₀ for K₂Cr₂O₇ 10-60 mg/L



Aquatic consumers - invertebrates


Daphnia magna



Artemia salina




Chironomus riparius



Potamopyrgus antipodarum



Ceriodaphnia dubia



Gammarus



Tamnocephalus platyurus



Lumbriculus variegatus



Hyalella azteca




Tubifex tubifex



Daphnia magna test (ISO 6341)

- 5 individuals per replicate (min 2 ml)
- no food
- 20°C; dark or 16h light / 8h dark
- 24h, 48h
- medium
 - $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 11,76 g/l
 - $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 4,93 g/l
 - NaHCO_3 2,59 g/l
 - KCl 0,23 g/l
 - 25 ml each to 1 L
 - pH 7.8
 - aeration






- no. of immobilized individuals
- validity:
 - $\text{O}_2 > 80 \%$ (2 mg/l)
 - mortality in control 10 %

Daphnia magna chronic test (ISO 10706)

- 10 juveniles (24h old) per replicate
- 50ml medium
- 3 times per week medium change
- 21 days

- 20 ± 2 °C
- pH 7-9
- dissolved O₂ > 3mg/l
- 16 h light / 8 h dark
- food - algae
- week controls: O₂, T, pH

Mortality, survival
Reproduction
Juveniles
Behavior




Daphnia magna acute vs chronic test

| Test type | Chronic (partial life cycle) | Acute 48 h |
|----------------------------------|--|---------------------------|
| Organisms | <i>D. magna</i> | <i>D. magna</i> |
| Age of test organisms | 24-h old | 24-h old |
| Number of organisms per chamber | 10 | 10 (minimum) |
| Experimental design | | |
| Test vessel type and size | 100 ml beakers | 250 ml |
| Test solution volume | 80 ml | 200 ml |
| Number of replicates per sample | 2 (minimum) | 3 (minimum) |
| Feeding regime | Various combinations of trout chow, yeast, alfalfa, green algae, and diatoms given in excess | Do not feed |
| Test duration | 21 days | 48 hr |
| Physical and chemical parameters | | |
| Water temperature | 20°C | 20 ± 2°C |
| Light quality | Ambient laboratory levels | Ambient laboratory levels |
| Light intensity | Up to 600 lux | 540 to 1080 lux |
| Photoperiod | 16 h light and 8 h dark (with 15- to 30-min transition) | 16 h light and 8 h dark |
| pH range | 7.0–8.6 | 7.0–8.6 |
| DO concentration | 40–100% | 60–100% |
| Aeration | Not necessary | none |
| Endpoint | Survival, growth, and reproduction | Immobilization |



Alternative: microbiotests



*Test organisms are included in the kits as "dormant eggs (cysts)" which can be hatched "on demand"


ROTOXKIT F chronic

Contains all the materials to perform three 48h reproduction assays



THAMNOTOXKIT F™ MICROBIOTESTS

With the crustacean *Thamnocephalus platyurus*




te
in
ia
la



OSTRACODTOXKIT F™ MICROBIOTESTS


FOR SEDIMENT TOXICITY TESTING

With the benthic crustacean *Heterocypris incongruens*



Chironomus riparius test (OECD 218)

- 10 larvae (cca 10d old) per beaker
- OECD sediment
- 100 ml sediment / 175 ml water
- $20 \pm 2^\circ\text{C}$; food, aeration
- 16h light / 8h dark; controlled pH, O_2
- 10 d
- survival and growth



A: the nylon screen
B: the inverted plastic cups
C: the lipless exposure beaker
D: the water exchange screen ports
E: water
F: sediment



Fish bioassays

Guppy, *Poecilia reticulata*



acute tests (96 h)
prolonged tests
embryolarval tests
chronic tests

- reproduction
- growth

Fathead minnow, *Pimephales promelas* (USA)



Specific endpoints –
genotoxicity, endocrine
disruption

Zebrafish, *Danio rerio* (syn. *Brachydanio rerio*)



(Rainbow) trout
(*Onchorhynchus sp.*)



Medaka, *Oryzias latipes*





Nile tilapia, *Oreochromis niloticus*




Fish bioassay acute (ISO 7346 1-3)

- *Brachydanio rerio*, ...
- 26°C
- medium:
 - pH 7,8 ± 0,2
 - CaCl₂·2H₂O, MgSO₄·7H₂O, NaHCO₃, KCl
 - dissolved O₂ > 90%
- 24, 48, 72, 96 h
- validity: mortality in control < 10%, O₂ etc.




Continuous:
-exposure is function of duration



Static renewal, with recovery:
-exposure is function of max conc'n



Static renewal, no recovery:
-exposure is function of cumulative concentration



Prolonged and chronic fish tests

Prolonged

- OECD 204 – 14d, ISO 10229 - 21 d
- semistatic or flow through
- food; controlled pH, O₂
- endpoints: breathing, gills, behavior, orientation, mortality
- 14-21d LC₅₀, NOEC, LOEC

Chronic





- OECD 210; US EPA OPPTS 850.1500
- 7 - 200 days
- starting with eggs or embryos or juveniles
- endpoints: survival, behavior, feeding, length, growth, weight, biochemical parameters, bioaccumulation

Embryolarval test (ISO 12890, OECD 210, 212)

- 100 embryos per replicate
- 6-7 days
- no food
- T, pH, O₂
- Endpoints: hatching, survival, morphology, behavior, weight, length, anomalies

FETAX – Frog Embryo Teratogenicity Assay Xenopus

- Choriogonadotropin → eggs after 9-12h
- 25 embryos per Petri dish, 10 ml test solution
- 24°C; pH 6.5-9
- 96h
- validity:
 - 6-aminonicotinamid
LC50 2,23 mg/ml, EC50 0,005 mg/ml
- mortality, growth, abnormalities
- Atlas of Abnormalities – John A. Bantle



FETAX



| |
|--------------------------------------|
| NaCl |
| NaHCO ₃ |
| KCl |
| CaCl ₂ |
| CaSO ₄ .2H ₂ O |
| MgSO ₄ |
| H ₂ O |
| pH |



Table 4.11 The Frog Embryo Teratogenesis Assay: *Xenopus* (FETAX)

| Test type | 96 h static renewal |
|----------------------------------|--|
| Organism | <i>Xenopus laevis</i> |
| Age of parent organism | Adult male: at least 2 years of age Adult female: at least 3 years of age |
| Size of parent organism | Adult male: 7.5–10 cm in crown-rump length Adult female: 10–12.5 cm in length |
| Feeding | Adult: three feedings per week of ground beef liver; liquid multiple vitamins should be added to the liver in concentrations from 0.05–0.075 cc/5 g liver |
| Experimental design | |
| Test vessel type and size | Adults: large aquarium or fiberglass or stainless steel raceways; side of tank should be opaque and at least 30 cm high. Breeding adults: 5- or 10-gallon aquarium fitted with a 1-cm mesh suspended approximately 3 cm from the bottom of the tank; nylon or plastic mesh is recommended; aquarium should be fitted with a bubbler to oxygenate the water; the top of aquarium should be covered with an opaque porous material such as a fiberglass furnace filter Embryos: 60-mm glass or 55-mm disposable polystyrene Petri dishes |
| Test solution volume | Adults: water depth should be 7-14 cm Embryos: 10 ml per dish |
| Exposure to test substance | Continuous throughout test |
| Replacement of test material | Every 24 h |
| Number of concentrations | 5 |
| Number of replicates per sample | 2 |
| Number of organisms per chamber | Adults: 4–6 per 1800 cm ² of water surface area Breeding adults: 2 Embryos: 25 |
| Test duration | 96 h |
| Physical and chemical parameters | |
| Temperature | Adult: 23 ± 3°C Embryos: 24 ± 2°C |
| Photoperiod | 12 h light / 12 h dark |
| pH range | 6.5 to 9 |
| TOC | 10 mg/l |
| Alkalinity and hardness | Between 16 and 400 mg/l as CaCO ₃ |
| Endpoint | Acute (mortality) and subacute (teratogenesis) |




Vibrio fischeri test (ISO 11348)

- 5-30 min
- luminiscence inhibition




- problem with particles and colour
→ flash test




Specific mechanisms of toxicity

- Cell lines (H4IIE.luc – diox.; MVLN, T47D.Luc – estrog.)
- Nuclear receptors (AhR, ER, AR, RAR/RXR)
- AhR/ER – luciferase reporter gene
- CALUX (Chemical Assisted LUCiferase eXpression)



Ligand (TCDD)



Adapted from Blankenship (1994)





Research centre
for toxic compounds
in the environment

Soil ecotoxicology bioassays

Exposure methods

- Tested chemical mixed with soil
 - Artificial soil (OECD, ISO)
 - Real soil (LUFA 2.2 ...)



- Topic applications, injections, forced feeding ... not so relevant



What is artificial soil ?



| Soil component | Content expressed on % dry mass basis |
|---|---------------------------------------|
| • Sphagnum peat (air dried), finely ground and with no visible plant remains | 10 |
| • Kaolinite clay (air dried), containing not less than 30 % kaolinite | 20 |
| • Industrial quartz sand (air dried), predominantly fine sand with more than 50 % by mass of particle size 0,05-0,2 mm (amount dependent on calcium carbonate required) | 70 |
| • Calcium carbonate (CaCO ₃ , pulverised, analytical grade) to obtain an initial pH of 6.0 ± 0.5 | 0.3-1.0 |

OECD 1984. Guideline for testing chemicals 207. Earthworm acute toxicity test.

- Is standard medium for many soil bioassays ...
- Is much more relevant than solution, agar, filter paper ...
- Should solve problem of high variability of natural soils ...
- Should resemble natural loamy soil ...
- Should enable the toxicity extrapolation to natural soils ...


LUFA soil (<http://lufa-speyer.de/>)

- Landwirtschaftliche Untersuchungs und Forschungsanstalt Speyer
- 4 EUR / 1 kg



| | LUFA 2.1 | LUFA 2.2 | LUFA 2.3 | LUFA 5M | LUFA 6S |
|--|-------------|-----------------|-----------------|-----------------|------------------|
| organic carbon (%) | 0.81 ± 0.21 | 2.16 ± 0.40 | 0.98 ± 0.05 | 1.29 ± 0.20 | 1.75 ± 0.11 |
| particles < 0.02 mm (%) | 8.2 ± 0.9 | 13.9 ± 1.1 | 22.7 ± 1.1 | 25.3 ± 1.8 | 65.1 ± 2.7 |
| pH (0.01M CaCl ₂) | 5.1 ± 0.4 | 5.4 ± 0.1 | 6.4 ± 0.6 | 7.2 ± 0.1 | 7.2 ± 0.1 |
| cation exchange capacity (meq/100g) | 4 ± 1 | 10 ± 1 | 8 ± 2 | 15 ± 3 | 22 ± 6 |
| water holding capacity (g/100g) | 33.2 ± 1 | 48.2 ± 5 | 34.4 ± 2 | 42.1 ± 4 | 40.7 ± 5 |
| weight per volume (g/1000ml) | 1404 ± 46 | 1197 ± 60 | 1291 ± 30 | 1212 ± 56 | 1264 ± 90 |
| Particle size (mm) distribution according to German DIN (in %): | | | | | |
| <0.002 | 3.0 ± 0.9 | 6.4 ± 0.9 | 9.4 ± 0.9 | 10.8 ± 1.3 | 42.1 ± 1.8 |
| 0.002 - 0.006 | 2.2 ± 0.7 | 3.5 ± 0.7 | 4.2 ± 0.8 | 5.4 ± 0.3 | 10.8 ± 0.7 |
| 0.006 - 0.02 | 2.9 ± 0.7 | 3.8 ± 0.7 | 9.1 ± 0.5 | 9.1 ± 0.5 | 12.1 ± 1.3 |
| 0.02 - 0.063 | 5.3 ± 1.8 | 5.4 ± 1.2 | 18.6 ± 2.3 | 19.5 ± 1.3 | 14.1 ± 2.5 |
| 0.063 - 0.2 | 27.0 ± 3.1 | 35.4 ± 2.3 | 29.3 ± 3.4 | 38.9 ± 1.0 | 8.7 ± 0.9 |
| 0.2 - 0.63 | 57.2 ± 4.3 | 44.8 ± 2.7 | 26.9 ± 0.7 | 14.9 ± 1.0 | 9.0 ± 0.3 |
| 0.63 - 2.0 | 2.4 ± 0.6 | 0.7 ± 0.1 | 2.5 ± 0.8 | 1.4 ± 0.1 | 3.2 ± 0.7 |
| soil type | sand (S) | loamy sand (LS) | loamy sand (LS) | silty sand (uS) | clayey loam (tL) |
| Particle size (mm) distribution according to USDA (in %) | | | | | |
| <0.002 | 3.0 ± 0.9 | 6.4 ± 0.9 | 9.4 ± 0.9 | 10.8 ± 1.3 | 42.1 ± 1.8 |
| 0.002 - 0.05 | 8.8 ± 1.8 | 12.2 ± 0.6 | 29.8 ± 3.0 | 27.5 ± 2.2 | 36.0 ± 2.3 |
| 0.05 - 2.0 | 88.2 ± 1.2 | 81.4 ± 1.2 | 60.8 ± 2.6 | 61.7 ± 3.2 | 21.9 ± 1.6 |
| soil type | sand | loamy sand | sandy loam | sandy loam | clay |

Earthworm bioassays




Earthworm acute toxicity test

- 500 g soil + 10 adult *Eisenia fetida*
- 14 days
- mortality and weight



Earthworm reproduction test ISO 11268-1


- 56 days
- 500 g soil + 10 adult *Eisenia fetida*
- horse manure as food
- juveniles extracted using water bath



Eisenia fetida reproduction test ISO 11268-2



Soil preparation



WHC measurement



Water added
Soil weighted to jars



10 adults to 1 jar



Weighting worms





10 adults from culture
Washed

E. fetida test – after 28 days

ISO 11268-2




Temperated room



Control of the jars, activity markers



Weighting the worms



Mortality assessment

E. fetida – 8 weeks

ISO 11268-2



After 20 min juveniles appear

Water bath, increasing temperature 40°C - 60°C



Collecting and counting juveniles



Sieving the soil



Hand sorting of cocoons



Counting

Avoidance test


ISO 17512-1

| | |
|----------------------|----------------------------|
| Guideline: | ISO/DIS 17512 (draft) |
| Species: | <i>E. andrei</i> |
| Substrate: | LUFA St. 2.2 standard soil |
| Duration: | 1 - 2 days |
| Parameter: | Behaviour of the worms |
| Test vessels: | Dual chamber |



Bioaccumulation test


| | |
|---|---|
| Test organism | <i>Eisenia fetida/andrei</i> , or <i>Enchytraeus albidus</i> , adult worms of similar size |
| Test substrate | spiked artificial soil based on OECD guideline No. 207 (OECD 1984) |
| Control substrate | uncontaminated artificial soil based on OECD guideline No. 207 (OECD 1984) |
| Biological parameters | concentration of test item in worms during uptake and elimination period |
| Endpoints | bioaccumulation factor, uptake rate coefficient, elimination rate constant |
| Test duration | uptake period: until steady state or 28 d; plus 10 day elimination period |
| Temperature | 20 ± 2°C |
| Light regime & light intensity | constant light; 400 to 800 lx |
| Test chambers | e.g. 250 mL/50 mL glass tubes; additional chambers of appropriate size for chemical analyses |
| Feeding during exposure | food added to soil directly after spiking; additional feeding once per week during exposure periodic addition of deionised water |
| Water | 4 days; spiked soil under test conditions ¹⁴ C-labelled/metal |
| Equilibration | if possible, test item dissolved in water mixed with dry soil; optional: coating of sand |
| Test item | in worms and soil |
| Spiking of soil | 1 plus control(s) to be fixed (expressed in Bq or mg kg ⁻¹ soil dry weight) |
| Analysis of test item | at least 3 per sampling date |
| Number of test concentrations | <i>Eisenia</i> : 1; <i>Enchytraeus</i> : 20 |
| Test concentrations | 4 samples after soil preparation |
| Number of replicates per test concentration/sampling date | 4 samples at end of uptake phase* |
| Number of organisms per test chamber | mortality during test period ≤ 10% (earthworms), ≤ 20% (enchytraeids) |
| Determination of soil dry weight and organic carbon content in soil | Use of appropriate methods (e.g. nonlinear regression analysis, ANOVA, Dunnett's t-test) |
| Determination of lipid content in biota | |
| Validity of test | |
| Evaluation | |




Enchytraeidae



Avoidance test with *E. albidus*







Folsomia candida



Folsomia candida


ISO 11267



Caenorhabditis elegans test

ASTM: E2172-01 Standard
Guide for Conducting
Laboratory Soil Toxicity Tests
with the Nematode
Caenorhabditis elegans

ISO 10872



Mites


Hypoaspis aculeifer



predator

cont. soil


prey



Lactuca sativa root growth

ISO 11269-1

Germination tests with plants – testing toxicity in the aquatic media (*Sinapis alba*) or in soil (*Lactuca sativa*)




Higher plants chronic toxicity ISO 22030





Soil microbial assay according to OECD, ISO

Real uncontaminated agricultural soil with indigenous microflora:

$\text{pH}_{\text{KCl}} = 7 - 7.5$ $C_{\text{bio}} = 400 - 700 \mu\text{g}\cdot\text{g}_{\text{d.w.}}^{-1}$
 $C_{\text{org}} = 1.5\%$ $\text{BR} = 0.5 - 0.7 \mu\text{g CO}_2\text{-C}\cdot\text{h}^{-1}\cdot\text{g}_{\text{d.w.}}^{-1}$
sand = 70%



Effects on microbial respiration






Micro & Mesocosms

Expensive & time consuming (e.g. *Pesticide testing*)

Variable results (natural variability ...)

Higher ecological relevancy






Research centre
for toxic compounds
in the environment


Results of the bioassays and their use

Data from ecotoxicity tests




Dose(concentration) - response relationship

Concentration:




96-hour LC50 = 50 µg/L



Concentration and Dose

Note

Concentration and dose both refer to the amount of test material to which the test organism is subjected. Concentrations are used to describe the amount of test material in the testing environment (e.g., mg/L in water, mg/kg in soil or mg/kg in food). Doses are used to describe the amount of test material administered to a subject (e.g., mg/kg-bodyweight in an avian bolus study). Statistical methods for both types of studies are identical; however, interpretations are different. Although "concentration" is used throughout this document, all the statistical methods presented here also apply to studies in which a dose is used.



No Observed Effect Concentration (NOEC)
Lowest Observed Effect Concentration (LOEC)
EC_x (x % effects concentration)
LC_x (x % lethal concentration)

Bioassay vs real ecosystem

- Bioassay is only simplified ecosystem model
- species differs, matrix differs, single species vs multispecies, individuals vs populations



Aporrectodea caliginosa



Lumbricus terrestris



VS



Eisenia fetida – compost worm !!

Bioassay vs reality

Objective = to protect real ecosystems, safe concentration


→ need to extrapolate bioassay results to be valid for ecosystems

- how much information we have?
- uncertainty factors, 1, 10, 100, 1000



Extrapolate


| Data | Assessment factor |
|---|-------------------|
| L(E)C50 short-term toxicity tests | 1000 |
| NOEC for 1 long-term toxicity test | 100 |
| NOEC for additional long-term toxicity tests of 2 trophic levels | 50 |
| NOEC for additional long-term toxicity tests of 3 species of 3 trophic levels | 10 |



PNEC (limits, EQS)



Species sensitivity distribution (SSD)




HC5 = 95% protection level


Risk assessment with earthworms



Prüfung der Auswirkungen auf Regenwürmer



Labortest mit Kompostwurm



Kokons des Kompostwurms



einheimische Regenwurmart

1. Akute Toxizität (2 Wochen)

Bewertung: Mortalität, Körpergewicht
 $TER = \frac{LC50}{PEC} < 10$

2. Einfluss auf die Fortpflanzung (8 Wochen)

Bewertung: Anzahl der Jungtiere, Körpergewicht
 $TER = \frac{NOEC}{PEC} < 5$

3. Auswirkungen im Freiland (1 Jahr)

Bewertung: Individuenzahlen, Risiken für Populationen und Lebensgemeinschaften

Summary - Take home message

- **Bioassays are necessary addition to chemical analyses**
- **Lot of standardized/experimental bioassays to choose from**
- **Necessary know-how (lab tricks, weakpoints, when to use, interpretation, data evaluation ...)**
- **No single test gives all information - battery of bioassays**

Thanks for your attention!!!!



Jakub Hofman

hofman@recetox.muni.cz

<http://www.recetox.cz>