


Diverzita savců

Asi 5 500 recentních druhů

Nové objevy, popisy a nálezy
New discoveries, descriptions and finds



Podle:

Jan Hošek, 2007: Saola aneb
největší zoologické objevy
posledních let.
Scientia, Praha 2007, 215 str.

Počty žijících druhů obratlovců podle přehledů uveřejněných v posledních třech letech a jejich ohrožení podle Červeného seznamu IUCN.

<i>skupina</i>	<i>odhad počtu známých druhů</i>	<i>počet vymřelých druhů</i>	<i>počet ohrožených druhů</i>
savci (Mammalia)	5416	70	1093
ptáci (Aves)	9934	135	1206
plazi („Reptilia“)	8240	22	341
obojživelníci (Amphibia)	5918	34	1811
svaloploutví (Sarcopterygii)	8	0	1
paprskoploutvé ryby (Actinopterygii)	26848	80	1058
paryby (Chondrichthyes)	970	0	110
kruhoústí (Cyclostomata)	108	0	2
<i>celkem</i>	57442	342	5621

Počet recentních druhů savců


		N čeledí	N druhů
Monotremata		2	5
Marsupialia		21	330
Placentalia	Afrotheria		71
	Xenarthra		31
	Euarchontoglires		
	Glires		
		Lagomorpha	2
		Rodentia	28-30
			2300
	Euarchonta		
		Scandentia	20
		Dermoptera	2
		Primates	15
			370
	Laurasiatheria		
		Eulipotyphla	450
		Chiroptera	18
		Artiodactyla	10
		Cetacea	10
			80
		Perissodactyla	3
		Pholidota	8
		Carnivora	12
			280


1901

Okapia johnstoni

-  - g. horská
-  nda – pohoří Virunga
-  ageni – prase pralesní

-  ká, J-Etiopie





„ZLATÝ VĚK OBJEVŮ“

Kopytníci od roku 1900 (Ungulates)

1901	okapi pruhovaná (<i>Okapia johnstoni</i>) – Kongo, Uganda (Giraffidae) mazama yucatánský (<i>Mazama pandora</i>) – Mexiko chocholatka Weynsova (<i>Cephalophus weynsi</i>) – Afrika chocholatka ruwensorská (<i>Cephalophus rubidus</i>)	
1903	antilopka zakrská (<i>Neotragus batesi</i>) – rovníková Afrika	
1904	prase pralesní (<i>Hylochoerus meinertzhageni</i>) –	
1908	mazama venezuelský (<i>Mazama bricenii</i>) – J Amerika	
1910	nyala horská (<i>Tragelaphus buxtoni</i>) - J Etiopie (mazama anoa horský (<i>Bubalus quarlesi</i>) – Celebes (Sulawesi) (Bovidae)	
1911	dikdik somálský (<i>Madoqua piacentinii</i>) – Somálsko (Bovidae)	
1914	goral červený (<i>Nemorhaedus baileyi</i>) – Tibet, Barma (Bovidae)	
1918	chocholatka zanzibarská (<i>Cephalophus adersi</i>) – V Africe	
1929	kára (kabar) kabar (<i>Schus berezovskii</i>) – J Čína, Sibirský poloostrov (Muntiacidae)	
1930	pudu (<i>Pudu pudu</i>) – Argentinie (Cervidae)	
1932	muntžak (<i>Muntiacus muntiacus</i>) – J Asie (Cervidae)	
1935	goral (<i>Naemorhedus goral</i>) – J Asie (Capricornidae)	
1937	kupřej (<i>Bos sauveini</i>) – Zadní Indie (Bovidae)	
1959	mazama zakrslý (<i>Mazama chunyi</i>) – J Amerika (Bovidae)	
1963	nahur Schaeferův (<i>Pseidois schaeferi</i>) – Indonésie (Capricornidae)	

- 1981 kabar čínský (*Moschus fuscus*) – Čína (Bovidae)
- 1982 muntžak žlutý (*Muntiacus atherodes*) – Borneo (Cervidae)
- 1985 gazela jemenská (*Gazella bilkis*) – Jemen (Bovidae)
- 1987 prase floreské (*Sus heureni*) – J Asie (Suidae)
- 1990 muntžak gongšanský (*Muntiacus gongshanensis*) – J Asie (Tibet, Čína), (Cervidae)
- 1993 saola (*Pseudoryx nghetinhensis*)** - Vietnam-Laos (Bovidae)
- 1994** muntžak obrovský (*Megamuntiacus vuquangensis*) – Vietnam-Laos (Cervidae)
- lyrorožec - ling (*Pseudonovibos spiralis*)** – Vietnam, Kambodža (Bovidae)
- 1996 mazama bororo (*Mazama bororo*) – JV Brazílie (Cervidae)
- 1997 muntžak černý (*Muntiacus truongsonensis*) – stř. Vietnam (Cervidae)
- 1999 muntžak listový (*Muntiacus putaoensis*) – Myanmar (Barma) (Cervidae)
- 2003 buvolec zambijský (*Damaliscus superstes*) – Zambie, Kongo (Bovidae)
- 2005 kančil cejlonský (*Moschiola kathygre*) – Sri Lanka (Tragulidae – kančilovití)
- voduška konžská (*Cobus anselli*) – Kongo (Bovidae)
- 2008 jelínek mazama (*Mazama ochroleuca*) – Amazónie




kančil

1937

Bos sauvelli

kuprej





SEARCH & FILTER

GO

advanced search

Journal Home

Current issue

AOP

Archive

TOP 100 ARTICLES

Download PDF

References

Top 100 cited

Supplementary

More information

About this article

More articles like this

Top 100 cited

Previous | Next

letters to nature

Nature 363, 443 - 445 (03 June 1993); doi:10.1038/363443a0

A new species of living bovid from Vietnam

VU QUANG UND, PHAM BONG QUAO, HOANGH NGOC CHINH DOTOUC, PETER ARCT AND CRAIG JOHN BLACKMAN¹

Forest Inventory and Planning Institute, Ministry of Forestry, Hanoi, Vietnam

Asian Bureau for Conservation, 18/E Capital Building, 175-191 Lockhart Road, Wan Chai, Hong Kong

To whom correspondence should be addressed.

IN May 1992 a joint survey by the Ministry of Forestry and World Wide Fund for Nature of the Vu Quang Nature Reserve, Ha tinh province, found three sets of long straight horns of a new bovid (Mammalia, Artiodactyla) in hunters' houses¹. None of the specimens had dentition. On four follow-up visits by Vietnamese scientists new specimens were discovered and surveys of forests in neighbouring Nghe an province revealed more localities and some partial specimens. In all, we have examined more than 20 specimens. Three have complete upper skulls and dentitions, two have lower jaws and dentitions. Three complete skins have been collected. The specimens are distinct in appearance, morphology and DNA sequence and cannot be ascribed to any known genus. Only two bovid genera are known from this part of Asia, *Eos* and *Naemorhedus* = *Capricornis*^{2,3}. A new genus and species are therefore



Copyright Mark Kostich


Muntiacus vuquangensis – muntžak obrovský 1994



Muntiacus truongsonensis – muntžak černý 1997



Muntiacus putaoensis – muntžak listový 1999



Pseudonovibos spiralis – ling 1994




Figure 1. Phylogenetic tree reconstructed from the combination of cytochrome b and lactoferrin sequences. Shown is a bootstrap 50 % majority-rule consensus tree performed with the maximum parsimony (MP) method [26]. *Antilocapra americana* (family Antilocapridae), *Cervus elaphus* and *Odocoileus hemionus* (family Cervidae) were used to root the bovid tree. The analyses were conducted with either equal weighting, or differential weighting of the nucleotide transformations (see section 5.2). Bootstrap proportions were computed after 1 000 replicates. These are indicated below branches, to the left of the slash for the weighted MP analysis and to the right for the unweighted MP analysis. The type and position of the molecular signatures characteristic of the different groups of species (i.e. exclusive synapomorphies), are indicated on the branches as shown in the key.



Moschiola kathygre – kančil cejlonský 2005

Kytovci od roku 1900 (Cetacean, Whale)


- 1908 vorvaňovec australský (*Mesoplodon bowdoini*)
- 1912 sviňucha jižní (*Australophocaena dioptrica*)
- 1913 vorvaňovec tmavý (*Mesoplodon mirus*)
- 1918 delfínowec čínský (*Lipotes vexillifer*)
- 1926 vorvaňovec Longmanův (*Indocetus pacificus*)
- 1934 delfín Graffmanův (*Stenella graffmani*)
- 1937 vorvaňovec Shepherdův (*Tasmacetus shepherdii*)
- 1956 plískavice saravacká (*Lagenodelphis hosei*)
- 1958 vorvaňovec japonský (*Mesoplodon gingkodens*)
sviňucha kalifornská (*Phocoena sinus*)
- 1963 vorvaňovec kalifornský (*Mesoplodon californicus*)
- 1991 vorvaňovec peruánský (*Mesoplodon peruvianus*)
- 2002 vorvaňovec Perrinův (*Mesoplodon perrini*) – DNA
- 2003 plejtvák Omurův (*Balaenoptera amurai*) – DNA
- 2005 orcela tupoploutvá (*Orcaella heinsohni*)

Letouni po roce 2000

2000	3 druhy
2001	6 druhů, v Evropě <i>Myotis alcathoe</i> – netopýr alkathoe (menší, nymfin)
2002	8 druhů
2003	3 druhy
2004	10 druhů, <i>Plecotus gaisleri</i> – S Afrika, <i>Pipistrellus hanaki</i> - Libye
2005	13 druhů
2006	9 druhů
2007	1 druh

Ostatní savci po roce 1990

1995	prase vietnamské (<i>Sus bucculentus</i>) – Laos, popis 1892, Lazarus taxon králík Timminsův (<i>Nesolagus timminsi</i>) – Laos
1996	khanyou (<i>Laonastes aenigmaemus</i>) – stř. Laos – skalní krysa (Laonastidae), Lazarus taxon, bazální linie hystricognáthních hlodavců
1997	langur duk (<i>Pygathrix nemaeus</i>) – Vietnam (Cercopithecidae) cibetka tainguenská (<i>Viverra tainguensis</i>) – Vietnam, Annamity
2001	slon pralesní (<i>Loxodonta cyclotis</i>) – kryptický druh, DNA, dnes nepřijímán



• BIODIVERZITA





Figure 2. Global distribution of new mammals described since 1992. The distribution is overlaid on currently recognized regions of high threat and irreplaceability. Variable levels of shading indicate the number of global biodiversity conservation templates that prioritize the region (Brooks et al. 2005).

SAVCI – distribuce nových druhů objevených po 1992



Jižní Amerika

Primáti

1998 *Callithrix humilis* – kosman (*Callithrichidae*, drápkaté opice)

2000 *Callithrix manicorensis*

Callithrix acariensis

2003 *Callicebus bernhardi* – titi (*Cebidae* – malpovití)

Callicebus stephennashi

? *Callicebus sp.nov.* 1

Callicebus sp.nov. 2

Pithecia sp.nov. 1 - chvostan

Pithecia sp.nov. 2

Saguinus sp.nov. 1 - tamarín

Saguinus sp.nov. 2

Ateles sp.nov. 1 - chápan

Ateles sp.nov. 2

Cacajao sp.nov. - uakari

Lagothrix sp.nov. 1 - chápan

Lagothrix sp.nov. 2

Callithrix sp.nov. - kosman

Callicebus sp. nov. - titi

Saimiri sp.nov. - kotul

Ostatní

Agouti sp. nov. – paka (Agoutidae)



Eira sp.nov. – hyrare, brazilská kuna (Mustelidae)


Nasua sp.nov. – nosál (Procyonidae)

Myrmecophaga sp.nov. – mravenečník (Myrmecophagidae)

Pteronura sp.nov. – vydra (Mustelidae)

Panthera sp.nov. – jaguár (Felidae)





Marc van Roosmalen

24.7. 1947, Tilburg, Holandčan žijící v brazilském Manau, primatolog

<http://www.marcvanroosmalen.org/news.htm>



marcvanroosmalen.org

ENTER

NAMED NEW SPECIES

Black-crowned dwarf marmoset

Callibella humilis

Rio Acarí Amazonian marmoset

Mico (Callithrix) acariensis

Sataré Amazonian marmoset

Mico (Callithrix) saterei

Rio Manicoré Amazonian marmoset

Mico (Callithrix) manicorensis

Stephen Nash's titi monkey

Callicebus stephennashi

HRH Prince Bernhard's titi monkey

Callicebus bernhardi

Giant collared peccary

Pecari maximus

Van Roosmalen's dwarf porcupine

Sphiggurus roosmalenorum - kuandu

Dwarf manatee - kapustňák

Trichechus pygmaeus sp. nov.

Lecythidaceae (Brazilnut Family)

Lecythis oldemanii sp. nov.

Black dwarf lowland tapir

Tapirus pygmaeus sp. nov.

Van Tienhoven's fair brocket deer

Mazama tienhoveni sp. nov.

NEW SPECIES SEARCH

Arboreal giant anteater - *Myrmecophaga* sp. nov.

White-throated black jaguar - *Panthera* sp. nov.

Black giant otter - *Pteronura* sp. nov.

Orange coati-mundè - *Nasua* sp. nov.

Orange tayra - *Eira* sp. nov.

Black woolly monkey - *Lagothrix* sp. nov.

Cruz Lima's saddleback tamarin monkey

Saguinus (fuscicollis) cruzlimai sp. nov.

Rio Pauini white bald-headed uacari

Cacajao (calvus) sp. nov.

Rio Aripuanã green-backed squirrel monkey

Saimiri (ustus) sp. nov.

Rio Mamurú titi monkey - *Callicebus (moloch)* sp. nov.

Upper Xingú Amazonian marmoset monkey

Mico (Callithrix) sp. nov.

Orange woolly monkey - *Lagothrix* sp. nov.

Long-limbed black spider monkey - *Ateles* sp. nov.

Silvery bellied spider monkey - *Ateles* sp. nov.

Eastern saddleback tamarin monkey

Saguinus (fuscicollis) orientalis sp. nov.

Rio Purús collared titi monkey *Callicebus (torquatus)* sp. nov.

Upper Rio Xingú titi monkey - *Callicebus (moloch)* sp. nov.

Grey saki monkey - *Pithecia* sp. nov.

Southbank Rio Negro saki monkey

Pithecia (Pithecia) sp. nov.

A NEW SPECIES OF LIVING PECCARY (MAMMALIA: TAYASSUIDAE)
FROM THE BRAZILIAN AMAZON

Marc G. M. VAN ROOSMALEN, Lothar FRENZ, Pim VAN HOOFT, Hans H. DE IONGH & Herwig LEIRS



Pecari maximus 2007

– největší pekari

A NEW SPECIES OF LIVING MANATEE FROM THE AMAZON

Shallow clear-water adapted dwarf manatee is already on the verge of extinction

Marc G.M. van Roosmalen¹, Pim van Hooft² & Hans H. de Iongh³

¹ AAPA Manaus-Amazonas, Brazil

² Wageningen University and Research Centre, Resource Ecology Group, Bornsesteeg 69, 6708PD Wageningen, The Netherlands

³ Leiden University, Institute of Environmental Sciences, PO Box 9518, 2300RA Leiden, The Netherlands



(LEFT) *The Rio Arauazinho harbors dwarf manatee and numerous other new mega-fauna species. Marc proposes this be the cornerstone for a new Brazilian National Park.*

Trichechus bernhardi 2007

- nejmenší kapustňák

A NEW SPECIES OF LIVING BROCKET DEER
(MAMMALIA: CERVIDAE) FROM THE BRAZILIAN AMAZON

Marc G. M. VAN ROOSMALEN & Pim VAN HOOFT



(LEFT) *Mazama ochroleuca* sp. nov. redrawn from plate depicting *Mazama (gouazoupira) nemorivaga* (Eisenberg, 1989).

(ABOVE) Two spikes of *Mazama americana* above, one of *Mazama ochroleuca* sp. nov. below.

Here we report on the existence of a new species of even-toed ungulate in the Brazilian Amazon, which we name *Mazama ochroleuca* sp. nov., the fair brocket deer. It is intermediate in size between the two known species of brocket deer, *Mazama americana* and *Mazama (gouazoupira) nemorivaga*, and occurs in sympatry with both. Preliminary mitochondrial partial cytochrome b sequences of fair brocket deer compared with that of the sympatric and morphologically most related grey brocket deer {*Mazama (gouazoupira) nemorivaga*} revealed a sequence difference of 3.7%. Divergence time is therefore estimated at 1.0 million years before present. As in other brocket deer, fair brocket deer seem to live solitary or in pairs. In view of recent developments in the Rio Aripuanã basin where it lives and due to its limited distribution, we consider the fair brocket deer highly endangered.

KEY WORDS

New species, Artiodactyla, Cervidae, *Mazama ochroleuca* sp. nov., fair brocket deer, Brazilian Amazon

Mazama ochroleuca 2008

A NEW SPECIES OF LIVING LOWLAND TAPIR (MAMMALIA: TAPIRIDAE) FROM THE BRAZILIAN AMAZON

Marc G.M. van Roosmalen / AAPN Manaus-Amazonas, Brazil



(LEFT) Adapted rendering of a black dwarf lowland tapir.
(ABOVE) On the left a skull of Brazilian lowland tapir, on the right the skull of black dwarf lowland tapir.

Here we report on the existence of a new species of odd-toed ungulate in the Brazilian Amazon, which we name *Tapirus pygmaeus* sp. nov., the black dwarf lowland tapir. It is much smaller than the since long known Brazilian lowland tapir, *Tapirus terrestris*. As in other tapirs, dwarf tapirs seem to live solitary or in pairs. In view of recent developments in the interfluves where it lives and due to its limited distribution and apparent rariness, we consider the dwarf tapir highly endangered.

KEY WORDS

New species, Perissodactyla, Tapiridae, *Tapirus pygmaeus* sp. nov., black dwarf lowland tapir, Brazilian Amazon

Tapirus pygmaeus 2008 – nejmenší tapír

Současné tempo popisování


- RYBY cca 150 druhů ročně
- OBOJŽIVELNÍCI cca 70 druhů ročně
- PTÁCI cca 5 - 10 druhů ročně
- SAVCI cca 40 – 50 druhů ročně

1992 – 2005


ptáci vs. savci

83 : 341

Log of Birds (+) and Mammals (o) described per year



Log of Birds (-) and Mammals (- -) described per year



REEDER ET AL.—NEW MAMMALS CONTINUE TO BE DISCOVERED






Figure 1. Cumulative and decadal descriptions of taxonomically valid extant mammal species.

Patterson 2001




Patterson 2001




Chiroptera


Patterson 2001



Patterson 2001




Patterson 2001





Rodentia

Patterson 2001



Patterson 2001







Andy

Cuscomys ashaninka

– činčilák ašaninka (Abrocomidae – činčilákovití) – 1999 L.H. Emmonsová



Abrocoma oblativa Eaton 1916
Cuscomys oblativus

Cuscomys ashaninka

1992 – 2005

341 nový druh savců

1992 – 2005

Rodentia 155 druhů



Mallomys sp. nov. – krysa (Murinae)

1992 – 2005

Chiroptera 78 druhů



Styloctenium mindorensis Esselstyn 2007 – kaloň, nízinné lesy Filipín

1995 – 2007

36 nových druhů primátů

2006

Tarsius lariang – nártoun (kombovití – Galagonidae)

Microcebus jollyae – maki (makiovití - Cheirogaleidae)

Microcebus mittermeieri

Microcebus simmonsi

Microcebus mamaratra

Lepilemur aeaeclis – lemur (lemurovití noční – Megaladapidae)

Lepilemur randrianasoli

Lepilemur sahamalazensis

Lepilemur ahmansi

Lepilemur betsileo

Lepilemur fleuretae

Lepilemur grewcocki

Lepilemur hubbardi

Lepilemur jamesi

Lepilemur milanoii

Lepilemur petteri

Lepilemur seali

Lepilemur tymerlachsoni

Lepilemur wrighti

Avahi peyrierasi – avahi (indriovití – Indridae)

Cebus queirozi – malpa (malpovití – Cebidae)

2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis _____
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansi
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae - maki
Microcebus mittermeieri —
Microcebus simmonsi
Microcebus mamilatra
Lepilemur aeeclis - lemur
Lepilemur randrianasoli
Lepilemur sahamalzensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi - malpa



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansi
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalzensis
Lepilemur ahmansonii _____
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalzensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi _____
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang _____
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus queirozi



2006

Tarsius lariang
Microcebus jollyae
Microcebus mittermeieri
Microcebus simmonsi
Microcebus mamaratra
Lepilemur aeeclis
Lepilemur randrianasoli
Lepilemur sahamalazensis
Lepilemur ahmansonii
Lepilemur betsileo
Lepilemur fleuretae
Lepilemur grewcocki
Lepilemur hubbardi
Lepilemur jamesi
Lepilemur milanoii
Lepilemur petteri
Lepilemur seali
Lepilemur tymerlachsoni
Lepilemur wrighti
Avahi peyrierasi
Cebus flavius




Lophocebus kipunji *Rungwecebuss*



Tanzánie 2005

mangabej kipundži – kočkodanovití (Cercopithecidae)





Tim Davenport / WCS

because of the terrain, thick secondary forest, and the animal's cryptic nature, sightings were infrequent and poor. It was not until December 2003, during work in the contiguous Livingstone Forest, that the monkey was clearly observed and recognized as a new species of mangabey.

Ndundulu population. The Udzungwa Mountains (Udzungwas: 10,000 km², 07°40'S to 08°40'S and 35°10'E to 36°50'E) lie 350 km to the northeast of Rungwe-Livingstone (4, 5). Supporting circa (ca.) 1017 km² of fragmented forest (6), the Udzungwas receive a maximum annual rainfall of roughly 2200 mm and were previously thought to hold 10 primate species (7), including the endemic Sanje mangabey, *Cercocebus sanjei*, discovered in 1979 (8).

Two populations of the Sanje mangabey are known from the Udzungwas (7, 9). During visits from 1991 to 2000, ornithologists working in the Ndundulu Forest Reserve (Fig. 1) reported a third population of the Sanje mangabey (10). Subsequent surveys failed to confirm the presence of this species in Ndundulu (7, 9, 11) and led to our intensified surveys in July and September 2004. During these surveys, Sanje mangabeys were not encountered or heard. However, on 7 July 2004, the new species of mangabey was discovered. It now seems certain that the ornithologists had misidentified the new species of mangabey as the Sanje mangabey.

The researchers working on each of these two new populations of mangabeys did not become aware that a second population was known until October 2004.

Lophocebus kipunji Ehardt, Butynski, Jones, and Davenport sp. nov.



Fig. 2. Holotype: adult male highland mangabey *Lophocebus kipunji* in the type locality, Rungwe-Livingstone, Tanzania. [Photograph by T.R.B. Davenport]

live individual should be collected at this time to serve as the holotype. The Rungwe-Livingstone population is designated the source population for physical specimens in support of the holotype.

Paratype. Adult in photograph (Fig. 3). Sex not known. Photograph taken in Ndundulu Forest Reserve (07°48'45"S 36°31'05"E), Udzungwa Mountains, Tanzania.



Fig. 3. Paratype: adult highland mangabey *Lophocebus kipunji*, Ndundulu Forest Reserve, Tanzania. [Photograph by T. Jones]

tail off-white. Crown with very long, broad, erect crest of hair. Eyelids black, not contrasting with color of face. Adults emit a distinctive, loud, low-pitched "honk-bark" (Fig. 4). Arboreal. Found only at high altitudes (1300 m up to 2450 m asl) and low-temperature tolerant; temperatures in Rungwe-Livingstone drop to at least -3°C.

Description. A primarily brown, medium-sized, long-tailed, arboreal monkey. Muzzle elongated. Facial skin, including eyelids, black. Suborbital fossae "tear line" pronounced. Eyes brown. Pelage light to rufous brown except as follows: center of ventrum and distal half of tail, white to off-white; hands and feet, black; lower forelimbs, dark brown;






Fig. 2. Holotype: adult male highland mangabey *Lophocebus kipunji* in the type locality, Rungwe-Livingstone, Tanzania. [Photograph by T.R.B. Davenport]

Holotype. Adult male in [REDACTED] (Fig. 2). Photograph taken in the type locality at 9°07'S 33°44'E (12). The number of individuals in each of the two populations of this species is undoubtedly very small; no live individual should be collected at this time to serve as the holotype. The Rungwe-Livingstone population is designated the source population for physical specimens in support of the holotype.

Lophocebus kipunji



Rungwecebus kipunji



TANZANIA

Protected Areas

(not an authority on boundaries)




Galagoides udzungwensis – kombá

1996

Congosorex phillipsorum - bělozubka

2005





Rhynchocyon udzungwensis Rathbun 2008

Ital Rovero – 2005 fotopasti, Dr. Rathbun – Kalifornská akademie věd - 2008
Afrotheria – Macroscelidea – bercoun = elephant shrew, 1 kg, 30 cm






Dr. Rathbun







Národní přírodovědné muzeum Paříž

Genneta bourloni 2003 – ženetka Bourlonova
coll. 1959 – podle kůže, stř. Z Afrika



...na trhu







Neofelis nebulosa diardi – nový druh, pův. poddruh levharta obláčkového
Borneo 2006, 1/3 terestrických savců na Borneu - endemiti







© Tristan Savatier - www.loupiote.com





1995

...fotopasti



© The Natural History Museum, London

Nesolagus netscheri – králík krátkouchý (Leporidae),
Sumatra, od 1929 nezvěstný



1998: *Nesolagus netscheri*



Nesolagus netscheri



Nesolagus timminsi 2000 – králík Timminsův

Trh v Laosu 1995 - Timmins, 2000 - popsán Rusy, endemit Anamitského pohoří na hranici Laosu a Vietnamu



Nesolagus timminsi 2000



Laonastes aenigmamus 2005
- khanyou (Laonastidae), skalní krysa (Laos=kámen, skála –
řecky, tajuplná skalní myš z Laosu)

Nové čeledi na základě nálezů nových druhů

1905



Calomyscidae – křečkové myší (také jako podčeledě – Calomyscinae)
– blízký a střední východ

1918



Lipotes vexillifer – delfínovec čínský (ex. 2007)

vymřelá čeleď Lipotidae (Irinnidae nebo Platanistidae) - delfínovcovití

1974




Craseonycteridae



Extinct relatives
Diatomys shantungensis fossil (20 mya)
from Shandong, China

Laonastes aenigmamus (extant)
Khammouan Province, Laos

Huchon et.al. 2007



Pectinator spekei – gundi somálský

Ctenodactylus gundi – gundi saharský (Ctenodactylidae)




Massoutiera mzabi – g. východní





Ctenodactylus gundi

- ...fotopasti



1998
Pseudoryx nghetinhensis



2007
Muntiacus vuquangensis

Bornejská kočkoliška (2003)



1. puchol hnědý *Diplogale hosei* (Viveridae) – netypické zbarvení
2. nový druh cibetky (2006)
3. poletucha Thomasova *Aeromys thomasi* (Sciuridae)
Meijaard et al., Mammal Review 2006


















Lutreola thomasi vliegende oehoorn
Evan Hartweg 56

3




6

Aeromys thomasi



Lampros thomasi slagerde oehoorn
Evan Hartweg 186





IUCN Small Carnivore Specialist Group - Mohd. Azlan J.

2003 *Diplogale hosei* – puchol hnědý (Borneo)



Daniela De Luca / WCS

2006 *Bdeogale jacksoni* – mangusta Jacksonova (Herpestidae)



2007 *Dicerorhinus sumatrensis* – nosorožec sumaterský



© Ben Collen/ ZSL

Stealth Cam 02-03-2008 00:46:48

2008 *Hexaprotodon liberiensis* – hrošík liberijský

© Ben Collen/ ZSL



Stealth Cam 02-11-2008 08:02:38

Hexaprotodon liberiensis



ZSL

Hexaprotodon liberiensis



2003 *Catopuma badia* –
kočka bornejská



Thylacinus cynocephalus
1936 - ?




Image: David Fleay




Figure 1 Species richness of mammals across Europe. This illustrative map was prepared using Worldmap software and shows species richness (number of species per grid square) with one level of smoothing. The entire atlas dataset has been used, including introduced species. There are some differences between the *Atlas Flora Europaea* UTM grid used by this software and the one used in the Atlas, particularly around the south-eastern borders of Europe and also in the treatment of some island groups.

