

CG920 Genomics

Lesson 5

Gene Expression and Chemical Genetics

Jan Hejátko

Functional Genomics and Proteomics of Plants,
Mendel Centre for Plant Genomics and Proteomics,
Central European Institute of Technology (CEITEC), Masaryk University, Brno
hejatko@sci.muni.cz, www.ceitec.muni.cz



INVESTICE DO ROZVOJE VZDĚLÁVÁNÍ

Tato prezentace je spolufinancována
Evropským sociálním fondem
a státním rozpočtem České republiky

Literature

- Literature sources for Chapter 05:
 - Surpin, M. and Raikhel, N. (2004) Traffic jams affect plant development and signal transduction. *Nature Reviews/Molecular Cell Biology* 5,100-109
 - Zouhar, J., Hicks, G.R. and Raikhel, N.V. (2004) Sorting inhibitors (Sortins): Chemical compounds to study vacuolar sorting in Arabidopsis. *Proceedings of the National Academy of Sciences of the U.S.A.*, 101, 9497–9501
 - Nevo-Dinur, K., Nussbaum-Shochat, A., Ben-Yehuda, S., and Amster-Choder, O. (2011). Translation-independent localization of mRNA in *E. coli*. *Science* 331, 1081-1084.
 - Lecuyer, E., Yoshida, H., Parthasarathy, N., Alm, C., Babak, T., Cerovina, T., Hughes, T.R., Tomancak, P., and Krause, H.M. (2007). Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. *Cell* 131, 174-187.
 - Schonberger, J., Hammes, U.Z., and Dresselhaus, T. (2012). In vivo visualization of RNA in plants cells using the lambdaN(22) system and a GATEWAY-compatible vector series for candidate RNAs. *The Plant journal : for cell and molecular biology* 71, 173-181.

Outline



- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis
 - Quantitative analysis of gene expression
 - DNA and protein chips
 - Next generation transcriptional profiling
- Regulation of gene expression in the identification of gene function by gain-of-function approaches
 - T-DNA activation mutagenesis
 - Ectopic expression and regulatable gene expression systems
- Chemical genetics

Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene


Gene expression

- **Transcriptional fusion with a promoter region**
 - Identification and cloning of the promoter region of the gene
 - Preparation of recombinant DNA carrying the promoter and the reporter gene (uidA, GFP)




Gene expression

- **Transcriptional fusion with a promoter region**
 - Identification and cloning of the promoter region of the gene
 - Preparation of recombinant DNA carrying the promoter and the reporter gene (uidA, GFP)
 - Preparation of transgenic organisms carrying this recombinant DNA and their histological analysis



GUS reporter in mouse embryos





LacZ activity marks the cells of the developing myotome.

Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene

Gene expression

- **Translational fusion of coding region of analysed gene with a reporter gene**
 - Identification and cloning of the promoter and coding region of the analysed gene
 - Preparation of a recombinant DNA carrying the promoter and the coding sequence of the studied gene in fusion with the reporter gene (uidA, GFP)




Gene expression


- **Translational fusion of coding region of analysed gene with a reporter gene**
 - Preparation of transgenic organisms carrying the recombinant DNA and their histological analysis
 - Compared to transcriptional fusion, translation fusion allows analysis of e.g. Intercellular localization of gene product (protein) or its dynamics



PIN1-GFP in *Arabidopsis*



Histone 2A-GFP in *Drosophila* embryo by PAM





Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases


Gene expression

- Analysis of expression using Genevestigator (**AHP1** and **AHP2**, *Arabidopsis*, Affymetrix ATH 22K Array)




Gene expression

- Analysis of expression using Genevestigator (*AHP1* and *AHP2*, *Arabidopsis*, Affymetrix ATH 2K Array)



Gene expression

- Analysis of expression using Genevestigator (**AHP1** and **AHP2**, Arabidopsis, Affymetrix ATH 22K Array)





Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis

Expression Maps - RNA




High-Resolution Expression Map in Ara



Microarray expression profiles of 19 fluorescently sorted GFP-marked lines were analyzed (3–9, 23, 24). The colors associated with each marker line reflect the developmental stage and cell types sampled. Thirteen transverse sections were sampled along the root's longitudinal axis (red lines) (10). CC, companion cells.

Expression Maps - RNA


- High-Resolution Expression Map in Arabidopsis Root



(A) The majority of enriched GO terms (hierarchically clustered) are associated with individual cell types (blue bar). A smaller number are present across multiple cell types (green bar). (fig. S2) (B) GO category enrichment for hair cells confirms a previous report (15). Enriched cis-elements and an enriched TF family were also identified. (C) From the top 50% of varying probe sets, 51 dominant radial patterns were identified. Pattern expression values were mean-normalized (rows) and \log_2 transformed to yield relative expression indices for each marker line (columns). Marker line order is the same for all figures; see table S1 for marker line abbreviations. (D) Pattern expression peaks were found across one to five cell types. (E to G) Patterns where expression is enriched in single and multiple cell types support transcriptional regulation of auxin flux and synthesis. In all heat maps with probe sets, expression values were mean-normalized and \log_2 transformed. Expression is false-colored in representations of a root transverse section, a cut-away of a root tip, and in a lateral root primordium. (E) Auxin biosynthetic genes (*CYP79B2*, *CYP79B3*, *SUPERROOT1*, and *SUPERROOT2*) are transcriptionally enriched in the QC, lateral root primordia, pericycle, and phloem-pole pericycle ($P = 1.99E^{-11}$, pattern 5). All AGI identifiers and TAIR descriptions are found in table S14. (F) Auxin amido-synthases *GH3.6* and *GH3.17* that play a role in auxin homeostasis show enriched expression in the columella, just below the predicted auxin biosynthetic center of the QC ($P = 8.82E^{-4}$, pattern 13). (G) The expression of the auxin transporter, *PIN-FORMED2*, and auxin transport regulators (*PINOID*, *WAG1*) are enriched in the columella, hair cells, and cortex ($P = 1.03E^{-4}$, pattern 31).

Expression Maps - Proteins

□ Human Protein Atlas



Schematic flowchart of the Human Protein Atlas. For each gene, a signature sequence (PrEST) is defined from the human genome sequence, and following RT-PCR, cloning and production of recombinant protein fragments, subsequent immunization and affinity purification of antisera results in immunospecific antibodies. The produced antibodies are tested and validated in various immunoassays. Approved antibodies are used for protein profiling in cells (immunofluorescence) and tissues (immunohistochemistry) to generate the images and protein expression data that are presented in the Human Protein Atlas (Ponten et al., *J Int Med*, 2011).

Expression Maps - Proteins

- Human Protein Atlas
(<http://www.proteinatlas.org/>)


THE HUMAN PROTEIN ATLAS

ABOUT & HELP

SEARCH ? »

Search Clear Fields »

e.g. CD44, ELF3, KLK3, or use Fields to search specific fields such as protein_class:Transcription factors or chromosome:X



dictionary: histology of esophagus

News

Protein evidence according to [Fagerberg et al](#) is summarized in the [chromosome progress diagram](#).

Version: **11.0**
Atlas updated: 2013-03-11
[release history](#)

15156 genes with protein expression profiles based on **18707** antibodies.


*Knut och Alice
Wallenbergs
Stiftelse*

The Human Protein Atlas project is funded by the Knut & Alice Wallenberg foundation.




Expression Maps - Proteins


- Human Protein Atlas
(<http://www.proteinatlas.org/>)




SUBCELLULAR LOCATION SUMMARY ? »




Show image »

Main location(s)	Nucleus but not nucleoli
Additional location(s)	
Staining summary	Localized to the nucleus but excluded from the nucleoli.
Reliability (APE)	 High
Antibodies in assay	CAB039238, CAB039239


MORE SUBCELL DATA















NORMAL TISSUE & ORGAN SUMMARY ? »



Show image »

Expression summary	Fractions of cells showed weak nuclear and/or cytoplasmic expression.	
Tissue specificity	Expressed in 11 out of 82 cell types	
Reliability (APE)	 High	
Antibodies in assay	CAB002973, CAB039238, CAB039239	

Organ	No of cell types	Protein expression
CNS (brain)	11	
Hematopoietic (blood)	8	
Liver and pancreas	5	
Digestive (GI-tract)	13	
Respiratory (lung)	4	
Cardiovascular	1	
Female tissues	13	
Placenta	2	
Male tissues	5	
Urinary tract (kidney)	3	
Skin and soft tissues	14	
Endocrine tissues	3	

MORE TISSUE DATA

Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis
 - Quantitative analysis of gene expression
 - DNA and protein chips

DNA Chips


■ DNA chips

- Method, which provides quick comparison of a large number of genes/proteins between the test sample and control
- Oligo DNA chips are used the most



▪ There are commercially available kits for the whole genome

- company Operon (Qiagen), 29.110 of 70-mer oligonucleotides representing 26.173 genes coding proteins, 28.964 transcripts and 87 microRNA genes of *Arabidopsis thaliana*
- Possibility of use for the preparation of photolithography chips – facilitation of oligonucleotide synthesis e.g. for the whole human genome (about $3,1 \times 10^9$ bp) it is possible to prepare 25-mers in only 100 steps, by this technique



- Chips not only for the analysis of gene expression, but also for e.g. Genotyping (SNPs, sequencing with chips, ...)

Affymetrix ATH1 *Arabidopsis* genome array

Critical Specifications	
Number of arrays	One
Number of sequence represented	>24,000 gene sequences
Feature size	18 μ m
Oligonucleotide probe length	25-mer
Probe pairs/sequence	11
Control sequences	<i>E. coli</i> genes <i>bioB</i> , <i>bioC</i> , <i>bioD</i> . <i>B. subtilis</i> gene <i>lysA</i> . Phage P1 <i>cre</i> gene. <i>Arabidopsis</i> maintenance genes GAPDH, Ubiquitin, and Actin
Detection sensitivity	1:100,000*

*As measured by detection in comparative analysis between a complex target containing spiked control transcriptions and a complex target with no spikes.

DNA Chips

- DNA chips, analysis of results
 - For the correct interpretation of the results, good knowledge of advanced statistical methods is required
 - It is necessary to include a sufficient number of controls and repeats
- Control of accuracy of the measurement (repeated measurements on several chips with the same sample, comparing the same samples analysed on different chips with each other)
- Control of reproducibility of measurements (repeated measurements with different samples isolated under the same conditions on the same chip – comparing with each other)
- Identification of reliable measurement threshold
- Finally comparing the experiment with the control or comparing different conditions with each other -> the result
- Currently there's been a great number of results of various experiments in publicly accessible databases

Expression of 195M6T7 in response to chemical treatment

Home | About TAIR | Sitemap | Contact | Help | Order | Login

Search | Tools | Arabidopsis Info | News | Links | FTP | Stocks

Gene Search

Experiment: Aluminum Stress

Experiment Summary | Samples | Slides & Datasets | Array Design | View All

Slide Details

Slide (name : description)	External ID	Replicate (id :name)	Replicate type	Reverse replicate	Sample	Experimental variables	Label	Get Data
HoekengaS7 (*) : Aluminum Stress 1 [strong spatial bias]	AFGC: 7304	63: Aluminum Stress	technical		7304_Cy3.7305_Cy5	no treatment (pool of 3, 8, and 24 hours)	Cy3	Download
					7304_Cy5.7305_Cy3	Aluminum (50 5M AlCl3, pool of 3, 8, and 24 hours)	Cy5	
HoekengaSc Aluminum Stress 2 [strong spatial bias]	AFGC: 7305	64: Aluminum Stress	technical	63	7304_Cy5.7305_Cy3	Aluminum (50 5M AlCl3, pool of 3, 8, and 24 hours)	Cy3	Download
					7304_Cy3.7305_Cy5	no treatment (pool of 3, 8, and 24 hours)	Cy5	


Protein Chips

- Protein chips
 - Chips with high density containing 10^4 proteins
 - Analysis of protein-protein interactions, kinase substrates and interactions with small molecules
 - Possibility of using antibodies – more stable than proteins

Protein Chips

■ Identification of proteins interacting with integrin $\alpha_{IIb}\beta_3$ cytoplasmic domain of platelets

- Expression of cytoplasmic part as a fusion peptide biotin-KVGFFKR
- Analysis of binding to the protein chip containing 37.000 clones of *E. coli* expressing human recombinant proteins
- Confirmation of interaction by pull-down analysis of peptides and by coprecipitation of whole proteins as well (e.g. chloride channel Icn)
- Other use: e.g. in the identification of kinase substrates, when substrates are bound to the chip and exposed to kinases in the presence of radiolabeled ATP (786 purified proteins of barely, of which 21 were identified as CK2 α kinase substrates; Kramer et al., 2004)






Lueking et al., 2005

Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis
 - Quantitative analysis of gene expression
 - DNA and protein chips
 - **Next generation transcriptional profiling**

Next Gen Transcriptional Profiling

- *Transcriptional profiling* via *RNA sequencing*



Library Preparation
~2 h [15 min hands-on (Nextera)]
< 6 h [< 3 h hands-on (TruSeq)]



Cluster Generation
~5 h (<10 min hands-on)



Sequencing by Synthesis
~1.5 to 11 days



CASAVA
2 days (30 min hands-on)

Sequencing by Illumina and
number of transcripts determination

Results of –omics Studies vs Biologically Relevant Conclusions

- Transcriptional profiling yielded more than **7K differentially regulated genes...**

Dii et al., unpublished

gene	locus	sample_1	sample_2	status	value_1	value_2	log2(fold_change)	test_stat	p_value	q_value	significant
AT1G07795	1:2414285-2414967	WT	MT	OK	0	1,1804	1.79769e+308	1.79769e+308	6.88885e-05	0,00039180	1 yes
HRS1	1:4556891-4558708	WT	MT	OK	0	0,696583	1.79769e+308	1.79769e+308	6.61994e-06	4.67708e-05	yes
ATMLO14	1:9227472-9232296	WT	MT	OK	0	0,514609	1.79769e+308	1.79769e+308	9.74219e-05	0,00053505	5 yes
NRT1.6	1:9400663-9403789	WT	MT	OK	0	0,877865	1.79769e+308	1.79769e+308	3.2692e-08	3.50131e-07	yes
AT1G27570	1:9575425-9582376	WT	MT	OK	0	2,0829	1.79769e+308	1.79769e+308	9.76039e-06	6.647e-05	yes
AT1G60095	1:22159735-22162419	WT	MT	OK	0	0,688588	1.79769e+308	1.79769e+308	9.95901e-08	9.84992e-07	yes
AT1G03020	1:698206-698515	WT	MT	OK	0	1,78859	1.79769e+308	1.79769e+308	0,00913915	0,0277958	yes
AT1G13609	1:4662720-4663471	WT	MT	OK	0	3,55814	1.79769e+308	1.79769e+308	0,00021683	0,00108079	yes
AT1G21550	1:7553100-7553876	WT	MT	OK	0	0,562868	1.79769e+308	1.79769e+308	0,00115582	0,00471497	yes
AT1G22120	1:7806308-7809632	WT	MT	OK	0	0,617354	1.79769e+308	1.79769e+308	2.48392e-06	1.91089e-05	yes
AT1G31370	1:11238297-11239363	WT	MT	OK	0	1,46254	1.79769e+308	1.79769e+308	4.83523e-05	0,00028514	3 yes
APUM10	1:13253397-13255570	WT	MT	OK	0	0,581031	1.79769e+308	1.79769e+308	7.87855e-06	5.46603e-05	yes
AT1G48700	1:18010728-18012871	WT	MT	OK	0	0,556525	1.79769e+308	1.79769e+308	6.53917e-05	0,00037473	6 yes
AT1G59077	1:21746209-21833195	WT	MT	OK	0	138,886	1.79769e+308	1.79769e+308	0,00122789	0,00496816	yes
AT1G60050	1:22121549-22123702	WT	MT	OK	0	0,370087	1.79769e+308	1.79769e+308	0,00117953	0,0048001	yes
AT4G15242	4:8705786-8706997	WT	MT	OK	0,00930712	17,9056	10,9098	-4,40523	1.05673e-05	7.13983e-05	yes
AT5G33251	5:12499071-12500433	WT	MT	OK	0,0498375	52,2837	10,0349	-9,8119	0	0	yes
AT4G12520	4:7421055-7421738	WT	MT	OK	0,0195111	15,8516	9,66612	-3,90043	9.60217e-05	0,000528904	yes
AT1G60020	1:22100651-22105276	WT	MT	OK	0,0118377	7,18823	9,24611	-7,50382	6.19504e-14	1.4988e-12	yes
AT5G15360	5:4987235-4989182	WT	MT	OK	0,0988273	56,4834	9,1587	-10,4392	0	0	yes


Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis
 - Quantitative analysis of gene expression
 - DNA and protein chips
 - Next generation transcriptional profiling
- Regulation of gene expression in the identification of gene function by gain-of-function approaches
 - T-DNA activation mutagenesis

Gain-of-Function Approaches


- Methods for identification of gene function using gain-of-function approaches
 - T-DNA activation mutagenesis
 - Method enabling isolation of dominant mutants by random insertion of constitutive promoter, resulting in overexpression of the gene and therefore in corresponding phenotypic changes
 - First step: preparation of mutant library prepared by transformation of a strong constitutive promoter or enhancer
 - Next step: search of interesting phenotypes
 - Identification of the affected gene, e.g. by plasmid-rescue

Activation Mutagenesis




Isolation of *CK11* gene

- Tatsuo Kakimoto, *Science* 274 (1996), 982-985 *
- Isolation of the gene using activation mutagenesis



- Mutant phenotype is a phenocopy of exogenous application of cytokinins (*CK11*, *CYTOKININ INDEPENDENT 1*)




Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis
 - Quantitative analysis of gene expression
 - DNA and protein chips
 - Next generation transcriptional profiling
- Regulation of gene expression in the identification of gene function by gain-of-function approaches
 - T-DNA activation mutagenesis
 - Ectopic expression and regulatable gene expression systems


Regulated Expression Systems

- Regulatable gene expression systems
 - Time- or site-specific regulation of gene expression, leading to a change in phenotype and thereby identification of the natural function of the gene
 - pOP system


Regulated Expression Systems



Regulated Expression Systems





Regulated Expression Systems



Regulated Expression Systems

- Regulatable gene expression systems
 - Time- or site-specific regulation of gene expression, leading to a change in phenotype and thereby identification of the natural function of the gene
 - pOP system
 - UAS system



Outline

- Methods of gene expression analysis
 - Qualitative analysis of gene expression
 - Preparation of transcriptional fusion of promoter of analysed gene with a reporter gene
 - Preparation of translational fusion of the coding region of the analysed gene with reporter gene
 - Use of the data available in public databases
 - Tissue- and cell-specific gene expression analysis
 - Quantitative analysis of gene expression
 - DNA and protein chips
 - Next generation transcriptional profiling
- Regulation of gene expression in the identification of gene function by gain-of-function approaches
 - T-DNA activation mutagenesis
 - Ectopic expression and regulatable gene expression systems
- **Chemical genetics**

Chemical Genetics

- New trends
 - Chemical genetics
 - „chemical genetics“ – more than **50.000/89.631** records in PubMed database (16.10. **2008/29.10. 2015**, an increase of **65 %**)

The screenshot displays the PubMed search interface. At the top, the NCBI logo is on the left, and the PubMed logo is in the center, with the text "A service of the U.S. National Library of Medicine and the National Institutes of Health" and "www.pubmed.gov" below it. On the right, there are links for "NCBI" and "SIGN IN | REGISTER". Below the header, a navigation bar includes "All Databases", "PubMed", "Nucleotide", "Protein", "Genome", "Structure", "OJPM", "PMC", and "Journals". The search bar contains "PubMed" and "for chemical genetics", with "Go" and "Clear" buttons. There are also links for "Advanced Search (beta)" and "Save Search". Below the search bar, there are buttons for "Limits", "Preview/Index", "History", "Clipboard", and "Details". The "Display" section shows "Summary" selected, "Show 20", "Sort By", and "Send to". The results section shows "All: 50407" and "Review: 5562". The first three results are listed:

- 1: [Luchowski E, Matheva EG, Grysinski J, Isperichig EA, Patankar L, Lucio G, Bomilo J, Grysinski Z.](#) [Refiled Articles](#)
Single Molecule Studies of Multiple-Fluorophore Labeled Antibodies. Effect of Homo-FRET on the Number of Photons Available Before Photobleaching. *Curr Pharm Biotechnol*. 2008 Oct;9(3):411-20. PMID: 18833493 [PubMed - in process]
- 2: [Kobayashi M, Ito Y, Ito Y, Shima H.](#) [Refiled Articles, Links](#)
Five cases of beta-ureidopropionase deficiency detected by GC/MS analysis of urine metabolome. *J Mass Spectrom*. 2008 Oct 14. [Epub ahead of print] PMID: 18833477 [PubMed - as supplied by publisher]
- 3: [Zhou M, Peng Z, Hsu-Lyly P, Wu H.](#) [Refiled Articles, Links](#)
A conserved C-terminal thirteen amino acid motif of Gap1 is required for the Gap1 function and necessary for biogenesis of a serine-rich glycoprotein of *Streptococcus parasanguinis*. *Infect Immun*. 2008 Oct 13. [Epub ahead of print] PMID: 18832249 [PubMed - as supplied by publisher]


On the right side, there is a "Recent Activity" box with a search icon and the text "cism.kalozsika (50407) PubMed".

Chemical Genetics

- New trends
 - Chemical genetics
 - „**chemical genetics**“ – more than **50.000/89.631** records in PubMed database (16.10. **2008**/29.10. **2015**, **an increase of 65 %**)
 - Like in the case of genetics, there are also „**forward**“ and „**reverse**“ genetics approaches
 - Unlike in „classical“ genetics approaches, **the subject of study** is not a gene, but a **protein**
 - Chemical genetics tries to identify either the **target protein** after a chemical treatment and after following phenotypic changes („**forward**“ **chemical genetics**) or **chemicals able to interact with protein of interest** („**reverse**“ **chemical genetics**)
 - For that purpose there are carried out **searches in the libraries** of various **chemicals** (thousands of entries, commercially available)
 - example: **analysis of endomembrane transport** in plants



Chemical Genetics



- Analysis of mechanisms of endomembrane transport by chemical genetics approaches
 - In plants cells there occur very dynamic processes mediated mainly by endomembrane transport (see film, GFP targeting to the ER)



Chemical Genetics

- Analysis of mechanisms of endomembrane transport by chemical genetics approaches
 - In plants cells there occur very dynamic processes mediated mainly by endomembrane transport (see film, GFP targeting to the ER)
 - Endomembrane transport is an important regulatory mechanism in signal transduction and regulation of cellular processes





Richter et al., *E J Cell Biol* (2010)

In the figure, there is simplified scheme of vesicle trafficking pathways, regulated by GNOM and its closest relative, GNOM-LIKE1 (GNL1).

Secretory and membrane proteins are synthesised at the ER (blue) and passed onto the Golgi apparatus (green) by anterograde trafficking in COPII-coated vesicles.

The retrograde route from the Golgi apparatus to the ER is regulated by the ARF-GEFs GNOM (GN) and GNL1, which regulate the recruitment of COPI coats to the Golgi membrane. On the secretory route, proteins are transported to the sorting station, the trans-Golgi network (TGN; lilac).

From there, proteins are either transported to the vacuole (grey) via multivesicular bodies (MVB, also called prevacuolar compartment, PVC, which corresponds to the late endosome; deep blue) or trafficked to the plasma membrane (PM).


Plasma membrane proteins like the auxin efflux carrier PIN1 (red), which accumulates at the basal PM at steady state, are continually internalised and trafficked to the TGN, which resembles the early endosome (EE) in plants.

From the TGN, PIN1 is recycled to the plasma membrane via the recycling endosome (RE; light blue). This pathway is regulated by the ARF-GEF GNOM.

Chemical Genetics


■ Analysis of mechanisms of endomembrane transport by chemical genetics approaches

- By searching in the „library“ of chemicals there were identified those, that lead to the secretion of enzyme (carboxypeptidase Y) in yeast (*S. cerevisiae*) – this enzyme is normally transported to the vacuole via the endomembrane transport



- Analysis of changes in secretion using dot-blot and immunodetection of carboxypeptidase Y in the culture medium with monoclonal antibodies

Chemical structure of sortins




0 2.5 5 10 25 50 100 [mg/L]

Immunodetection of carboxypeptidase




Detection of vacuole phenotype (tonoplast shape) of yeast by staining with a specific color (MDY-64)



Chemical Genetics

■ Analysis of mechanisms of endomembrane transport by chemical genetics approaches

- By searching in the „library“ of chemicals there were identified those, that lead to the secretion of enzyme (carboxypeptidase Y) in yeast (*S. cerevisiae*) – this enzyme is normally transported to the vacuole via the endomembrane transport





- Analysis of changes in secretion using dot-blot and immunodetection of carboxypeptidase Y in the culture medium with monoclonal antibodies

- Identified compounds („sortins“) were able to induce similar changes in *Arabidopsis* as well – transport mechanisms are conserved in yeast and in plants

- For detailed identification of the molecular process affected by one of the identified „sortins“, the analysis of its influence on a secretion of a marker protein (AtCPY) was performed – sortin 1 specifically inhibits only this secretory pathway

- Identification of mutants with altered sensitivity to sortin 1 (hyper- or hypo-sensitive mutants) by EMS mutagenesis

Shape of plant vacuoles using EGFP:-TIP



Phenotype of seedlings in the presence of sortins

Chemical Genetics

- Analysis of mechanisms of endomembrane transport by chemical genetics approaches – **summary**

- GFP::d-TIP vacuole membrane (tonoplast) labelling and **identification of mutations leading to altered tonoplast morphology**
- **Chemical genetics in combination with classical genetics** – identification of proteins participating in regulation of endomembrane transport
- **Proteomics approaches** – identification and analysis of vacuole proteome

