

Population Structure

“Populační ekologie živočichů“

Stano Pekár


- ▶ **Demography** - study of organisms with special attention to stage or age structure
- ▶ processes are associated to age, stage or size

x .. age/stage/size category

p_x .. age/stage/size specific survival


$$p_x = \frac{S_{x+1}}{S_x}$$

m_x .. reproductive rate (expected average number of offspring per female)




- ▶ main focus on births and deaths
- ▶ immigration & emigration is ignored
- ▶ no adult survive
- ▶ one (not overlapping) generation per year
- ▶ egg pods over-winter
- ▶ despite high fecundity they just replace themselves

Annual species




- ▶ breed at discrete periods
- ▶ no overlapping generations

Biennial species



- ▶ breed at discrete periods
- ▶ adult generation may overlap

Perins (1965)



Perennial species

- ▶ breed at discrete periods
- ▶ breeding adults consist of individuals of various ages (1-5 years)
- ▶ adults of different generations are equivalent
- ▶ overlapping generations



Parus major

Age-size-stage life-table


► age/stage classification is based on developmental time

► size may be more appropriate than age (fish, sedentary animals)

► Hughes (1984) used combination of age/stage and size for the description of coral growth



Agaricia agaricites



Age-dependent life-tables

- ▶ show organisms' mortality and reproduction as a function of age

Static (vertical) life-tables


- ▶ examination of a population during one segment (time interval)

- segment = group of individuals of different cohorts
- designed for long-lived organisms

- ▶ ASSUMPTIONS:


- Birth rate and survival are constant over time
- population does not grow

- ▶ DRAWBACKS: confuses age-specific changes in e.g. mortality with temporal variation



x	S _x	D _x	l _x	p _x	q _x	m _x
1	129	15	1.000	0.884	0.116	0.000
2	114	1	0.884	0.991	0.009	0.000
3	113	32	0.876	0.717	0.283	0.310
4	81	3	0.628	0.963	0.037	0.280
5	78	19	0.605	0.756	0.244	0.300
6	59	-6	0.457	1.102	-0.102	0.400
7	65	10	0.504	0.846	0.154	0.480
8	55	30	0.426	0.455	0.545	0.360
9	25	16	0.194	0.360	0.640	0.450
10	9	1	0.070	0.889	0.111	0.290
11	8	1	0.062	0.875	0.125	0.280
12	7	5	0.054	0.286	0.714	0.290
13	2	1	0.016	0.500	0.500	0.280
14	1	-3	0.008	4.000	-3.000	0.280
15	4	2	0.031	0.500	0.500	0.290
16	2	2	0.016	0.000	1.000	0.280

Lowe (1969)



us elaphus

S_x .. number of survivors

D_x .. number of dead individuals

$$D_x = S_x - S_{x+1}$$

l_x .. standardised number of survivors

$$l_x = \frac{S_x}{S_0}$$

q_x .. age-specific mortality

$$q_x = \frac{D_x}{S_x}$$

p_x .. age-specific survival

$$p_x = \frac{l_{x+1}}{l_x}$$

Cohort (horizontal) life-table

- ▶ examination of a population in a cohort = a group of individuals born at the same period
- ▶ followed from birth to death
- ▶ provide reliable information
- ▶ designed for short-lived organisms
- ▶ only females are included

x	Sx	Dx	lx	px	qx	mx
0	250	50	1.000	0.800	0.200	0.000
1	200	120	0.800	0.400	0.600	0.000
2	80	50	0.320	0.375	0.625	2.000
3	30	15	0.120	0.500	0.500	2.100
4	15	9	0.060	0.400	0.600	2.300
5	6	6	0.024	0.000	1.000	2.400
6	0	0	0.000			



Vulpes vulpes


Stage or size-dependent life-tables

- ▶ survival and reproduction depend on stage / size rather than age
- ▶ age-distribution is of no interest
- ▶ used for invertebrates (insects, invertebrates)
- ▶ time spent in a stage / size can differ

Campbell (1981)


x	Sx	Dx	lx	px	qx	mx
Egg	450	68	1.000	0.849	0.151	0
Larva I	382	67	0.849	0.825	0.175	0
Larva II	315	158	0.700	0.498	0.502	0
Larva III	157	118	0.349	0.248	0.752	0
Larva IV	39	7	0.087	0.821	0.179	0
Larva V	32	9	0.071	0.719	0.281	0
Larva VI	23	1	0.051	0.957	0.043	0
Pre-pupa	22	4	0.049	0.818	0.182	0
Pupa	18	2	0.040	0.889	0.111	0
Adult	16	16	0.036	0.000	1.000	185

Lymantria dispar




Survivorship curves

- ▶ display change in survival by plotting $\log(l_x)$ against age (x)
- ▶ sheep mortality increases with age
- ▶ survivorship of lapwing (*Vanellus*) is independent of age but survival of sheep is age-dependent



Pearls (1928) classified hypothetical age-specific mortality:

- ▶ Type I .. mortality is concentrated at the end of life span (humans)
- ▶ Type II .. mortality is constant over age (seeds, birds)
- ▶ Type III .. mortality is highest in the beginning of life (invertebrates, fish, reptiles)








Birth rate curves

- ▶ fecundity - potential number of offspring
- ▶ fertility - real number of offspring

- ▶ semelparous .. reproducing once a life
- ▶ iteroparous .. reproducing several times during life

- ▶ birth pulse .. discrete reproduction
(seasonal reproduction)
- ▶ birth flow .. continuous reproduction





Key-factor analysis

- ▶ k-value - **killing power** - another measure of mortality

$$k = -\ln(p)$$

- ▶ k-values are additive unlike q

$$K = \sum k_x$$

- ▶ **Key-factor analysis** - a method to identify the most important factors that regulates population dynamics

- ▶ k-values are estimated for a number of years

- ▶ important factors are identified by regressing k_x on $\log(N)$

Leptinotarsa decemlineata


- ▶ over-wintering adults emerge in June → eggs are laid in clusters on the lower side of leaves → larvae pass through 4 instars → form pupal cells in the soil → summer adults emerge in August → begin to hibernate in September
- ▶ mortality factors overlap



Harcourt (1971)

Age interval	Numbers per 96 potato hills	Numbers 'dying'	'Mortality factor'	$\log_{10} N$	k -value	
Eggs	11 799	2531	Not deposited	4.072	0.105	(k_{1a})
	9268	445	Infertile	3.967	0.021	(k_{1b})
	8823	408	Rainfall	3.946	0.021	(k_{1c})
	8415	1147	Cannibalism	3.925	0.064	(k_{1d})
	7268	376	Predators	3.861	0.024	(k_{1e})
Early larvae	6892	0	Rainfall	3.838	0	(k_2)
Late larvae	6892	3722	Starvation	3.838	0.337	(k_3)
Pupal cells	3170	16	<i>D. doryphorae</i>	3.501	0.002	(k_4)
Summer adults	3154	-126	Sex (52% female)	3.499	-0.017	(k_5)
Female × 2	3280	3264	Emigration	3.516	2.312	(k_6)
Hibernating adults	16	2	Frost	1.204	0.058	(k_7)
Spring adults	14			1.146	2.926	(k_{total})

Summary over 10 years



- ▶ highest k-value indicates the role of a factor in each generation
- ▶ profile of a factor parallel with the K profile reveals the key factor
- ▶ emigration is the key-factor