

# MACROEVOLUTION






Fig. 161. Pelvis of Archaeopteryx.






Fig. 162. Pelvis of Apatornis.



Phyletic Gradualism




Morphology




Punctuated Equilibrium

Time



# RATE OF EVOLUTION PUNCTUATED EQUILIBRIA

Rate of evolution:



difference between trait values in time  $t_2$  and  $t_1$

Haldane (1949)

$$r = \frac{\ln x_2 - \ln x_1}{\Delta}$$

time interval  
 $t_2 - t_1$


1 darwin = e-fold change in a trait over 1 million years

G. G. Simpson:

evoluce bradytelic (slow)

horotelic (standard, eg. horses)

tachytelic (fast)



Haldane (1949): Tertiary horses – 0.04 darwins  
domestication –  $10^3$  darwins

Kuertén (1959): Holocene mammals – 12.6 darwins  
Pleistocene mammals – 0.5 darwins  
Tertiary mammals – 0.02 darwins

... differences caused by different time intervals

disadvantages: 1. factor  $e$  is not biologically natural

2. uses absolute time

3. does not take into account measured time interval



4. impossible to compare areas/volumes/linear dimensions

⇒ Haldane (1949), Gingerich (1993): 1 haldane = change measured in units of standard deviation per generation


# Theory of punctuated equilibria:

Stephen Jay Gould, Niles Eldredge (1972)

stasis vs. rapid change





S.J. Gould



N. Eldredge

# stasis vs. rapid change







## Mechanism?

peripatric speciation



macromutation – R. Goldschmidt, *The Material Basis of Evolution* (1940): „hopeful monsters“

# Peripatric speciation and punctuated equilibria











Species of different genera and classes has not changed by the same rate or to the same degree (see „living fossils“).

Periods during which species were changing were short relative to periods during which they stayed unchanged.

Except of (nonexisting) completely constant rate there is only variable rate – species change either in discrete steps (punctuationism) or gradually. Therefore, stasis is only an extreme case of slow evolution.




R. Dawkins: Blind watchmaker

Punctuational process typical for language evolution:  
changes play an important role in periods of divergence of a new language  
Bantu, Indo-European and Austronesian group: 10-33% differences  
connected with language splitting

### How to explain stasis?

genetic or ontogenetic constraints

habitat tracking – glacial/interglacial cycles



short-term local divergence – rapid changes spatially limited

# Relation between micro- and macroevolution

Steven M. Stanley (1975): macroevolution separated from microevolution

S.J. Gould (1980): „deposition of neo-Darwinism from the throne“, „effective death of neo-Darwinism“

Modern Synthesis narrow, extrapolationistic and reductionistic

Is macroevolution really different from microevolution?

evolution of horses

Darwin's finches

mammal evolution



evolution as „matryoshka“



# Evolution of horses:

2 dental dimensions

mean rate can be explained by acting of directional selection  
(sufficient 2 selective deaths/million of individuals/generation)


if  $N_e < 10^4$  individuals, can be explained by drift alone

likewise also other fossils



# Darwin's finches:

with known age of Galapagos enough time for diversification to 14 species  
(in fact more complicated – reversions, possible extinction of some species)




(b) The Galapagos finches

# Evolution of mammals from therapsid reptiles:

changes gradual


large differences between reptiles and mammals are adaptive in individual links  $\Rightarrow$  same mechanisms as in microevolution



# Relation of macroevolution and ontogeny

J. F. Meckel, E. Serres: embryos display traits of embryos of species preceding on the *Scala Naturae*

Ernst Haeckel – **biogenetic law** (= **recapitulation I.**): ontogeny recapitulates phylogeny (eg. gills during mammal embryonal development)





# RECAPITULATION IN MAN (?)






Fig. 1

× specialized larval forms (= non-terminal addition): zoëa of crabs, Müller's larva of echinoderms, caterpillar of butterflies etc.

terminal vs. non-terminal addition



# Karl Ernst von Baer – embryological laws:



Vertebrate embryos pass through stable stages during their development which are not identical to any animals species.

Embryos of related species are similar to each other and dissimilar to adults of ancestral species.

Developmental stages



Stage 1

Stage 2

Stage 3

Stage 4



Amphibian



Reptile




Mammal

Stage 1 embryos are most similar to each other

Stage 4 embryos are least similar to one another




# Karl Ernst von Baer – embryological laws:

1st law: General traits of a large animal group appear in the embryo earlier than special traits (eg. cartilage in bony fishes).



# General principles of ontogeny and evolution:



modularization and individualization: serial homology and homonymy



## General principles of ontogeny and evolution:

heterotopy = change of the position of a trait phenotypic expression  
(eg. photosynthesis in succulent stem; sesamoid bones – *patella*,  
ossified tendons in dinosaur tails, „panda’s thumb“)

■ Human thumb bones actually correspond to the sixth finger on the panda hand.




General principles of ontogeny and evolution:

heterochrony and allometry

## Heterochrony

<http://www.bio.miami.edu/dana/dox/heterochrony.html>



### Allometric Growth

Differential growth in organs and body parts



### Paedomorphy

Retention of juvenile structures

## Heterochrony:

		Somatic traits	Reproductive org.
peramorphosis			
paedomorphosis			


= change of timing of ontogenetic events:

1. speed of the process
2. timing of the process




# Heterochrony:

		Somatic traits	Reproductive org.
peramorphosis	hypermorphosis	--	deceleration
	acceleration	acceleration	--
paedomorphosis			



acceleration →  
peramorphosis



hypermorphosis →  
peramorphosis



hypermorphosis




*Megaceros giganteus*

# Heterochrony:

		Somatic traits	Reproductive org.
peramorphosis	hypermorphosis	--	deceleration
	acceleration	acceleration	--
paedomorphosis	progenesis	--	acceleration
	neoteny	deceleration	--



neoteny →  
pedomorphosis



progenesis →  
pedomorphosis

neoteny:




*Ambystoma mexicanum*





# Birds have paedomorphic dinosaur skulls

Bhart-Anjan S. Bhullar<sup>1</sup>, Jesús Marugán-Lobón<sup>2</sup>, Fernando Racimo<sup>1</sup>, Gabe S. Bever<sup>3</sup>, Timothy B. Rowe<sup>4</sup>, Mark A. Norell<sup>5</sup> & Arhat Abzhanov<sup>1</sup>



**Figure 4 | Similarity of embryonic *Alligator* and adult *Confuciusornis* skulls.** Superimposition of *Alligator* embryo skull (green) onto *Alligator* adult skull (red, left) and onto *Confuciusornis* adult skull (red, right), showing the nearly identical skull configuration of the latter two and indicating paedomorphic cranial morphology in *Confuciusornis*.







**Figure 5 | Summary of heterochrony and phylogeny in bird skull evolution.** A phylogenetic sequence with skull outlines set on deformation grids is depicted from the primitive stem-group archosaur *Euparkeria* to the modern



emu *Dromaius*. Heterochronic transformations referred to in the text are enumerated with Roman numerals. Major anatomical regions involved in heterochronic transformations are labelled.

# Neoteny in humans?

a)



b)



# Human neotenic features compared to the chimp<sup>\*)</sup> (Wikipedia):

## Head:

- rounded skull
- slender cranial bones
- reduced brow ridges
- large brain
- flat face
- broadened face
- hairless face
- hairs on top head
- large eyes
- ear shape
- small nose
- small teeth
- small upper and lower jaw



## Genitals:

- absence of baculum (*os penis*)
- presence of hymen
- anteriorly oriented vagina

## Limbs/posturer:

- legs longer than hands
- foot structure
- upright posture

„Naked“ body

<sup>\*)</sup> some of them are not, in fact, neotenic!

## Origin of macroevolutionary novelties:

change of function of a gene product:

pigment producing enzyme → change of coloration

digestive enzyme → change of sexual habits

loss of function:

genes suppressing own pathogenicity

deletion of host proteins recognized by parasites (eg. CCR5-Δ32

deletion in the *CCR5* gene → resistancy to the HIV and variola

.... 5-14% of Europeans, in Africans and Asiatics rare)

changes in gene regulation

prions – incorrect translation termination ⇒ bovine spongiform

encephalopathy,

scrapie of sheep and goats, kuru, Creutzfeld-Jakob disease in humans

role of gene duplications – more radical changes enabled


symbiosis, gene transfer (retroviruses)

homeotic genes



# Homeotic (*Hox*) genes

**William Bateson:** „homeosis“ = anatomical changes of large extent (eg. development of an extra finger, cervical vertebra instead of thoracic, limb in ectopic position)




# Homeotic (*Hox*) genes

Edward Lewis: **homeotic genes** = genes responsible for basic segmentation of multicellular animals – homeotic mutations do not change the number of segments but their identity



control of transcription of other genes (eg. *Ubx* probably regulates hundreds of „target“ genes)

determination of basic body segmentation

high evolutionary conservativeness



# Homeotic mutation





*Antennapedia*

mutation of the  
*Ultrabithorax* gene:  
3rd thoracic segment  
(T3) → T2

*Bithorax*

# Hox genes: basic antero-posterior body segmentation

linear clusters, same order as the segments



Hox gene effects are overlapping


*Drosophila*: 1 linkage group, 2 clusters:

*Antennapedia* (ANT-C)


*Bithorax* (BX-C)

vertebrates:

4 linkage groups








Mouse embryo



*Drosophila*


člověk





Homeobox: 180 bp → homeodomain

60 AA (expression regulation)





# Hox-genes are highly conservative

A

Scr group





Fruit fly	TKRQRTSYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKLKEH
Grasshopper	TKRQRTSYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEH
Beach hopper	TKRQRTSYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEH
Centipede	TKRQRTSYTRYQTLELEKEFHFNRYLTRRRRIEIAHSLCLSERQIKIWFQNRRMKWKKEH
Mite	TKRQRTSYTRYQTLELEKEFHFNRYLTRRRRIEIAHSLCLSERQIKIWFQNRRMKWKKEH
Leech	NKRTRTSYTRHQTLELEKEFHFNRYLSRRRRIEIAHVNLNLSERQIKIWFQNRRMKWKKDH
Sea urchin	SKRSRTAYTRYQTLELEKEFHFNRYLTRRRRIEIAHALGLTERQIKIWFQNRRMKWKKEH
Zebra fish	GKRARTAYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLSERQIKIWFQNRRMKWKKDN
Mouse	GKRARTAYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLSERQIKIWFQNRRMKWKKDN
Human	GKRARTAYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLSERQIKIWFQNRRMKWKKDN

Antp group


Fruit fly	RKRGRQTYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEN
Grasshopper	RKRGRQTYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEN
Beach hopper	RKRGRQTYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEN
Centipede	RKRGRQTYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEN
Spider	RKRGRQTYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEN
Leech	QKRTRQTYTRYQTLELEKEFYSNRYLTRRRRIEIAHSLALSERQIKIWFQNRRMKWKKEN
Sea urchin	GKRGRQTYTRQQTLELEKEFHFVTRRRRFEIAQSLGLSERQIKIWFQNRRMKWKREH
Zebra fish	GRRGRQTYTRYQTLELEKEFHFNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKEN
Mouse	GRRGRQTYTRYQTLELEKEFHYNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKES
Human	GRRGRQTYTRYQTLELEKEFHYNRYLTRRRRIEIAHALCLTERQIKIWFQNRRMKWKKES

# ParaHox genes


## MADS-box genes in plants




# 3 hypotheses of the origin of dipteran arrangement:



## Ubq protein expression:



original function:  
venation development



Ancestral insect  
hindwing

Promotes vein  
development




Dipteran  
haltere




Lepidopteran  
hindwing

loss of  
function ...

... but origin of  
halteres



Creates balloon  
shape



Promotes vein  
development




Controls scale  
morphology  
and color

in butterflies also  
scale morphology  
and wing coloration

# Evolution of thoracic segments in crustaceans – shift of anteroposterior boundary of expression of the *Ubx* gene:



thoracic segments: copepods – 6, lobster – 8, brine shrimps – 11 (ancestral)

movement → maxillipedes






shift of anteroposterior boundary of *Ubx* gene expression = position of transition of locomotory segments and maxillipedes

např. opossum shrimps: 2nd segment, prawns: 4th segment  
2nd thoracic limb of opossum shrimps = transitional segment between 1st (maxillipede) and 3rd limb (locomotion)



# Besides transcriptional factors also regulatory enhancers:



## Activators

The regulatory proteins bind to DNA at distant sites known as enhancers. When DNA folds so that the enhancer is brought into proximity with the transcription complex, the activator proteins interact with the complex to increase the rate of transcription.

## Repressors


These regulatory proteins bind to "silencer sites" on the DNA preventing the binding of activator to nearby enhancers and so slowing transcription.

## Basal factors

These transcription factors, in response to coactivators, position RNA polymerase at the start of a protein-coding sequence, and then release the polymerase to transcribe the mRNA

## Coactivators

These transcription factors transmit signals from activator proteins to the basal factors.



# Macroevolutionary trends species selection

trends: real  $\times$  passive (eg. wall effect)


Edward Drinker Cope: trend to bigger size

**Species selection:**



= preferential survival or proliferation of species

different extinction rates

(a) Differential extinction rate




Neutral  
Stabilizing selection



different speciation rates

(b) Differential speciation rate






the trait connected with differential survival or speciation

these features independent of natural selection

trait is heritable during speciation

SS favours only nonadaptive trends  
(otherwise = natural selection)



**Necessary to prove:**

higher speciation rate/lower extinction rate in lineages which deviate from the average in the direction of the trend

the trend and distribution of varied speciation/extinction rates are not caused by shift in fossil record

the trend and distribution of varied speciation/extinction rates are not caused by natural selection