

# Tutorial 5—Global Analysis

26.10.2021

1. Suppose  $M = \mathbb{R}^3$  with standard coordinates  $(x, y, z)$ . Consider the vector field

$$\xi(x, y, z) = 2\frac{\partial}{\partial x} - \frac{\partial}{\partial y} + 3\frac{\partial}{\partial z}.$$

How does this vector field look like in terms of the coordinate vector fields associated to the cylindrical coordinates  $(r, \phi, z)$ , where  $x = r \cos \phi$ ,  $y = r \sin \phi$  and  $z = z$ ? Or with respect to the spherical coordinates  $(r, \phi, \theta)$ , where  $x = r \sin \theta \cos \phi$ ,  $y = r \sin \theta \sin \phi$  and  $z = r \cos \theta$ ?

2. Consider  $\mathbb{R}^3$  with coordinates  $(x, y, z)$  and the vector fields

$$\xi(x, y, z) = (x^2 - 1)\frac{\partial}{\partial x} + xy\frac{\partial}{\partial y} + xz\frac{\partial}{\partial z}$$

$$\eta(x, y, z) = x\frac{\partial}{\partial x} + y\frac{\partial}{\partial y} + 2xz^2\frac{\partial}{\partial z}.$$

Are they tangent to the cylinder  $M = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 1\} \subset \mathbb{R}^3$  with radius 1 (i.e. do they restrict to vector fields on  $M$ )?

3. Suppose  $M = \mathbb{R}^2$  with coordinates  $(x, y)$ . Consider the vector fields  $\xi(x, y) = y\frac{\partial}{\partial x}$  and  $\eta(x, y) = \frac{x^2}{2}\frac{\partial}{\partial y}$  on  $M$ . We computed in class their flows and saw that they are complete. Compute  $[\xi, \eta]$  and its flow? Is  $[\xi, \eta]$  complete?

4. Let  $M$  be a (smooth) manifold and  $\xi, \eta \in \mathfrak{X}(M)$  two vector fields on  $M$ . Show that

(a)  $[\xi, \eta] = 0 \iff (\text{Fl}_t^\xi)^*\eta = \eta$ , whenever defined  $\iff \text{Fl}_t^\xi \circ \text{Fl}_s^\eta = \text{Fl}_s^\eta \circ \text{Fl}_t^\xi$ , whenever defined.

(b) If  $N$  is another manifold,  $f : M \rightarrow N$  a smooth map, and  $\xi$  and  $\eta$  are  $f$ -related to vector fields  $\tilde{\xi}$  resp.  $\tilde{\eta}$  on  $N$ , then  $[\xi, \eta]$  is  $f$ -related to  $[\tilde{\xi}, \tilde{\eta}]$ .

5. Consider the general linear group  $\text{GL}(n, \mathbb{R})$ . For  $A \in \text{GL}(n, \mathbb{R})$  denote by

$$\lambda_A : \text{GL}(n, \mathbb{R}) \rightarrow \text{GL}(n, \mathbb{R}) \quad \lambda_A(B) = AB$$

$$\rho_A : \text{GL}(n, \mathbb{R}) \rightarrow \text{GL}(n, \mathbb{R}) \quad \rho_A(B) = BA$$

left respectively right multiplication by  $A$ , and by  $\mu : \text{GL}(n, \mathbb{R}) \times \text{GL}(n, \mathbb{R}) \rightarrow \text{GL}(n, \mathbb{R})$  the multiplication map.

(a) Show that  $\lambda_A$  and  $\rho_A$  are diffeomorphisms for any  $A \in \mathbf{GL}(n, \mathbb{R})$  and that

$$T_B \lambda_A(B, X) = (AB, AX) \quad T_B \rho_A(B, X) = (BA, XA),$$

where  $(B, X) \in T_B \mathbf{GL}(n, \mathbb{R}) = \{(B, X) : X \in M_n(\mathbb{R})\}$ .

(b) Show that

$$T_{(A,B)} \mu((A, B), (X, Y)) = T_B \lambda_A Y + T_A \rho^B X = (AB, AY + XB)$$

where  $(A, B) \in \mathbf{GL}(n, \mathbb{R}) \times \mathbf{GL}(n, \mathbb{R})$  and  $(X, Y) \in M_n(\mathbb{R}) \times M_n(\mathbb{R})$ .

(c) For any  $X \in M_n(\mathbb{R}) \cong T_{Id} \mathbf{GL}(n, \mathbb{R})$  consider the maps

$$L_X : \mathbf{GL}(n, \mathbb{R}) \rightarrow T\mathbf{GL}(n, \mathbb{R}) \quad L_X(B) = T_{Id} \lambda_B(Id, X) = (B, BX).$$

$$R_X : \mathbf{GL}(n, \mathbb{R}) \rightarrow T\mathbf{GL}(n, \mathbb{R}) \quad R_X(B) = T_{Id} \rho_B(Id, X) = (B, XB).$$

Show that  $L_X$  and  $R_X$  are smooth vector field and that  $\lambda_A^* L_X = L_X$  and  $\rho_A^* R_X = R_X$  for any  $A \in \mathbf{GL}(n, \mathbb{R})$ . What are their flows? Are these vector fields complete?

(d) Show that  $[L_X, R_Y] = 0$  for any  $X, Y \in M_n(\mathbb{R})$ .