

## Tutorial 7-8—Global Analysis

1. Suppose  $E \rightarrow M$  is a (smooth) vector bundle of rank  $k$  over a manifold  $M$ . Then  $E$  is called *trivializable*, if it is isomorphic to the trivial vector bundle  $M \times \mathbb{R}^k \rightarrow M$ .
  - (a) Show that  $E \rightarrow M$  is trivializable  $\iff E \rightarrow M$  admits a global frame, i.e. there exist (smooth) sections  $s_1, \dots, s_k$  of  $E$  such that  $s_1(x), \dots, s_k(x)$  span  $E_x$  for any  $x \in M$ .
  - (b) Show that the tangent bundle of any Lie group  $G$  is trivializable.
  - (c) Recall that  $\mathbb{R}^n$  has the structure of a (not necessarily associative) division algebra over  $\mathbb{R}$  for  $n = 1, 2, 4, 8$ . Use this to show that the tangent bundle of the spheres  $S^1 \subset \mathbb{R}^2$ ,  $S^3 \subset \mathbb{R}^4$  and  $S^7 \subset \mathbb{R}^8$  is trivializable.
  
2. Let  $V$  be a finite dimensional real vector space and consider the subspace of  $r$ -linear alternating maps  $\Lambda^r V^* = L_{\text{alt}}^r(V, \mathbb{R})$  of the vector space of  $r$ -linear maps  $L^r(V, \mathbb{R}) = (V^*)^{\otimes r}$ . Show that for  $\omega \in L^r(V, \mathbb{R})$  the following are equivalent:
  - (a)  $\omega \in \Lambda^r V^*$
  - (b) For any vectors  $v_1, \dots, v_r \in V$  one has
 
$$\omega(v_1, \dots, v_i, \dots, v_j, \dots, v_k) = -\omega(v_1, \dots, v_j, \dots, v_i, \dots, v_k)$$
  - (c)  $\omega$  is zero whenever one inserts a vector  $v \in V$  twice.
  - (d)  $\omega(v_1, \dots, v_k) = 0$ , whenever  $v_1, \dots, v_k \in V$  are linearly dependent vectors.
  
3. Let  $V$  be a finite dimensional real vector space. Show that the vector space  $\Lambda^* V^* := \bigoplus_{r \geq 0} \Lambda^r V^*$  is an associative, unital, graded-anticommutative algebra with respect to the wedge product  $\wedge$ , i.e. show that the following holds:
  - (a)  $(\omega \wedge \eta) \wedge \zeta = \omega \wedge (\eta \wedge \zeta)$  for all  $\omega, \eta, \zeta \in \Lambda^* V^*$ .
  - (b)  $1 \in \mathbb{R} = \Lambda^0 V^*$  satisfies  $1 \wedge \omega = \omega \wedge 1 = \omega$  for all  $\omega \in \Lambda^* V^*$ .
  - (c)  $\Lambda^r V^* \wedge \Lambda^s V^* \subset \Lambda^{r+s} V^*$ .
  - (d)  $\omega \wedge \eta = (-1)^{rs} \eta \wedge \omega$  for  $\omega \in \Lambda^r V^*$  and  $\eta \in \Lambda^s V^*$ .

Moreover, show that for any linear map  $f : V \rightarrow W$  the linear map  $f^* : \Lambda^* W^* \rightarrow \Lambda^* V^*$  is a morphism of graded unital algebras, i.e.  $f^* 1 = 1$ ,  $f^*(\Lambda^r W^*) \subset \Lambda^r V^*$  and  $f^*(\omega \wedge \eta) = f^* \omega \wedge f^* \eta$ .

4. Let  $V$  be a finite dimensional real vector space. Show that:

(a) If  $\omega_1, \dots, \omega_r \in V^*$  and  $v_1, \dots, v_r \in V$ , then

$$\omega_1 \wedge \dots \wedge \omega_r(v_1, \dots, v_r) = \det((\omega_i(v_j))_{1 \leq i, j \leq r}).$$

In particular,  $\omega_1, \dots, \omega_r$  are linearly independent  $\iff \omega_1 \wedge \dots \wedge \omega_r \neq 0$ .

(b) If  $\{\lambda_1, \dots, \lambda_n\}$  is a basis of  $V^*$ , then

$$\{\lambda_{i_1} \wedge \dots \wedge \lambda_{i_r} : 1 \leq i_1 < \dots < i_r \leq n\}$$

is a basis of  $\Lambda^r V^*$ .

5. Let  $V$  be a finite dimensional real vector space. An element  $\mu \in L^r(V, \mathbb{R})$  is called *symmetric*, if  $\mu(v_1, \dots, v_r) = \mu(v_{\sigma(1)}, \dots, v_{\sigma(r)})$  for any vectors  $v_1, \dots, v_r \in V$  and any permutation  $\sigma \in S^r$ . Denote by  $S^r V^* \subset L^r(V, \mathbb{R})$  the subspace of symmetric elements in the vector space  $L^r(V, \mathbb{R})$ .

(a) For  $\mu \in L^r(V, \mathbb{R})$  show that

$$\mu \in S^r V^* \iff \mu(v_1, \dots, v_i, \dots, v_j, \dots, v_k) = \mu(v_1, \dots, v_j, \dots, v_i, \dots, v_k),$$

for any vectors  $v_1, \dots, v_r \in V$ .

(b) Consider the map  $\text{Sym} : L^r(V, \mathbb{R}) \rightarrow L^r(V, \mathbb{R})$  given by

$$\text{Sym}(\mu)(v_1, \dots, v_r) = \frac{1}{r!} \sum_{\sigma \in S^r} \mu(v_{\sigma(1)}, \dots, v_{\sigma(r)}).$$

Show that  $\text{Image}(\text{Sym}) = S^r V^*$  and that  $\mu \in S^r V^* \iff \text{Sym}(\mu) = \mu$ .

6. Let  $V$  be a finite dimensional real vector space and set  $S(V^*) := \bigoplus_{r=0}^{\infty} S^r V^*$  with the convention  $S^0 V^* = \mathbb{R}$  and  $S^1 V^* = V^*$ . For  $\mu \in S^r V^*$  and  $\nu \in S^t V^*$  define their symmetric product by

$$\mu \odot \nu := \text{Sym}(\mu \otimes \nu) \in S^{r+t} V^*.$$

By bilinearity, we extend this to a  $\mathbb{R}$ -bilinear map  $\odot : S(V^*) \times S(V^*) \rightarrow S(V^*)$ . Show that  $S(V^*)$  is an unital, associative, commutative, graded algebra with respect to the symmetric product  $\odot$ .

7. Suppose  $p : E \rightarrow M$  and  $q : F \rightarrow M$  are vector bundles over  $M$ . Show that their direct sum  $E \oplus F := \sqcup_{x \in M} E_x \oplus F_x \rightarrow M$  and their tensor product  $E \otimes F := \sqcup_{x \in M} E_x \otimes F_x \rightarrow M$  are again vector bundles over  $M$ .

8. Suppose  $E \subset TM$  is a smooth distribution of rank  $k$  on a manifold  $M$  of dimension  $n$  and denote by  $\Omega(M)$  the vector space of differential forms on  $M$ .

(a) Show that locally around any point  $x \in M$  there exists (local) 1-forms  $\omega^1, \dots, \omega^{n-k}$  such that for any (local) vector field  $\xi$  one has:  $\xi$  is a (local) section of  $E \iff \omega_i(\xi) = 0$  for all  $i = 1, \dots, n - k$ .

- (b) Show that  $E$  is involutive  $\iff$  whenever  $\omega^1, \dots, \omega^{n-k}$  are local 1-forms as in (a) then there exists local 1-forms  $\mu^{i,j}$  for  $i, j = 1, \dots, n-k$  such that

$$d\omega^i = \sum_{j=1}^{n-k} \mu^{i,j} \wedge \omega^j.$$

- (c) Show

$$\Omega_E(M) := \{\omega \in \Omega(M) : \omega|_E = 0\} \subset \Omega(M)$$

is an ideal of the algebra  $(\Omega(M), \wedge)$ . Here,  $\omega|_E = 0$  for a  $\ell$ -form  $\omega$  means that  $\omega(\xi_1, \dots, \xi_\ell) = 0$  for any sections  $\xi_1, \dots, \xi_\ell$  of  $E$ .

- (d) An ideal  $\mathcal{J}$  of  $(\Omega(M), \wedge)$  is called differential ideal, if  $d(\mathcal{J}) \subset \mathcal{J}$ . Show that  $\Omega_E(M)$  is a differential ideal  $\iff E$  is involutive.

9. Suppose  $M$  is a manifold and  $D_i : \Omega^k(M) \rightarrow \Omega^{k+r_i}(M)$  for  $i = 1, 2$  a graded derivation of degree  $r_i$  of  $(\Omega(M), \wedge)$ .

- (a) Show that

$$[D_1, D_2] := D_1 \circ D_2 - (-1)^{r_1 r_2} D_2 \circ D_1$$

is a graded derivation of degree  $r_1 + r_2$ .

- (b) Suppose  $D$  is a graded derivation of  $(\Omega(M), \wedge)$ . Let  $\omega \in \Omega^k(M)$  be a differential form and  $U \subset M$  an open subset. Show that  $\omega|_U = 0$  implies  $D(\omega)|_U = 0$ .

**Hint:** Think about writing 0 as  $f\omega$  for some smooth function  $f$  and use the defining properties of a graded derivation.

- (c) Suppose  $D$  and  $\tilde{D}$  are two graded derivations such that  $D(f) = \tilde{D}(f)$  and  $D(df) = \tilde{D}(df)$  for all  $f \in C^\infty(M, \mathbb{R})$ . Show that  $D = \tilde{D}$ .

10. Suppose  $M$  is a manifold and  $\xi, \eta \in \Gamma(TM)$  vector fields.

- (a) Show that the insertion operator  $i_\xi : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$  is a graded derivation of degree  $-1$  of  $(\Omega(M), \wedge)$ .

- (b) Recall from class that  $[d, d] = 0$ . Verify (the remaining) graded-commutator relations between  $d, \mathcal{L}_\xi, i_\eta$ :

- (i)  $[d, \mathcal{L}_\xi] = 0$ .
- (ii)  $[d, i_\xi] = d \circ i_\xi + i_\xi \circ d = \mathcal{L}_\xi$ .
- (iii)  $[\mathcal{L}_\xi, \mathcal{L}_\eta] = \mathcal{L}_{[\xi, \eta]}$ .
- (iv)  $[\mathcal{L}_\xi, i_\eta] = i_{[\xi, \eta]}$ .
- (v)  $[i_\xi, i_\eta] = 0$ .

**Hint:** Use (c) from 2.