

Microscopic Analysis of Urine

Authors:

RNDr. Miroslava Beňovská, Ph.D.

Mgr. Ondřej Wiewiorka

MUDr. Jana Tůmová

Specimen selections:

Team of employees in Department of Clinical Biochemistry, University Hospital Brno

Faculty of Medicine, Masaryk University

Department of Laboratory Methods

University Hospital Brno

Department of Clinical biochemistry

Created in collaboration with Service Center for E-Learning at MU,
Faculty of Informatics, Masaryk University, Brno 2014–2016

Tiskový výstup publikace vydané na Elportále MU (<http://elportal.cz/>)
<http://elportal.cz/webs/urine-analysis/>

Introduction

Semiquantitative urine analysis is one of the basic biochemical examinations. It consists of urine chemical properties determination by diagnostic strips and morphological examination of urinary elements. Both methods support each other in final determination of the result and both findings should correspond with each other. Urine analysis is required to be performed in one hour and is conducted from single sampling, preferably from the second urination in the morning. Nowadays, the analysis of urine is mostly automated. Manual microscopy is used only for determination of ambiguous or discrepant results.

The urine sediment for manual microscopy is prepared as follows: native urine sample is centrifuged in 2000 rpm and the supernatant is removed and the sediment resuspended to create a tenfold concentrated sample solution.

We use standardized staining (e.g. supravital staining) by Sternheimer for better recognition of elements. Staining reagent consists of 2 dyes (alcian blue and red pyronin B in 1:1 ratio). The staining reagent is added to the concentrated urine sample in 1:10 ratio.

In this database, we have sorted and described 3 variants of pictures.

1. Microscopic findings of stained sediment (10× concentrated urine sample with 400× magnification, Sternheimer staining)
2. Microscopic findings of native sediment (10× concentrated urine sample in 400× magnification)
3. Findings from automatic FUS-2000 analyzer (DIRUI)

In the following links the database is prepared in a format suitable for print. For better recognition, some elements are magnified.

Erythrocytes

Erythrocytes are red blood cells without nucleus, with size about 6–7 μm and a disk-like shape. They count amongst the smallest and the most common elements in urine. Their presence in urine (hematuria) can be macroscopic or microscopic (without visible red color).

Causes for haematuria:


- Renal (glomerulonephritis, kidney cancer)
- Prerenal (hemocoagulation aberrations, muscle traumas, burns)
- Subrenal (bleeding in urinary tract – infection, kidney stones, carcinoma)
- Exertion (physical stress, cold)

If the erythrocytes have a normal biconcave shape with smooth surface, they are called eumorphic erythrocytes. Erythrocytes that passed to urine through glomerular membrane might be damaged and their shape is changed – we call them dysmorphic erythrocytes.


Dysmorphic erythrocytes may have a tire shape (codocytes) or the erythrocyte membrane may have protrusions (acanthocytes).

Hedgehog shaped or crenated erythrocytes (echinocytes) don't count among the dysmorphic erythrocytes. They are deformed by erythrocyte dehydration in urine with high osmolality.


Stained sediment



Erythrocytes




Dysmorphic erythrocyte – codocyte




Dysmorphic erythrocytes – acanthocytes


Native sediment




Erythrocytes



Erythrocytes deformed by high osmolality (the cells lose water)



Dysmorphic cells – codocytes



Dysmorphic erythrocytes – acanthocytes

Pictures from FUS-2000 analyzer (DIRUI)

FUS-2000 hybridní močový analyzátor (CZ-MKO-150122-CSLV) -- Prac. list

Stav systému Prac. list Databáze QC Kalibrace Správa Nastavení Údržba Ukončení Odhlášení Nápověda (EN)

Ery. 2783/uL


Č. vzorku: 16

Leu. Bakterie
Ery.
Dl. epi. Krystaly ...
Shl. leu. Válce ...
Kulaté epi. Jiné ...
Artefakty Chemie ...


Zpět << >>

Standby Administrator moc 30°C 2016-03-31 18:07:03

Erythrocytes



Erythrocytes deformed by high osmolality



Dysmorphic erythrocytes – acanthocytes

Leukocytes

Neutrophil granulocytes are the most common leukocyte subcategory in urine. Their cells are round shaped with the size of 10–12 μm and with segmented nucleus in the center of the cell. The leukocyte nucleus may or may not be stained by the dye – vital leukocytes with undamaged membrane have colorless nucleus, cells with damaged membrane have their nucleus stained blue.

Other leukocyte types in urine such as **lymphocytes** (they have a large nucleus that fills almost the entire cell), **monocytes** (with nucleus in the shape of a horseshoe or a bean) and activated monocytes called **macrophages** can be rarely found. Diagnostic strip doesn't react with these leukocytes.

The granulocyte presence is typical especially for bacterial infections of urinary tract or kidneys. Semiquantitative detection with diagnostic strip is based on reaction with granulocyte esterase.

The presence of increased numbers of lymphocytes after renal transplantation is an important sign of kidney rejection.


Macrophages

Macrophages play important role in immune reaction. Their primary function is phagocytosis – absorption of elements inside their cells. They belong among mononuclear leukocytes, meaning they have just one non-segmented nucleus. They evolve from monocytes in tissues. The macrophages have increased amount of lysosomes and vacuolated cytoplasm. They may be seen with phagocytosed erythrocytes (erythrophages), lipid droplets (lipophages) or crystals.


Glitter cells

Some leukocytes are in hypotonic urine particularly distinguishable by the Brownian movement of granules inside their cells. This gave these neutrophil granulocytes the name "glitter cells". In some cases, especially in hypotonic urine, the leukocyte membrane may rupture and spill some of the cytoplasm outside of cell. We may observe this in samples of patients with interstitial nephritis.


Stained sediment




Leukocytes (granulocytes)




Leukocytes (granulocytes)




Granulocytes – glitter cells




Leukocyte (lymphocyte)




Monocytes (orange arrows) and activated monocytes (blue arrows)



Leukocytes (lymphocytes – arrows)




Leukocytes (monocytes), 1000× magnification




Leukocytes including macrophage with phagocytosed bacteria (arrow)


Native sediment



Leukocytes



Group of leukocytes




Granulocytes – glitter cells




Leukocytes including macrophages (arrows)


Pictures from FUS-2000 analyzer (DIRUI)




Leukocytes (granulocytes)



Leukocytes (including lymphocytes)



Leukocytes (including mononuclear cells)



Macrophage – activated monocyte


Epithelial cells

The outside or the inside of the organism surface is covered with epithelial cells. These cells can be found in urine and divided in several groups.


Squamous epithelial cells

These are very large unevenly shaped cells with easily visible nucleus inside. They originate from urethra or vagina. They are very common elements with minimal clinical impact.

Stained sediment




Squamous epithelial cells




Squamous epithelial cells

Native sediment




Squamous epithelial cells




Squamous epithelial cells

Pictures from FUS-2000 analyzer (DIRUI)



Squamous epithelial cells



Squamous epithelial cells


Transitional epithelial cells

These epithelial cells have different shapes and sizes depending on their origin. Round epithelial cells are most common. Their nucleus is in the middle of the cell, they are smaller than squamous epithelial cells and they come from the bladder or proximal segments of urethra in men. If they originate from deeper layers of epithelia or near the renal pelvis, they are smaller and more round.


Limited amount of round transitional epithelial cells may be normal, large amount accounts to urinary tract infection.

Another subcategory is called caudate cells. They come from deep layers of a bladder. The cells with two nuclei are also categorized as transitional epithelial cells. Large number of cells with two nuclei or asymmetric cells can be found in urine from patients with transitional cell (urothelial) carcinomas.


Stained sediment



Transitional epithelial cells




Transitional epithelial cells with two nuclei




Caudate transitional epithelial cells


Native sediment



Transitional epithelial cells




Transitional epithelial cells with two nuclei




Caudate transitional epithelial cell


Pictures from FUS-2000 analyzer (DIRUI)




Transitional epithelial cells



Transitional epithelial cells with cocci



Transitional epithelial cells with two nuclei




Caudate transitional epithelial cells

Renal epithelial cells


These cells have polyedric shape with non-segmented eccentric nucleus. They are the smallest epithelial cells, approximately twice the size of a leukocyte. They don't absorb water and swell, but keep their polyedric shape. Proximal tubular cells have granulated cytoplasm and sometimes may appear as small or fragmented granular casts. They are clinically significant elements in urine in cases of acute tubular necrosis or viral infection and drug or heavy metal toxicity.

Renal epithelial fragments constitute of several renal cells of collecting duct origin. Their presence in urine is considered to be clinically severe and indicates heavy renal tubular damage.


Stained sediment



Renal epithelial cells




Renal epithelial cells




Proximal renal
epithelial cell

Renal tubular epithelia




Proximal renal
epithelial cells

Renal tubular epithelia




Fragments of renal tubular epithelia


Native sediment



Renal epithelial cells




Renal tubular epithelia probably from proximal tubules




Fragments of renal tubular epithelia


Pictures from FUS-2000 analyzer (DIRUI)



Renal tubular epithelia



Renal tubular epithelia



Renal tubular epithelia and fragments of renal tubular epithelia

Casts


These elements are formed in the kidney tubules by Tamm-Horsfall mucoprotein precipitation. The mucoprotein is secreted from renal tubule cells. Their formation is reinforced by acidic pH in urine, higher concentration of plasmatic proteins, dehydration and excessive physical activity. Their shape copies the shape of a tubule with defined outer line, parallel sides and round ends.

We distinguish hyaline, cellular, granular, wax, lipid, bacterial and combined casts. The cast is classified as cellular or granular cast only if the amount of material inside takes up to 1/3 of its volume. Otherwise, it is called hyaline cast. The cast goes through different stages of development with increasing time in the kidney tubule: cell cast → granular cast → waxy cast.


Hyaline casts

Hyaline casts are formed only by Tamm-Horsfall glycoprotein without any other elements or their fragments inside. Due to their composition, they are almost undetectable in native sediment.

Stained sediment




Hyaline cast




Hyaline casts

Native sediment




Hyaline cast




Hyaline cast

Pictures from FUS-2000 analyzer (DIRUI)



Hyaline cast




Hyaline cast


Cell casts

Cellular casts comprise of cells that may occur in renal tubules (leukocytes, erythrocytes and tubular epithelia) entrapped in Tamm-Horsfall mucoprotein matrix. We recognize leukocyte casts, erythrocyte casts and renal tubular epithelial cell casts. In some cases, the cells in the cast cannot be accurately determined and the cast is therefore categorized as cellular.

Stained sediment




Erythrocyte cast




Hyaline cast with erythrocytes




Leukocyte cast



Hyaline cast with leukocytes inside



Renal epithelial cell cast




Renal epithelial cell casts

Native sediment




Erythrocyte cast



Leukocyte cast




Renal epithelial cell cast




Renal epithelial cell cast


Pictures from FUS-2000 analyzer (DIRUI)



Erythrocyte cast



Leukocyte casts and hyaline casts with leukocytes inside




Renal epithelial cell casts and hyaline casts with renal epithelia inside


Granular casts

Granules inside these casts are formed after decomposition of cells in the cast or a tubule. Granular casts may vary from coarsely granular including cell particles to finely granular that are turning to waxy casts. Small number of these casts may occur after intensive physical activities (patients who were exposed to cold conditions or cold hardening). Increased concentration of granular casts is strongly pathological.

Stained sediment




Granular cast



Granular cast

Native sediment




Granular cast




Granular cast

Pictures from FUS-2000 analyzer (DIRUI)



Granular casts




Granular casts


Waxy casts

Waxy casts are the clinically most serious type of casts and they are also called the casts of renal failure. They occur in patients with chronic kidney diseases. Their structure is homogenous, they have the biggest size and their endings are often broken. Sometimes they are partially made of granular matter. They indicate tubule damage.

Stained sediment



Waxy cast



Waxy cast

Native sediment




Waxy cast



Waxy cast

Pictures from FUS-2000 analyzer (DIRUI)




Waxy cast

Fatty casts

Fatty casts in protein matrix contain fat inclusions. They are related to strong insufficiency, nephritic syndrome, diabetics and mercury intoxication.

Stained sediment




Fatty cast


Bacterial casts

Bacterial casts are composed of bacteria in a protein matrix. They may be found in acute pyelonephritis or intrinsic renal infection. Bacterial casts should be seen in association with loose bacteria, leukocytes, and leukocyte casts. Their occurrence is extremely rare, due to their fragility and also commonly used antibiotic treatment.


Stained sediment



Bacterial cast




Bacterial cast




Bacterial cast

Native sediment



Bacterial cast

Pictures from FUS-2000 analyzer (DIRUI)




Bacterial cast


Pseudocasts

Pseudocasts are structures of aggregates that resemble and may be mistaken for casts because of their shape. They are without diagnostic significance. Such structures include mucus threads, leukocytes entrapped in mucus, rolled squamous epithelial cells, aggregates of amorphous urates, calcium oxalate crystals and so on. Artifacts don't belong in this category.


Stained sediment



Pseudocasts – mucus




Pseudocasts – group of leukocytes




Pseudocast – bilirubin crystals

Native sediment




Pseudocast – ammonium urate crystals (arrow)


Pictures from FUS-2000 analyzer (DIRUI)



Pseudocasts – microcrystals on mucus



Pseudocast – uric acid microcrystals



Pseudocasts – group of leukocytes


Microorganisms

Bacteria


Bacteria are single celled organisms. They have shapes of spheres (cocci) or rods and usually reach size of a few micrometers.

Small amount of bacteria in urine is quite common finding. Urine should be analyzed within one hour, otherwise the bacteria multiply. They tend to form chains or occur as longer fibers. Antibiotic treatment may cause bacteria cell wall disintegration producing unusually long rod-shaped elements called bacteria protoplasts. Larger quantities of bacteria can often be found in pathological urine with leukocytes.

Stained sediment




Rod-shaped bacteria that remained unstained after dyeing




Cocci (detail) and protoplast bacteria (arrow) – stained after dyeing


Native sediment



Chained bacteria – cocci




Bacteria – cocci




Bacteria – rods


Pictures from FUS-2000 analyzer (DIRUI)



Bacteria – cocci



Bacteria – rods




Fibers of bacteria


Yeast

Yeasts are quite common pathological finding. They are unicellular microorganisms with typical round or oval shape. They multiply by budding and in some cases they form pseudomycelium (fibrous form of yeast). They may occur in urine of immunodeficient or immunosuppressed patients and may also be present in urine of diabetics, because glucose contributes to their growth. The most common species in urine is *Candida albicans*.

Stained sediment




Yeast



Yeast pseudomycelium

Native sediment




Yeast




Yeast pseudomycelium

Pictures from FUS-2000 analyzer (DIRUI)



Yeast and yeast pseudomycelium



Yeast pseudomycelium

Crystals


The presence of crystals and amorphous microcrystalline deposits in urine is not considered a significant clinical finding. However, both parameters are determined and their amount is evaluated.

Crystals have various crystalline structure and they occur in many forms. The pH of urine is important factor for their formation and structure, although it is sometimes difficult to distinguish them even then. In that case, the elements could be classified only as crystals - without further specification.


The most frequent crystals in urine are: oxalate and uric acid (in acidic urine) or phosphate (in alkaline urine). Rarely found crystals in urine: bilirubin, cysteine, leucine, tyrosine or drug. We distinguish two types of amorphous microcrystals – amorphous urates in acidic urine and amorphous phosphates in alkaline urine.

Oxalates

Stained sediment




Calcium oxalate dihydrate (envelope-shaped)




Calcium oxalate monohydrate (oval and biscuit-shaped)

Native sediment




Calcium oxalate – monohydrate (oval form) and dihydrate




Calcium oxalate – monohydrate (oval form and biscuit-shaped) and dihydrate


Pictures from FUS-2000 analyzer (DIRUI)



Calcium oxalate dihydrate (envelope-shaped)



Calcium oxalate monohydrate




Calcium oxalate – monohydrate (oval form) and dihydrate (evelope-shaped)

Uric acid

Various forms of crystals of uric acid


Stained sediment




Uric acid (lemon-shaped)



Uric acid (barrel-shaped)




Uric acid (needle form)



Uric acid

Native sediment




Uric acid (lemon-shaped)



Uric acid (barrel-shaped)




Uric acid (needle form)




Uric acid


Pictures from FUS-2000 analyzer (DIRUI)




Uric acid (lemon-shaped)



Massive uric acid




Uric acid




Uric acid

Ammonium urate

Stained sediment




Ammonium urate




Ammonium urate

Native sediment



Ammonium urate

Pictures from FUS-2000 analyzer (DIRUI)




Ammonium urate


Triple phosphate

Ammonium magnesium phosphate crystals (Triple phosphate).


Stained sediment




Triple phosphate (coffin-shaped)



Triple phosphate



Triple phosphate crystal




Triple phosphate (arrows)


Native sediment




Triple phosphate (coffin-shaped)



Triple phosphate




Triple phosphate




Triple phosphate (arrows)


Pictures from FUS-2000 analyzer (DIRUI)



Triple phosphate



Triple phosphate (coffin-shaped)




Triple phosphate (coffin-shaped)


Calcium phosphate

These crystals can be differentiated from uric acid crystals by polarization microscopy. Unlike uric acid, calcium phosphate doesn't turn polarized light.


Stained sediment



Calcium phosphate



Calcium phosphate (needle-shaped and star-shaped druses)




Calcium phosphate

Native sediment



Calcium phosphate




Calcium phosphate (needle-shaped and star-shaped druses)




Calcium phosphate

Pictures from FUS-2000 analyzer (DIRUI)




Calcium phosphate (needle-shaped)




Calcium phosphate (star-shaped druses)

Bilirubin

Stained sediment




Bilirubin




Bilirubin

Native sediment




Bilirubin



Bilirubin


Pictures from FUS-2000 analyzer (DIRUI)




Bilirubin

Cystine

Stained sediment




Cystine




Cystine

Native sediment




Cystine




Cystine

Pictures from FUS-2000 analyzer (DIRUI)




Cystine




Cystine

Amorphous microcrystals

Stained sediment




Amorphous microcrystals




Amorphous microcrystals

Native sediment




Amorphous microcrystals




Amorphous microcrystals

Pictures from FUS-2000 analyzer (DIRUI)



Amorphous microcrystals




Amorphous microcrystals

Lipid particles

Oval fat bodies are renal tubular epithelial cells or macrophages filled with fat globules. The nuclei of these cells are barely visible. Fat bodies are accompanied by the presence of free fat droplets and possibly by hyaline casts with fat inclusions or fatty casts. Oval fat bodies and free fat droplets (globules) in urine are considered to be an exceptional and serious finding. The presence of fat with biological origin in urine is called lipiduria and indicates severe renal dysfunction. It is associated with serious damage and necrosis of renal epithelial cells as in nephrotic syndrome, advanced diabetes mellitus or some kinds of poisoning.


Fat globules may be seen in the urine as a contamination from oiled catheter or vaginal creams. There are no fat bodies and casts in sample in this case. Examples are shown in the Artifacts category.

Stained sediment




Macrophage and fat droplets

Native sediment




Macrophage with fat droplets and sperm

Pictures from FUS-2000 analyzer (DIRUI)



Fat droplets




Squamous epithelial cells with fat droplets

Mucus


Mucus is secreted by glands in urinary tract and vagina. Its quantity could be increased with inflammatory conditions. It is common constituent in urine with no diagnostic significance.

Stained sediment




Mucus

Native sediment



Mucus

Pictures from FUS-2000 analyzer (DIRUI)




Mucus


Sperms

The finding of sperms is common in man's urine. It is insignificant in women's samples as long as they are not underage girls.

Stained sediment



Sperms




Sperms

Native sediment




Sperms




Sperms

Pictures from FUS-2000 analyzer (DIRUI)



Sperms




Sperms


Artifacts

We can find some elements in urine that don't come from patient's body. They have no clinical significance, but it is important to recognize them to avoid confusion with another element. They can have various shapes. Textile and paper fibers and oil droplets are the most common artifacts.


Stained sediment



Artifact




Artifact




Fat droplets (artificial contamination)

Native sediment




Artifact




Fat droplets (artificial contamination)

Pictures from FUS-2000 analyzer (DIRUI)



Artifact




Fat droplets (artificial contamination)


Interesting findings

Erythrocytes x yeast

Occasionally, it is difficult to distinguish between single yeast and erythrocyte due to their similar shape and size in the pictures from automatic analyzer and in native sediment. The stained sediment can be used for determinative analysis, because erythrocytes are typically stained pink, but yeasts are colorless.




Yeast between erythrocytes




Erythrocytes and yeast together

Erythrocytes x oxalates

If erythrocytes and oxalates are both in urine, it is sometimes difficult to distinguish them in the automatic analyzer properly due to their similar size and shape. This applies particularly to calcium oxalate monohydrate (ovoid form). In the stained sediment on the other hand, the erythrocytes are pink and oxalates colorless.




Calcium oxalate monohydrate crystals next to erythrocytes automatically classified as erythrocytes in FUS-2000




Calcium oxalate monohydrate crystals next to erythrocytes

The gradual transformation of pathological casts


Exceptionally, all stages of cast transformations could be observed in one sample.




Cellular cast




Transition phase of cast from cellular to granular (enlargement ×600)




Transition phase of cast from cellular to granular



Granular cast




Transition phase of cast from granular to waxy and waxy cast




Waxy cast

The finding of fat particles in patient with nephrotic syndrome


The elements listed below were found in urine of patient with nephrotic syndrome.



Hyaline cast with fat inclusions of fat droplets



Squamous epithelial cell and oval fat body




Fatty cast and free fat droplets

Elements in hypotonic urine


A patient with acute urine bladder inflammation was treated by hydration of the bladder, causing his urine to be hypotonic.

In the urine sample, we found neutrophil granulocytes which are called “glitter cells” because of the rapid Brownian movement of granules inside their cells. The low urine osmolality caused ruptures in the cell membranes of several of these leukocytes which resulted in cytoplasm spilling outside of the cells. These cells are sometimes called “winged leukocytes”.


The hypotonic urine from the same patient also caused erythrocytes to swell and their size matched that of the leukocytes. The same size resulted in miscategorization of some erythrocytes in the leukocyte category by the FUS-2000 analyzer software leading to a discrepant finding.




Leukocytes – some of them glittering cells (pointer)



Swollen erythrocytes in hypotonic urine




Leukocytes (glittering cells) with erythrocytes




Leukocytes (glittering cells) with erythrocytes

Sample contaminated by feces


The urine analyzed with FUS-2000 showed some dark asymmetric particles miscategorized as squamous epithelial cells. Further investigation with microscopy in stained and native sediment confirmed them to be stool particles. This rare contamination of feces may occur in patients with fistula of bladder.



Fecal particles (black elements) miscategorized in squamous epithelia section




Fecal particles (brown elements) with squamous cells




Fecal particles

Intracellular Bacteria


Bacteria, most commonly *E. coli*, may be sometimes observed inside epithelial cells of the bladder in samples of patients with urinary tract infection. Insufficient antibiotic treatment does not eradicate bacteria inside the cells and may cause future chronic infections.



Bacteria inside squamous epithelia




Bacteria inside squamous and transitional epithelia




Bacteria inside squamous and transitional epithelia (cropped from the picture above)

Atypical cells

Occasionally, atypical cells may be found with urine sediment microscopy. In this case, the cells are probably of epithelial origin with variable shape, size and multiple nuclei. Such finding should be reported to the responsible physician. The patient was diagnosed with bladder cancer.



Atypical cells



Atypical cells

Table of contents

Introduction.....	2
Erythrocytes	3
Leukocytes.....	9
Epithelial cells.....	18
Squamous epithelial cells.....	18
Transitional epithelial cells.....	22
Renal epithelial cells.....	27
Casts	34
Hyaline casts.....	34
Cell casts.....	38
Granular casts.....	45
Waxy casts.....	49
Fatty casts.....	52
Bacterial casts.....	53
Pseudocasts.....	56
Microorganisms.....	60
Bacteria.....	60
Yeast	65
Crystals	69
Oxalates.....	70
Uric acid.....	74
Ammonium urate	80
Triple phosphate	82
Calcium phosphate.....	88
Bilirubin	93
Cystine.....	96
Amorphous microcrystals	99
Lipid particles	102
Mucus.....	105
Sperms.....	107

Artifacts	111
Interesting findings.....	115
Erythrocytes x yeast	115
Erythrocytes x oxalates	117
The gradual transformation of pathological casts	119
The finding of fat particles in patient with nephrotic syndrome	122
Elements in hypotonic urine	124
Sample contaminated by feces	127
Intracellular Bacteria	129
Atypical cells.....	131
Table of contents.....	133