

IA038 Types and Proofs

4. The Curry-Howard Isomorphism

Jiří Zlatuška

March 14, 2023

3.1.2 Terms

$[x_i : T_i]^R$ Hypothesis [and its parcel]

$$\frac{[x : U]^R}{\frac{\vdots}{v : V} \rightarrow \text{-I}^R}
 \quad
 \frac{t : U \rightarrow V \quad u : U}{tu : V} \rightarrow \text{-E}$$

$\lambda x. v : U \rightarrow V$

$$\frac{\neg u : U \quad v : V}{\langle u, v \rangle : U \times V} \times \text{-I}
 \quad
 \frac{t : U \times V}{\pi^1 t : U} \times \text{-1E}
 \quad
 \frac{t : U \times V}{\pi^2 t : V} \times \text{-2E}$$

3.1.2. Terms using another formulation

$$\begin{array}{c}
 \frac{x_i : T_i \vdash x_i : T_i}{\text{Id}_p} \quad \text{Axiom [parcel } p\text{]} \\
 \frac{\Gamma, x : U \vdash v : V}{\Gamma \vdash \lambda x. v : U \rightarrow V} \rightarrow\text{-I} \quad \frac{\Gamma \vdash t : U \rightarrow V \quad \Delta \vdash u : U}{\Gamma, \Delta \vdash tu : V} \rightarrow\text{-E} \\
 \frac{\Gamma \vdash u : U \quad \Delta \vdash v : V}{\Gamma, \Delta \vdash \langle u, v \rangle : U \times V} \times\text{-I} \quad \frac{\Gamma \vdash t : U \times V}{\Gamma \vdash \pi^1 t : U} \times\text{-1E} \quad \frac{\Gamma \vdash t : U \times V}{\Gamma \vdash \pi^2 t : V} \times\text{-2E}
 \end{array}$$

or expressed in logic notation

$$\begin{array}{c}
 \frac{}{x_i : T_i \vdash x_i : T_i} \text{Id}_p \quad \text{Axiom [parcel } p\text{]} \\
 \frac{\Gamma, x : U \vdash v : V}{\Gamma \vdash \lambda x. v : U \rightarrow V} \rightarrow\text{-I} \quad \frac{\Gamma \vdash t : U \rightarrow V \quad \Delta \vdash u : U}{\Gamma, \Delta \vdash tu : V} \rightarrow\text{-E} \\
 \frac{\Gamma \vdash u : U \quad \Delta \vdash v : V}{\Gamma, \Delta \vdash \langle u, v \rangle : U \wedge V} \wedge\text{-I} \quad \frac{\Gamma \vdash t : U \wedge V}{\Gamma \vdash \pi^1 t : U} \wedge\text{-1E} \quad \frac{\Gamma \vdash t : U \wedge V}{\Gamma \vdash \pi^2 t : V} \wedge\text{-2E}
 \end{array}$$

3.1.4 Conversion

Expressed using Natural Deduction derivation trees:

$$\frac{\begin{array}{c} [x : U]^x \\ \vdots \\ v : V \\ \hline \lambda x.v : U \rightarrow V \end{array}}{\frac{u : U}{(\lambda x.v)u : V}} \rightarrow\text{-I}^x \quad \rightarrow\text{-E}$$

$$v[u/x] : V$$

$$\frac{\begin{array}{c} U \\ \vdots \\ V \\ \hline U \Rightarrow V \end{array}}{V} \Rightarrow\text{-I} \quad \Rightarrow\text{-E} \Rightarrow V$$

$$\frac{\begin{array}{c} U \\ \vdots \\ V \\ \hline U \Rightarrow V \end{array}}{V} \Rightarrow\text{-I} \quad \Rightarrow\text{-E} \Rightarrow V$$

$$\frac{\begin{array}{c} u : U \quad v : V \\ \hline \langle u, v \rangle : U \times V \end{array}}{\pi^1 \langle u, v \rangle : U} \times\text{-I} \quad \times\text{-E} \Rightarrow u : U$$

$$\frac{\begin{array}{c} U \quad V \\ \hline U \wedge V \end{array}}{U \wedge V} \wedge\text{-I} \quad \wedge\text{-1E} \Rightarrow U$$

$$\frac{\begin{array}{c} u : U \quad v : V \\ \hline \langle u, v \rangle : U \times V \end{array}}{\pi^2 \langle u, v \rangle : U} \times\text{-I} \quad \times\text{-2E} \Rightarrow v : V$$

$$\frac{\begin{array}{c} U \quad V \\ \hline U \wedge V \end{array}}{U \wedge V} \wedge\text{-I} \quad \wedge\text{-2E} \Rightarrow V$$

Conversion expressed using alternative logical system for ND

$$\begin{array}{c}
 \text{A} \quad \frac{\Gamma, x : U \vdash v : V}{\Gamma \vdash \lambda x.v : U \rightarrow V} \rightarrow\text{-I} \quad \Delta \vdash u : U \quad \rightarrow\text{-E} \quad \Rightarrow \quad \boxed{\Gamma, \Delta \vdash t[u/x] : V} \\
 \Gamma, \Delta \vdash (\lambda x.t)u : V \\
 \frac{\Gamma \vdash u : U \quad \Delta \vdash v : V}{\Gamma, \Delta \vdash \langle u, v \rangle : U \times V} \times\text{-I} \quad \times\text{-1E} \quad \Rightarrow \quad \boxed{\Gamma \vdash u : U} \\
 \frac{\Gamma \vdash u : U \quad \Delta \vdash v : V}{\Gamma, \Delta \vdash \langle u, v \rangle : U \times V} \times\text{-I} \quad \times\text{-2E} \quad \Rightarrow \quad \boxed{\Delta \vdash v : V}
 \end{array}$$

The diagram illustrates the conversion of terms between different logical systems. It shows three main components: 1) A conversion rule for function application (\rightarrow -I and \rightarrow -E), 2) A conversion rule for pair formation (\times -I and \times -1E), and 3) A conversion rule for projection (\times -I and \times -2E). Red annotations and boxes highlight specific parts of the derivation, such as the resulting terms and the intermediate steps.

A few bits of history dates

1934 Gentzen's Natural Deduction

1940 Church's Lambda-Calculus

1956 Prawitz published normalization of Natural deduction proofs directly (Gentzen used Sequent Calculus, even though a direct proof in ND by him was recently discovered in his writings¹)


1956 Curry and Feys published Combinatory Logic, a system based purely on combinators

without variables; the types of basic combinators (I, K, S) corresponded to Hilbert axioms for Propositional Logic


1969 Howard combined results of Prawitz, Curry and Feys into the correspondence/isomorphism

1980 Howard's work published in "To H. B. Curry", a festschrift for Curry's 80 birthday

¹ Plato and Gentzen: Gentzen's Proof of Normalization for Natural Deduction, The Bulletin of Symbolic Logic, Vol. 14, No. 2, June 2008, 240-257



Some essential pieces of the correspondence



Aside: Combinatory Logic and Hilbert-style systems

Combinatorial terms over alphabet consisting of constants $\mathbf{K}_{A \Rightarrow (B \Rightarrow A)}$, $\mathbf{S}_{(A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C))}$, for every $A, B, C \in \text{Typ}$, and typed variables $x_T \in \mathcal{C}^T$:

$$\mathbf{K}_{A \Rightarrow (B \Rightarrow A)} \in \mathcal{C}^{A \Rightarrow (B \Rightarrow A)}$$

$$\mathbf{S}_{(A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C))} \in \mathcal{C}^{(A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C))}$$

$$x \in \mathcal{C}^T \quad x \text{ var of type } T$$

Hilbert system

Two axiom schemes

$$A \Rightarrow (B \Rightarrow A)$$

$$((A \Rightarrow (B \Rightarrow C)) \Rightarrow ((A \Rightarrow B) \Rightarrow (A \Rightarrow C)))$$

and Modus Ponens:

$$\frac{A \quad A \Rightarrow B}{B}$$

Example

Proof of $A \Rightarrow A$ in Hilbert system:

$$(\mathbf{S}_{(A \Rightarrow (B \Rightarrow A) \Rightarrow A) \Rightarrow ((A \Rightarrow (B \Rightarrow A)) \Rightarrow (A \Rightarrow A))} \mathbf{K}_{A \Rightarrow ((B \Rightarrow A) \Rightarrow A)} \mathbf{K}_{A \Rightarrow (B \Rightarrow A)} \in \mathcal{C}^{A \Rightarrow A}$$

Hilbert system proof:

$$\frac{\frac{(A \Rightarrow ((B \Rightarrow A) \Rightarrow A)) \Rightarrow ((A \Rightarrow (B \Rightarrow A)) \Rightarrow (A \Rightarrow A)) \quad A \Rightarrow ((B \Rightarrow A) \Rightarrow A)}{(A \Rightarrow (B \Rightarrow A) \Rightarrow (A \Rightarrow A))} \quad A \Rightarrow (B \Rightarrow A)}{A \Rightarrow A}$$

This corresponds to a term $(\mathbf{SK})\mathbf{K} : A \Rightarrow A$:


$$\frac{\frac{\mathbf{S}_{(A \Rightarrow ((B \Rightarrow A) \Rightarrow A)) \Rightarrow ((A \Rightarrow (B \Rightarrow A)) \Rightarrow (A \Rightarrow A))} \mathbf{K}_{A \Rightarrow ((B \Rightarrow A) \Rightarrow A)} \quad \mathbf{K}_{A \Rightarrow (B \Rightarrow A)}}{\mathbf{SK} \in \mathcal{C}^{(A \Rightarrow (B \Rightarrow A)) \Rightarrow (A \Rightarrow A)}} \quad \mathbf{K}_{A \Rightarrow (B \Rightarrow A)}}{(\mathbf{SK})\mathbf{K} \in \mathcal{C}^{A \Rightarrow A}}$$

Conversion between proofs in Hilbert system and Natural Deduction follows from

1. $\lambda x.x = (\mathbf{SK})\mathbf{K}$
2. $\lambda x.M = \mathbf{K}M$, for $x \notin \text{FV}(M)$,
3. $\lambda x.MN = S(\lambda x.M)(\lambda x.N)$.

$$\mathbf{KML} \rightarrow M$$

Example of term normalization



Natural Deduction definition of conversion/proof simplification

$$\frac{\frac{[x : U]^x}{\vdots} \quad v : V}{\lambda x.v : U \Rightarrow V} \Rightarrow I^x \quad u : U \quad \frac{}{\Rightarrow E} \quad \Rightarrow \quad v[u/x] : V$$

$$\frac{u : U \quad v : V}{\langle u, v \rangle : U \wedge V} \wedge I \quad \frac{}{\pi^1 \langle u, v \rangle : U} \wedge 1 E \quad \Rightarrow \quad u : U$$

$$\frac{u : U \quad v : V}{\langle u, v \rangle : U \wedge V} \wedge I \quad \frac{}{\pi^2 \langle u, v \rangle : U} \wedge 2 E \quad \Rightarrow \quad v : V$$

$$\begin{array}{c}
\frac{[z : B \times A]^z}{\pi^2 z : A} \times\text{-2E} \quad \frac{[z : B \times A]^z}{\pi^1 z : A} \times\text{-1E} \\
\hline
\frac{\langle \pi^2 z, \pi^1 z \rangle : A \times B}{\lambda z. \langle \pi^2 z, \pi^1 z \rangle : (B \times A) \rightarrow (A \times B)} \times\text{-I} \\
\hline
(\lambda z. \langle \pi^2 z, \pi^1 z \rangle)(\langle y, x \rangle) : A \times B \quad \rightarrow\text{-E} \\
\hline
\beta\text{-conversion} \\
\hline
\frac{\frac{[y : B]^y [x : A]^x}{\langle y, x \rangle : B \times A} \times\text{-I}}{\pi^2 \langle y, x \rangle : A} \times\text{-2E} \quad \frac{\frac{[y : B]^y [x : A]^x}{\langle y, x \rangle : B \times A} \times\text{-I}}{\pi^1 \langle y, x \rangle : A} \times\text{-1E} \\
\hline
\frac{\pi^2 \langle y, x \rangle, \pi^1 \langle y, x \rangle : A \times B}{\langle \pi^2 \langle y, x \rangle, \pi^1 \langle y, x \rangle \rangle : A \times B} \times\text{-I} \\
\hline
\downarrow\downarrow \text{pairing} \\
\frac{[x : A]^x [y : B]^y}{\langle x, y \rangle : A \times B} \times\text{-I}
\end{array}$$

Using logic notation

$$\frac{\Gamma, x : U \vdash v : V}{\Gamma \vdash \lambda x.v : U \Rightarrow V} \Rightarrow\text{-I} \quad \Delta \vdash u : U \quad \Rightarrow\text{-E} \implies \Gamma, \Delta \vdash (\lambda x.t)u : V$$

$$\frac{\Gamma \vdash u : U \quad \Delta \vdash v : V}{\Gamma, \Delta \vdash \langle u, v \rangle : U \wedge V} \wedge\text{-I} \quad \wedge\text{-1E} \implies \Gamma \vdash u : U$$

$$\frac{\Gamma \vdash u : U \quad \Delta \vdash v : V}{\Gamma, \Delta \vdash \langle u, v \rangle : U \wedge V} \wedge\text{-I} \quad \wedge\text{-2E} \implies \Delta \vdash v : V$$

Proof simplification using the logic-based system

$$\begin{array}{c}
 \frac{}{z : B \wedge A \vdash z : B \wedge A} \text{Id}_z \quad \frac{}{z : B \wedge A \vdash z : B \wedge A} \text{Id}_z \\
 \frac{}{z : B \wedge A \vdash \pi^2 z : A} \wedge\text{-2E} \quad \frac{}{z : B \wedge A \vdash \pi^1 z : A} \wedge\text{-1E} \\
 \frac{}{z : B \wedge A \vdash \langle \pi^2 z, \pi^1 z \rangle : A \wedge B} \wedge\text{-I} \\
 \frac{}{\vdash \lambda z. \langle \pi^2 z, \pi^1 z \rangle : (B \wedge A) \Rightarrow (A \wedge B)} \Rightarrow\text{-I}_z \\
 \frac{}{x : A, y : B \vdash (\lambda z. \langle \pi^2 z, \pi^1 z \rangle)(\langle y, x \rangle) : A \wedge B} \Rightarrow\text{-E}
 \end{array}$$

\Downarrow β -conversion

$$\begin{array}{c}
 \frac{}{y : B \vdash y : B} \text{Id}_y \quad \frac{}{x : A \vdash x : A} \text{Id}_x \\
 \frac{}{x : A, y : B \vdash \langle y, x \rangle : B \wedge A} \wedge\text{-I} \quad \frac{}{y : B \vdash y : B} \text{Id}_y \quad \frac{}{x : A \vdash x : A} \text{Id}_x \\
 \frac{}{x : A, y : B \vdash \pi^2 \langle y, x \rangle : A} \wedge\text{-2E} \quad \frac{}{x : A, y : B \vdash \langle y, x \rangle : B \wedge A} \wedge\text{-1E} \\
 \frac{}{x : A, y : B \vdash \pi^1 \langle y, x \rangle : A} \wedge\text{-I} \\
 x : A, y : B \vdash \langle \pi^2 \langle y, x \rangle, \pi^1 \langle y, x \rangle \rangle : A \wedge B
 \end{array}$$

$\Downarrow\Downarrow$ pairing

$$\frac{}{x : A \vdash x : A} \text{Id}_x \quad \frac{}{y : B \vdash y : B} \text{Id}_y \\
 \frac{}{x : A, y : B \vdash \langle x, y \rangle : A \wedge B} \wedge\text{-I}$$

