

MONITOROVACÍ A DROGOVÉ ZÁVISLOSTI



Methamphetamine use in the Czech Republic



History of methamphetamine

- 1950s Misuse of psychostimulant medicines (Psychoton)
- 1960s Misuse phenmetrazine
- Since 1970s methamphetamine (pervitin)
 (manufactured home-made in a "kitchen" labs)
- Pervitin was mostly made either of ephedrine coming from ephedrine company in Roztoky nearby Prague or of Solutan containing pseudoephedrine
- Small group of users/producers with division of roles in the production (glass, precursors, chemicals...)
- Since 1990s the main problem drug in the Czech Cepublic



Prevalence of methamphetamine use in the general population

- General population survey (2008)
 - Life time prevalence: 5% in the age group 25-34: 9 %
 - Last year prevalence: 2%
 - Last month prevalence: 1%
- ESPAD 2007
 - 3,5% a decline since 1999
- Slight increase in the use of pervitin within nightlife settings
 - Last month prevalence in 2003 14%, in 2007 16%




Problem drug use of methamphetamine

- In the long term: total estimates of 30 000 PDU
 - 95% injecting users
 - 2/3 20 000 PDU preferring pervitin
 - Significant increase in PDU estimates in 2009 to 37 400 (from 32 500 in 2008)
- In 2008 21 000 problem pervitin users
- In 2009 25 000 problem pervitin users
 - increase by 4 000 in comparison with 2008
- Usti nad Labem region (4 000), Prague (3 000)



Problem Drug Use (PDU)

Number of problem drug users per 1,000 inhabitants aged 15-64






Treatment

- Pervitin users accounted for 60% of all the treatment demands
- **2009**:
 - 5 209 in total, 1/3 women
 - the average age: 25 years, 20% people under 19 years
 - secondary drugs: 55% cannabis, 15% opiates.
- Psychiatric co morbidity:
 - acute toxic psychosis, anxious and depressive disorders
 - higher level of irritability (attacks of anger), paranoid syndromes, unexpected mood/emotional changes, pathological jealousy, memory and concentration disorders, social phobia etc.



Treatment

Number of first treatment demands according to drug used, 1997-2009





Pervitin related infections and deaths

- Fatal overdoses 18 cases (19 in 2008) (direct drug related deaths) in terms of pervitin
- HIV prevalence in IDUs very low: app. 1 pro mile
 - 7 new HIV positive IDUs
 - At the end of 2009: 1 344 HIV positive cases, 65 out of them IUD (app. 5 %)
- HCV prevalence in IDUs: 20 40%
 - univariate analysis: prevalence of HCV lower in pervitin than in opiate users
 - multivariate analysis: after adjustment for duration of injecting and needle sharing no statistical difference




Services for pervitin users

- No specific interventions for pervitin users are generaly implemented in practise nowadays
- Harm reduction services
 - low thresholds programmes (about 95 facilities)
 - Distribution of gelatine capsules per os alternative for pervitin use
 (drug user put pervitin into capsule and use it by a mouth instead of injecting)
- Treatment modalities
 - inpatient and outpatient facilities (various forms)
 - Psycho-pharmacotherapy for treatment of psychiatric comorbidity
 - **Sporadic methamphetamine substitution** by methylphenidate (Ritalin) and/or by vigabatrin (Sabril) lack of evaluation, just case reports
 - Psychotherapy is based especially on cognitive-behavioral therapy
 - Sociotherapy, after treatment care, supportive accommodation and work
- CA tests incentive therapy based on behavioral therapy
- P-centrum tests psychotherapy program based on 12 steps (AA)



Gelatine capsules for oral use of pervitin






Primary drug related criminality (generally)

- drug use is not illegal/punished
- possession of drugs for personal use: up to 2g
- 2 500 persons prosecuted
- 90% of them is charged
- 1 400 is sentenced (in total for all types of drugs)
- Proportion of drug crimes related to production and trafficking: 80%
- Proportion of drug crimes related to pervitin: 50%







Secondary criminality

- committed by drug users to obtain money for buying drugs
- 20% of total criminality related to drug users, mostly stealing things from cars

Criminality committed under influence of drugs

- 2% of all crime (7000 in total) committed under non alcoholic drugs, mostly influence of pervitin
- For comparison: 18% is committed under influence of alcohol



Production and market with pervitin nowadays

- Estimated consumption in 2008
 - 4,5 tons with 80% purity, 0,1 tons for export mainly to Germany
- Until 2004 pervitin made of ephedrine
 - afterwards stopped production of ephedrine in Roztoky
 - ephedrine replaced by pseudoephedrine only (e.g. Modafen, Neurofen Stop Grip)
- In 2009
 - restriction for sales of medicaments containing pseudoephedrine in pharmacies
 - due to this measure higher level of import of medicaments containing pseudoephedrine especially from Poland
 - estimation: 95% of all medicines intended for pervitin production at present
- Increased illegal import of pharmaceuticals containing pseudoephedrine from abroad (Poland)








Methamphetamine production

Seizures of laboratories, precursors and methamphetamine

Sales of pharmaceuticals containing pseudoephedrine in the Czech pharmacies – decrease in 2009



Seizures	2007	2008	2009
Ephedrine (g)	1,185	1,677	6,023
Pseudoephedrine (g)	218	_	_
Modafen® (tablets)	3,480	7,876	840
Nurofen® StopGrip (tablets)	11,948	21,785	876
Panadol® Plus Grip (tablets)	72	17,021	1,224
Paralen® Plus	_	_	1,440
Acatar® (tablets)	_	_	3,508
Cirrus® (tablets)	_	_	6
Ibuprofen® (tablets)	_	_	80
Ibuprom® (tablets)	_	_	22,080
Sudafed® (tablets)	_	_	12,231
Cooking labs	388	434	342
Pervitin (g)	5,978	3,799	3,599



Price and purity of methamphetamine

- Only/mostly retail market level
- Average price stable:
 - 1 000,- CZ crowns (about 40 €) per 1g
- Average purity:
 - 70% (in the long term) pervitin is often cut by piracetam
- Sporadically wholesale market level
 - 500 000,- CZ crowns (about 20 000 €) per 1kg



Medicines containing pseudoephedrine in illicit lab





Summary and prospects

■ In 2008 the Government Council for Drug Policy Coordination established the **methamphetamine working group**, an inter agency, multidisciplinary body with representatives from important state authorities, institutions, academic bodies, NGO s etc.

Discussed topics:

- **Treatment** (new specific interventions: e.g. contingency management therapy, substitution)
- **Harm reduction** (eg. gelatine capsules)
- Supply reduction (measures to decrease of availability of precursors and other chemicals (e.g. red phosphorus)



Conclusion

- Methamphetamine (pervitin) is historically the main problem drug in the Czech Republic
- Produced and consumed (almost exclusively) domestically
- Made of ephedrine in past and pseudoephedrine from medicines nowadays
- Though sale of medicines containing pseudoephedrine has been limited and decreased since 2009, it has had practically NO IMPACT on supply, price, level of use of pervitin
- Import of medicines containing pseudoephedrine from abroad as source for pervitin production
- Measures in neighbouring countries? At EU level?
- Risk of increased use of other stimulants? Cocaine? New synthetic cathinones and phenetylamines?



Thanks ©

mravcik.viktor@vlada.cz www.drogy-info.cz www.focalpoint.cz www.emcdda.europa.eu