### Climate change and fossil fuels

Filip Černoch <u>cernoch@mail.muni.cz</u>



# Explaining the climate change

- "How could scientists predict the climate in 100 years when they cannot predict the weather tomorrow?"
- Climate: atmospheric conditions over a long period of time (years to centuries).
- Weather: short time (minutes to weeks).
- Consequences for prediction climate undergoes more gradual change (than weather) and is easier to predics.




### 1) The planet's temperature is rising

- •Over the past 130 years the global average temperature has increased by 0,8°C (more than half of that in last 35 years).
- •Ocean accounts for more than 90% of the energy accumulated between 1971-2010.
- •Ancient ice samples (from Antarctica and other places) their layers are dated and gas bubles inside are analysed.
  - CO2 concentration is measured by infrared spectroscopy or mass spektromectry.
  - Isotope ratios of water molecules are measured to determine historical temperatures.


### The planet's temperature is rising

- Earth's climate has always fluctuated. The cooler period ice ages or glacial periods, the warmer period
  - interglacial periods.
    - Orbital variations (Milankovitch cycles)
    - Solar output
    - Volcanism
    - Plate tectonics
- The rate of change has become more dramatic since the Industrial Revolution = anthropogenic origins.
  - Problems with adaptation the change is too fast.












# Observed change in surface temperature 1901 - 2012






# 2) CO<sub>2</sub> level is increasing (also methane and nitrous oxide)

- $CO_2$  concentration increased by 40% since pre-industrial time. The ocean absorbed about 30% of this increase, causing ocean acidification.
- Also methane (150% increase) and nitrous oxide (20%).








# 3) We are responsible for the increase in $CO_2$

- Human CO<sub>2</sub> emissions (20 billion tons/y) are small compared to natural emission (776 billion tons/y).
- But natural absorptions (788 billion tons/y) roughly balance natural emisssions.
- Carbon 12 isotope to carbon 13 isotope ratio increases (isotope = different atoms with the same chemical behavior but with different masses).





Global carbon cycle. Numbers represent reservoir mass (carbon stocks) and annual carbon exchange fluxes. Black numbers and arrows indicate reservoir mass and exchange fluxes estimated for the time prior to the Industrial Era (about 1750). Red indicate annual **CENTER FOR** anthropogenic fluxes averaged over the 2000-2009.

# 4) Increased CO<sub>2</sub> is the primary driver of greenhouse effect

- Inbound solar radiation has short wavelenghts and high energy contents. This radiation passes through the atmosphere. Some energy is absorbed by the ground (warming it up). Some energy is reflected back to the space.
- That reflected radiation has lower energy levels and longer wavelengths. 80% of the outgoing radiation is trapped in the lower troposphere.
- Energy trapped in the troposphere wams the surface.
- More GHGs in the atmosphere trap more outbound solar radiation, thus warming the planet anthropogenic climate change.



Sunlight passes through the atmosphere and warms the Earth's surface. This heat is radiated back toward space.


> Most of the outgoing heat is absorbed by greenhouse gas molecules and re-emitted in all directions, warming the surface of the Earth and the lower atmosphere.



# 4) Increased CO<sub>2</sub> is the primary driver of greenhouse effect

• CO2 traps infrared radiation (thermal radiation). Proven by laboratory experiments and satellites (satelit data from 1970; direct experimental evidence) that find less heat escaping out to space over the las few decades.





# Climate change controversy

- Positive/negative feedbacks examining different period throughout Earth's history shows that feedbacks amplify or diminish any initial warming.
- Positive feedback
  - Warming keeps more water in the air and more wapour traps more heat
  - Warming releases carbon (methane) in the arctic from thawing permafrost. Or from hydrates (water ice containing methane in its structure).
  - Drying rainforest, forest fires. Desertification.
  - Albedo feedback.
- Negative feedback
  - More water vapour causes more clauds, reflecting sunlight.
  - Increase in the overal amount of greenery increased plants photosynthesis

### Climate change controversy

- About 700 800 Gt of CO2 in the atmosphere. Up to 1700Gt in permafrost in Siberia, Canada...
- Higher temperature leads to faster thawingwhich leads to faster releasing of CO2, which leads to increase in the temperature....
- With about 3°C expected average temperature increase (=  $6^{\circ}$ C in arctic, due to the uneven distribution) about 30 85% of near-surface permafrost is to melt.
- According to the UNEP, by 2100 about 43-135Gt of CO2 may be released, by 2020 something between 246-415Gt.
- Unreliable data about deeper layers of permafrost.
- Plus methane.




#### Summary

- Earth's climate has undergone changes over long periods of time (several ice ages, period of warming).
- Previous changes were dramatic but gradual (thousands of years).
- Today's change is extremely fast and the pace is increasing. Until 250 years ago the highest rate of temperature increase recorded was approximately 0,003°C/y. For the last ten years, it is 0,017°C
- Global warming vs. climate change. The first suggests that Earth's climate is warming on average, but it is not fully true. Factors such as precipitation and evaporation are also changing. And these changes often affect climate patterns elsewhere in the world.

### Summary

- There is scientific consensus on
  - correlation between the concentration of  $CO_2$  and temperature.
  - that humans release anthropogenic compounds into the environment, resulting in previously unseen rises in atmospheric gas concentrations and temperature.
- There is continuous debate on
  - the proportion of changes caused by this anthropogenic compound vs. other causes.







#### Climate change impacts

- Melting ice
  - The vast majority of the world's glaciers are melting faster than are replenished.
  - 1/3 of North Pole's ice sheets melted since 90s.
- Accelerated sea level rise, increase coastal flooding
  - 20 cm in the last century (40% thermal expansivity, 60% melting of the land ice).
  - Actual rate 3mm/y.
  - Problem for low-lying communities (such as Bangladesh).
  - Problem from some smaler islands (such as Kiritimati in the Pacific, no more than 2m above sea level at the highest point)




#### Climate change impacts

- Increase in extreme weather events
  - Climate change increases certain types of extreme weather events heat waves, coastal flooding, extreme precipitation events, more severe droughts.
  - Temperature average kinetic energy of the molecules within a substance = the more radiation trapped in the atmosphere the higher temperature is.



#### umber of Climate-related Disasters Around the World (1980-2011)



## Climate change impacts

• Health impacts

- Increased air pollution, a longer and more intense allergy seasons, the spread of insect-borne diseases, more frequent heat waves, flooding = costly risks to public health.
- Food problems and water
  - According to IPCC  $1^{\circ}C = 65$  million people starving.
  - Increase of the temperature of more than  $2^{\circ}C = 3$  billion people without water supply.
  - Between 18-35% of plant and animal species is committed to extinction by 2050 (oceans are absorbing much of the  $CO_2$  in the air, which leads to ocean acidification destabilising the whole oceanic food chain). An estimated 1 billion people depend on the ocean for more than 30% of their animal protein.
  - Climate refugees.
- And others...




# Climate refugees

- No legal internationally recognised definition of climate refugee.
- Not covered by UN 1951 Refugee Convention.
- Numbers are increasing. World Bank expects only in the regions of Sub-Saharan Africa, South Asia, and Latin America climate change could fore more than 143 million people to move internaly, which eventualy be reflected in general migration.
- Problem in numbers and resources?




#### Climate refugees

• Global Compact for Safe, Orderly, and Regular Migration





#### The Relationship Between the Level of Greenhouse Gas Stabilization and Eventual Temperature Change



**Eventual Temperature change (relative to pre-industrial)** 



### Possible Effects of Climate Change

|                         | Eventual Temperature Rise Relative to Pre-Industrial Temperatures   |   |  |   |   |  |  |
|-------------------------|---|---|--|---|---|--|--|
| Type of Impact          | 1°C   | 2°C   | 3°C  | 4°C   | 5°C   |  |  |
| Freshwater<br>Supplies  | Small glaciers in the<br>Andes disappear,<br>threatening water supplies<br>for 50 million people                          |   | Serious droughts in<br>nesouthern Europe every 10<br>years. 1–4 billion more<br>people suffer water<br>shortages | Potential water supply<br>decrease of 30–50% in<br>southern Africa and<br>Mediterranean         | Large glaciers in Himalayas<br>possibly disappear, affecting<br>¼ of China's population |  |  |
| Food and<br>Agriculture | Modest increase in yields<br>in<br>temperature regions  | Declines in crop yields in<br>tropical regions (5–10% in<br>Africa) | 150–550 million more<br>people at risk of hunger.<br>Yields likely to peak at<br>higher latitudes                | Yields decline by 15–35% in<br>Africa. Some entire regions<br>out of agricultural<br>production | n Increase in ocean acidity<br>possibly reduces fish stocks                             |  |  |
| Human Health            | At least 300,000 die each<br>year from climate–related<br>diseases.<br>Reduction in winter<br>mortality in high latitudes |   | d 1–3 million more<br>potentially people die<br>annually from malnutrition                                       | Up to 80 million more<br>people exposed to malaria i<br>Africa                                  | Further disease increase and<br>nsubstantial burdens on<br>health care services         |  |  |
| Coastal Areas           | Increased damage from<br>coastal flooding   | Up to 10 million more<br>people exposed to coastal<br>flooding      | Up to 170 million more<br>people exposed to coastal<br>flooding  | Up to 300 million more<br>people exposed to coastal<br>flooding                                 | Sea-level rise threatens<br>major cities such as New<br>York, Tokyo, and London         |  |  |
| Ecosystems              | At least 10% of land<br>species facing extinction.<br>Increased wildfire risk   | 15–40% of species potentially face extinction                       | 20–50% of species<br>potentially face extinction<br>Possible onset of collapse<br>of Amazon forest               | Loss of half of Arctic tundr<br>Widespread loss of coral<br>reefs                               | aSignificant<br>extinctions across the globe<br>CENTER FOR<br>RGY STUDIES               |  |  |

#### Climate change impacts by region

|        | People affected each year<br>by 2080s by storm surges<br>with sea-level rise of<br>about 38cm assuming<br>constant protection<br>mechanisms (evolving<br>protection mechanisms) <sup>a</sup> | Estimated climate<br>refugees due to sea-<br>level rise (slr) <sup>b</sup> | Vulnerability to<br>tropical cyclones <sup>c</sup>   | People at risk of wa-<br>ter stress by 2085<br>due to a temperature<br>increase of 2–3<br>(depending on<br>population level) <sup>d</sup> | Estimates related to<br>drought and water<br>stress <sup>e</sup>   | Additional num-<br>ber of people at<br>risk of hunger by<br>the 2080s <sup>f</sup> |
|--------|--|--|--|---|--|--|
| Africa | Southern<br>Mediterranean:<br>13 million (6 million)   | Egypt: 12 million<br>by 2050   | Southeast<br>Africa: low to<br>moderate risk   | North Africa:<br>155–599 million  | <ul><li>14 African</li><li>countries currently</li><li>experience water</li><li>scarcity.</li><li>Expected to rise to</li><li>24 countries by</li><li>2030</li></ul> | Total: 23-200  |
|        | West Africa: 36 million<br>(3 Million)   | Nigeria: 6–11 mil-<br>lion by 2050   |  | South and<br>East Africa:<br>15–529 million   |  |  |
|        | East Africa: 33 million<br>(5 million)   |  |  | West Africa:<br>27–517 million  |  |  |
| Asia   | South Asia: 98 million (55 million)  | Bangladesh: 26<br>million by 2050  | Major urban cen-<br>ters: moderate toSouth Asia:<br>39–812 millionhigh riskWest Asia:<br>95–492 millionate riskCentral Asia: |   | Millions at risk due<br>to the glacier melt<br>in the Himalayas.   | West Asia:<br>5–134 million<br>Southeast Asia:<br>2–44 million                     |
|        | Southeast Asia: 43 mil-<br>lion (21 million)   | China: 73 million  |  |   |  |  |
|        |  | India: 20 million<br>by 2050   |  | 50–60 percent of<br>world population  | 2-44 mmon  |  |
|        |  |  | East Asia: moder-<br>ate to high risk  | 14–228 million<br>East Asia:  | live in the larger<br>Himalaya-Hindu<br>Kush region and<br>could be affected by<br>water stress  |  |
|        |  |  |  |   |  |  |
|        |  |  | South East Asia:<br>moderate to high<br>risk   | 41–1577 in worst<br>case scenario   |  |  |



#### Climate change impacts by region

| Latin<br>America          | N/A   | Venezuela: 56,000<br>assuming 1m slr<br>and no adapta-<br>tion measures<br>Uruguay: 13,000<br>assuming 1m slr<br>and no adapta-<br>tion measures | Central America:<br>low to high risk<br>Northern Latin<br>America: low risk | Central America:<br>5–246 million<br>South America:<br>72–272 million in<br>the worst-case sce-<br>nario | Glacier melt in the<br>South American<br>Andes could cause<br>water stress under<br>37 million people<br>by 2010 and 40 mil-<br>lion by 2050 | Total:<br>5–85 million |
|---------------------------|---|--|---|--|--|------------------------|
| Small<br>island<br>states | Caribbean: 1,350,000<br>(560,000)   | 1 million  | Caribbean: low to<br>moderate risk  | Caribbean:<br>0–73 million   | Water availability<br>could become too<br>low during low<br>rainfall seasons   | N/A.                   |
|                           | Indian Ocean:<br>920 thousand<br>(460,000)<br>Pacific: 290,000<br>(160,000) |  | Indian Ocean: low<br>to moderate risk                                       |  |  |                        |
|                           |   |  | Pacific: low to high<br>risk  |  |  |                        |
|                           |   |  |   |  |  |                        |



#### Sources

- IEA: CO<sub>2</sub> Emission from Fuel Combustion
- IPCC: Climate Change 2013: The Physical Science Basic
- Wagner, G.; Weitzman, M.L.(2015: Climate Shock: The Economic Consequences of a Hotter Planet
- Figueres, Ch.-Ivanova, H.M.: Climate Change: National Interests or a Global Regime?
- IEA: CO<sub>2</sub> Emission from Fuel Combustion
- Carbon Brief
- Center for Climate and Energy Solutions
- Harris, J.M.; Roach, B.; Codur, A-M.(2017): The Economics of Global Climate Change. A GDAE module
- Ritchie, H.; Roser, M.: CO2 and other Greenhouse Gas Emissions. Our World in Data
- Biermann, F.; Boas, I.(2010): Preparing for a Warmer World: Towards a Global Governance System to Protect Climate Refugees
- Burrows, K.; Kinney,L.(2015): Exploring the Climage Change, Migration and Conflict Nexus
- Carbon Pricing Leadership
- UNEP: Policy Implications of Warming Permafrost
- Sceptical Science: What does past climate change tell us about global warming?
- IJRC (2018): UN Member States adopt global compacts on migration and refugees.
- World Bank (2018): Groundswell: Preparing for internal climate migration.
- FiveThirtyEight (2018): The World Isn't Ready for Climate Refugees.

