Types of prevention Risk assessment

Mgr. A. Peřina, Ph.D.

Dept. of Public Health

Faculty of Medicine, Masaryk University

Health

• A state of complete physical, mental and social well-being and not merely the absence of disease or infirmity (WHO, 1948).

 Health is neither born nor cultivated in hospitals but it arises and developes wherever people live, play and work, relax and grow old.

- Pillars of health care:
 - 1. Health education
 - 2. Disease prevention
 - 3. Health protection + health promotion

Health protection and promotion


- Health protection: summary of activities and measures focused on creation and protection healthy living and working conditions aimed at the prevention of infectious and non-infectious diseases
- Health promotion: summary of activities helping people to strengthen and improve their health and control their determinants of disease
- Sectors involved: health sector, ministries of environment, local development, agriculture, industry, trade, labor and social issues, transport, culture, ministry of defense, ministry of education and sport...

Prevention


Type of prevention	Tasks	Responsibility	
Primary	Influencing disease determinants and reducing health risks	Ministries of Health, Environment, Agriculture, Social Affairs, Education etc.	
Secondary	Preventive examinations, screening programs	Ministry of Health Ministry of Health, M. of Social Affairs	
Tertiary	Treatment of diseases and mitigation of their consequences		
Quartery	Mitigate the effects of unnecessary or excessive treatment interventions	Ministry of Health	

 BreatheLife mobilizes cities and individuals to protect people's and Planet health from the effect of air polution.

Over 80% of the world's cities have air pollution levels over the WHO guideline for safe air



Antibiotic resistance:
 new WHO
 recommendations are
 trying to help
 maintaining the
 effectiveness of
 antibiotics used in
 human medicine by
 limiting the unnecessary
 use in animals.



Global Hearts" is an initiative to averting the global threat of cardiovascular disease, the world's leading causes of death.




• "WHO's Global School Health Initiative", also known as "Health Promotion School": An initiative to promote health and education at local, national, regional and global level. The aim is to improve the health of students, school staff, families and other community members through schools.



 Global Action Plan "Healthy enterprise" addresses all aspects of workers' health: primary prevention of occupational hazards, protection and promotion of health at work, working conditions and health services for employees of enterprises.



 SaveLIVES: road safety package, evidencebased inventory focusing on speed limits, transport infrastructure, vehicle safety, traffic law and regulations and actions increasing the chance of survival after a traffic accident.



The preventive and therapeutic components of medicine are complementar y to one another.



"What fits your busy schedule better, exercising one hour a day or being dead 24 hours a day?"

Why prevention when we can treat the disease?

- Effective prevention reduces the incidence and prevalence of serious diseases.
- In most cases, treatment has side effects that can be avoided by effective prevention.
- Consistently healthy individual is economically active.
- Consistently healthy individual is less sensitive to social, cultural and other differentiation.
- Satisfactory environment fulfills well the functions of production, recreation and regeneration.
- The zero option means the growing cost of medical technology, the possibilities of which can be (once) exhausted (eg. antibiotics and current resistance, organ transplants and ethical issues etc.).

Health risks assessment



Hazard vs Risk

- Hazard
 - Characterizes properties of agents
 - Pathogenicity, toxicity...

... It may, it could...

- Risk
 - Determines the probability of adverse changes in health status
 - It is a mathematical function of hazard.

Probability

- Meaning in conversation
- Relative frequency of the phenomenon

 The result can be predicted on the basis of a statistical analysis of a large number of iterations.

However:

- the life expectancy in the population does not indicate the length of an individual's life.
- the likelihood of genetic diseases do not indicate the occurrence of this disease in particular newborns



Pifalls in risk assessment

- Risk assessment (among others) = probability of hazard application + emotions
 - the public knows very little about probability and overestimates its importance
 - experts (mostly) know very little about emotions; experts must be fully aware of the fact, that
 - emotions are measurable, as well as the probability is
 - emotions can be affected as well as the probability
 - emotions are a legitimate part of the risk

Prioritization of health risks

• Why?

There are a lot of risks that occur in the human environment, but only some of them have a direct impact on human health.





Risk Assessment

Attention focuses on human!

Hazard identification: can the agent (or active factor) harm health?

- 1. Dose effect relationship: what is the numerical relationship between the exposure and the effect on health?
- 2. Exposure assessment (evaluation): how important is the contact of the individual / population with the agent?
- 3. Risk characterization: can the assumption of an adverse effect on health be confirmed?

Types of hazard (will be discussed in more detail in specific


seminars)

- Biological agents
 - Pathogenic microorganisms (see epidemiological seminars)
 - Non-pathogenic microorganisms related to health
 - Toxins as by-products of decomposing and primarily non-pathogenic microflora (fungi and aflatoxins)
- Chemicals
- Irritating, toxic, mutagenic, teratogenic and carcinogenic effects
- Physical factors
 - Noise, vibration
 - Nonionizing and ionizing radiation: Special features of therapeutic use: benefit / risk ratio
 - Microclimate, unilateral strain of muscle groups

Types of dose-effect relationship

- Agents with threshold effect
 - Existence of a safe dose
 No Observed Adverse Effect Level
- Agens with no threshold effect
 - Non-existence of a safe dose, the probability of an adverse health effect increases
 - "Slope factor" defined as the size of the inclination angle of no-threshold line (see the figure)

Dose-Effect Curves



Exposure assessment


- Dose offered
 - It corresponds to the concentration of the agent in the environment (ie in air, water, food, soil), converted to a unit of mass, volume or area of the matrix.
- Dose absorbed
 - It depends on the speed of diffusion and capacity of receptor.
 - Ingestion, inhalation, contact with skin or mucous membranes
 - Comment: besides the concentration, the duration of exposure can also determine the size of the effect.
- Effective dose
 - Defined by the concentration of agents in the target organ

Exposure assessment – methods

- Indirect methods
 - 1. Environmental monitoring: The amount of agent in the matrix multiplied by the average matrix intake by the exposed person:
 - Average lung volume (22 m3/person/day)
 - Average water consumption per person (1,9 liter/day)
 - Amount of food consumed per person (e.g. Food pyramid)
 - The average length of stay in the swimming pool
 - Inaccuracy! Interindividual differences are significant!
 - 2. Exposure scenario or questionnaire survey: A rough estimation of the exposure can be specified, most often in a well defined population group (typically school pupils, members of the army ...)

Estimation of individual exposure based on the food pyramid may be difficult.





CLOSE ENOUGH.

Exposure assessment – methods II

- Direct methods
 are preferred but are generally less accessible.
 - Personal monitoring:
 - 24 hrs recall, double portion method...
 - Personal dosimetry healthcare workers
 - Biological monitoring
 - Biomarkers of exposure (genotoxicology of DNA adduct)
 - Biomarkers of effect (measurable pathophysiological changes in organs)
 - Biomarkers of sensitivity (measurable susceptibility to health impairment)

Risk characterization

- 1. Harmful to health has not been confirmed
- 2. Exposure to harmful factor reduces the level of well-being (health in a broader sense)
 - Example: The source of environmental noise has forced the use of space (more demanding activities are moved to a quieter part of the building).
- 3. Exposure to a harmful factor poses a threat to health in the longer term, with the factor being considered at most as one of several disease factors (long-term and multifactorial health effects)
- 4. Exposure to harmful agents poses an immediate threat to human health or lives
 - (See the Czech "methanol affair" in 2012.)

Risk matrix

Graphic tool for expressing the risk according to the probability of exposure and the size of the effect.

		Impact					
		Negligible	Minor	Moderate	Significant	Severe	
— Likelihood ——	Very Likely	Low Med	Medium	Med Hi	High	High	
	Likely	Low	Low Med	Medium	Med Hi	High	
	Possible	Low	Low Med	Medium	Med Hi	Med Hi	
	Unlikely	Low	Low Med	Low Med	Medium	Med Hi	
	Very Unlikely	Low	Low	Low Med	Medium	Medium	

Uncertainty in health risk assessment

- Errors in exposure estimation
 - Most common
 - Due to the complexity of the presence of the agent in the matrix (water, soil, air, food, occupational exposure)
 - The size of individual effective dose in the target organ of the individual
- Interspecies extrapolation
 - In many cases, toxicological data exists only on animal or tissue models
 - Can be addressed with the development of epidemiological methods of work

Epidemiology


- Method of work used to study the distribution of determinants of diseases in the population.
 - Epi demios (Greek) = Among the people
- Descriptive, analytical, experimental and interventional epidemiology use statistical methods; a separate field of biostatistics is being developed.

Focus on the study of infectious and non-infectious diseases.

Epidemiology in health risk assessment


Benefits

- fundamental role in the scientific concept of protecting public health
- no need for interspecies extrapolation
- exposure to the harmful agent may be studied in context
- no need to extrapolate experimental data to the low dose range



Epidemiology in health risk assessment


- Problems: transferability of results
- Internal validity of epidemiological studies
- Hill's criteria of causality (Sir Austin Bradford Hill, 1897 1991)
 - ✓ Strength of association: Even a weak association does not rule out causality if it is weakened by unrecognized confounders
 - ✓ Consistency: Inconsistency with other epidemiological studies does not exclude causality, the effect may only occur under special circumstances
 - ✓ The causality does not assume the specificity of the effect
 - √ The chronology (time sequence) of exposure and effect is a prerequisite!



Epidemiology in health risk assessment II.

Hill's criteria of causalitycontinuation

- ✓ Biological gradient: its absence does not exclude causality, many dependencies can be in the shape of the letter "J"
- ✓ Biological plausibility: the inability of pathophysiological explanation of the phenomenon may be just the result of the current scientific knowledge
- ✓ Coherence: Incompatibility with "established" theories does not exclude causality
- ✓ Experimental evidence: its absence does not exclude causality, as the experiment can also be prevented by ethical reasons
- ✓ Analogy: its absence can only be a manifestation of a lack of scientific imagination



Epidemiology in health risk assessment III.

- Bias in epidemiological studies
 - ✓ an error in estimating the effect being studied as a result of poor planning or execution of epidemiological studies. It may cause the distortion of the results in terms of both positive or negative.
- Basic types of bias in epidemiological studies
 - ✓ Selective bias: e.g. respondents willing to collaborate are not a representative sample of the population
 - ✓ Information bias is a sign of poor classification in a statistical survey
 - ✓ Confounding bias is a manifestation of the interfering "third" factor
 of the disease.

Health risk assessment in doctor's everyday practice

Not only in prevention, but also in the diagnosis and treatment of diseases, doctors should ask the following questions (according to the US EPA):

- ✓ What symptoms can be causal in terms of exposure to environmental or occupational hazards?
- ✓ Is there a safe exposure level for the selected agent?
- ✓ Is there a possibility that some people will be exposed to significantly different exposure doses?
- ✓ Is it an exposure of an increasingly vulnerable (susceptible) population group (such as children, pregnant women, sick people, the elderly, people with occupational exposure)?

Public health protection and qualitative research

- Qualitative research always just as a complement to epidemiological methods of work
- It enables us to understand the social, cultural, economic and behavioral aspects of public health protection

- Epidemiological methods: "How many? How much?"
 - ✓ Calculation of frequency, confidence intervals, the probability of the estimation error (magic "p" value)
- Qualitative research: how and why?
 - √ From the Latin word "Qualis" (= How? What?)
 - √ Verbal analysis of relationships and contexts





Presented by: Susan Abbott

Conclusions

Lost health can be restored medically.

But:

 Creation of health, health protection and health promotion in some ways exceeds the capacity of clinical medicine.

- The basis for the protection and promotion of health is a health risk assessment process built on a scientific basis.
- Principles of health risk assessment are very useful in clinical practice.