

# Urinary tract diseases

## Male genital system

V. Žampachová  
I. ÚP



# Congenital anomalies

- **Extrophy of the bladder** – non-closure of anterior wall (bladder +/- abdominal, symphysis pubis; infections incl. pyelonephritis; ↑ risk of ca)
- **Congenital stenosis** – ureteropelvic junction, double/bifid ureter → hydronephrosis
- **vesicoureteral / ureteropelvic reflux**



# Urinary tract obstruction

- increased susceptibility to urolithiasis
  - increased susceptibility to infection
  - risk of hydronephrosis
- 
- Combination of inborn + acquired risk factors





# Vesico-ureteric reflux

- Incompetence of the vesico-ureteral valve
- Combination of congenital defect (short intravesical part of ureter, 1-2% of children)
- ↓ ureteral contractility in infection
- acquired in bladder atonia (spinal cord injury)



# Vesico-ureteric reflux








cystoureterogram

# Intrarenal reflux

- Upper and/or lower renal papillae
- Progression of infection into the kidney tissue
- No reflux – usually no ascending infection






# Obstruction causes

- **Intrinsic** – luminal obstruction (stone, blood clot, necrotic papilla, tumor or its part)
- **Wall stenosis or dysfunction** (inborn, inflammation, postinflammatory, tumor, ...)
- **Extrinsic** – external compression, some causes common for both sexes, some different




# Obstruction causes

- **In males:** prostatic hyperplasia, prostatic ca, urethral stenosis, phimosis + complications
- **In females:** pregnancy, cervical ca (+ therapy), uterine myoma, ovarian tumor, uterine prolapse
- **in both:** chronic inflammation/fibrosis (retroperitoneal fibrosis), tumor (colorectal ca, LN,...), aortic aneurysm



Massive hematuria from renal calculi,  
tumors, or papillary necrosis



# Urinary calculi


- Usually of renal origin
- Stones > 5 mm cannot pass into ureter
- Renal colic – pain + spasms during the passage of a stone along the ureter
- Chronic dull pain lumbar – lower pelvic region
- ↑ risk of obstruction
- Repeated infection
- Epithelial transformation: squamous metaplasia




# Urinary calculi


- **Calcium** containing stones: commonest, laid down in an acid urine.
- **Complex triple phosphate** stones: often associated with urinary infection, in an alkaline urine.
- **Mixture of uric acid and urate-uric acid** stones, 20% of patients with gout, in an acid urine. Pure uric acid stones radiolucent.
- **Cystine** stones: in primary cystinuria, important in childhood.

# Nephrolithiasis



## Pyelolithiasis in situ






# Urocystolithiasis




# Urate nephropathy

- Hyperuricemic disorders (urate crystals formation) may lead to 3 forms of injury:
- **Acute urate nephropathy** in patients with haematologic malignancies, commonly during chemotherapy (extensive cell breakdown – release of nucleic acids – urate crystals in tubules – acute renal failure)
- **Chronic urate nephropathy** – in gout. Urate crystals surrounded by foreign body giant cells, tubulo-interstitial nephritis
- **Urate stones**

# Urate crystals



# Urate crystals - polarization



# Acute pyelonephritis


- Common purulent renal inflammation, bacterial infection by *Escherichia coli*, *Proteus*, *Klebsiella*, *Enterobakter*
- **Ascending** infection by urine reflux in urinary tract inflammation
- **Descending (haematogenous)** infection in septicaemia, rare



# Acute pyelonephritis

## HEMATOGENOUS INFECTION

Common agents:  
*Staphylococcus*  
*E. coli*




# Acute pyelonephritis


- Facilitated by DM, gout, all causes of obstructive uropathy (e.g. nephrolithiasis, tumors, urinary tract anomalies incl. vesicoureteric and intrarenal reflux)
- Instrumental interventions (catheterization, cystoscopy)
- GROSS: enlarged kidney, cortical and medullary abscesses
- MICRO: purulent neutrophilic exudate in tubules and interstitium, oedema



# Acute pyelonephritis (4)




# Acute pyelonephritis



# Acute pyelonephritis

- Pyonephros
- Papillary necrosis in diabetics
- Peri- and paranephritic abscess




Papillary  
necrosis





# Chronic pyelonephritis

- Uni- or bilateral chron. tubulo-interstitial renal inflammation with scarring
- 10-20% end-stage kidney
- **Obstructive PN** - repeated infections
- **Reflux nephropathy –vesicoureteric and/or pelveorenal reflux** (from lower and upper pole calyces into renal parenchyme)





Chronic pyelonephritis




5 cm


# Chronic pyelonephritis




# Chronic pyelonephritis



# Chronic pyelitis



# Xanthogranulomatous pyelonephritis



- Uncommon form of chronic pyelonephritis with accumulation of foamy macrophages in interstitium
- Yellowish focal lesions in macroscopy, diff. dg. x renal carcinoma



# Hydronephrosis, hydroureter

Renal tissue  
atrophy, renal  
insufficiency



# Tumors of the renal pelvis

**Primary:** transitional cell neoplasia, mostly papillary  
transitional cell carcinoma (invasive, non-invasive)

Rarely other types of carcinoma (squamous cell,  
neuroendocrine, adenocarcinoma)

Other types (mesenchymal, melanoma, ...)


**Secondary:** local progression from the kidney  
metastasis




# Transitional cell ca

- in pelvis or ureter less common than bladder
- possible multiple dysplastic foci
- histopathology similar to bladder ca
- possible porogenous seeding
- invasion into kidney and surrounding tissues
- diff. dg. x renal cell ca


## Invasive urothelial carcinoma – renal pelvis



# Transitional cell ca of the renal pelvis



# Transitional cell ca of the renal pelvis



Concurrent urothelial ca of pelvis + bladder possible



# Ureter

- **congenital anomalies** (double/bifid ureter, obstruction of the ureteropelvic junction, lesion of the vesicoureteric junction, diverticula)
- **obstruction**
- **inflammation** (acute, chronic incl. ureteritis cystica - preneoplastic)
- **neoplasia and pseudotumorous lesions** (transitional cell tu, squamous cell ca, fibroepithelial polyp, etc.)




# Ureteritis cystica

- Special form of chronic ureteritis
- Numerous cysts in the ureteral mucosa, solid or cystic nests of transitional or metaplastic glandular epithelium.
- Terminal stage of chronic inflammations
- ↑ risk of ca




## Chronic ureteritis cystica



# Cystic ureteritis



# Cystic ureteritis




# Tumors, tumor-like lesions

- Fibroepithelial polyp: small, loose stroma + epithelium
- Benign: rare, mesenchymal tumors
- Malignant: transitionall cell carcinoma

progression of malignancy  
from surrounding tissues



# **Transitional cell carcinoma of the ureter**



# Urinary bladder

- **congenital anomalies** (exstrophy, diverticula, persistent urachus)
- **cystitis** (acute, chronic, special forms – malakoplakia, tbc, schistosomiasis,etc.)
- **metaplasia** – usually in cystitis (c. cystica, c. glandularis), squamous m.- leukoplakia
- **tumors and pseudotumorous lesions**
- **miscellaneous** (calculi, fistulae, prolapse)



# Exstrophy


- nonclosure of anterior bladder + abdominal wall incl. missing layers
- bladder may be opened to the outside
- ascending renal infection
- epithelial metaplasia, ↑ risk of ca
- commonly in combination with other congenital anomalies

# Diverticula

- **Congenital:** uncommon, solitary. Wall defect; intrauterine urinary obstruction
- **Acquired:** multiple. Most common in prostatic hyperplasia, concurrent with cystitis
- ↑ risk of infection, stones, perforation



## Urocystolithiasis + chronic cystitis + diverticula



# Acute cystitis

- highly common in females (short urethra, perineal connection with anus)
- mostly fecal bacteria, mixed flora
- risk factors – urine pH, hormonal status, iatrogenic
- usually purulent (leucocytes, blood in urine), urging, pain; may have systemic signs
- complications – ureteral spread, ulcers, rare phlegmona, pseudomembranous infl



# Haemorrhagic cystitis


Klepnutím lze upravit styly předlohy textu.

Druhá úroveň


Třetí úroveň

Čtvrtá úroveň

Pátá úroveň



# Ulcerative cystitis




# Chronic cystitis

- epithelial transformations – polyps, epithelial hyperplasia (Brunn's nests, reactive atypia – x neoplastic), metaplasia (squamous - leukoplakia, glandular)
- neoformation of lymphatic follicles in stroma
- in obstruction + muscular hyperplasia, diverticuli
- acute exacerbations, stone formation
- may be risk factor for neoplasia
- diff. dg. x neoplasia




# Leukoplakia




Copyright © 2002, Elsevier Science (USA). All rights reserved.




# Chronic cystitis with squamous metaplasia



# Chronic cystitis with squamous metaplasia



# Glandular cystitis



# Granulomatous cystitis

- **specific**
  - parasites
  - TB, incl. BCG vaccine as treatment for bladder ca
  - other
- **nonspecific**
  - foreign body reaction, incl. post-treatment (endoresection)



# Granulomatous cystitis


# Schistosomiasis

- chronic parasitic inflammation
- endemic in Africa, Middle East
- possible traveller infection, worm lives for up to 20 years
- granulomatous reaction to eggs + fibrosis
- strictures, squamous metaplasia
- ↑↑ risk of squamous ca



# Schistosomiasis

# Schistosomiasis



# Transitional cell (urothelial) tumors

## exophytic:

- papilloma (benign), inverted papilloma
- papillary urothelial neoplasm of low malignant potential (PUNLMP)
- non-invasive papillary urothelial carcinoma – low grade, high grade
- invasive urothelial carcinoma – low grade, high grade



# **Transitional cell (urothelial) tumors**

- **Flat lesions**

## **Intraepithelial neoplasia**

- dysplasia – low grade intraurothelial neoplasia (LG IUN)
- high grade (HG IUN) - carcinoma in situ (CIS)

## **Invasive carcinoma**




# Bladder epithelial tumors - other

- squamous cell carcinoma
- adenocarcinoma
- small cell carcinoma (neuroendocrine ca)
- mixed ca
- secondary tumors – prostatic ca, cervical ca, rectal ca




# Bladder non-epithelial tumors


- Melanocytic
- Mesenchymal (benign, malignant sarcomas)
- Other




Papilloma-  
papillary carcinoma



Invasive papillary  
carcinoma



Flat noninvasive  
carcinoma




Flat invasive  
carcinoma

**Figure 21–9. Four morphologic patterns of bladder tumors.**

from Robbins' Pathology of Diseases





# Urothelial papilloma



- Rare
- Solitary
- Exophytic / endophytic (inverted)
- Normal urothelium – no atypia, usual number of layers, superficial differentiation od umbrella cells

# Papillary urothelial neoplasm of low malignant potential

- Slight increase in the number of cell layers, normal stratification, minimal increase in nuclear size and density
- Basal sporadic mitotic activity
- Risk of recurrence, possible progression into ca



# Papillary urothelial neoplasm of low malignant potential




# Non-invasive papillary urothelial carcinoma


- low grade
- high grade
- cytonuclear atypias of a carcinoma
- no stromal invasion
- histological code of a ca in situ (8130/2)  
for low grade ca
- High grade ca – code for carcinoma  
(8130/3)




# Non-invasive papillary urothelial carcinoma, low grade



# Non-invasive papillary urothelial carcinoma, high grade



# Non-invasive papillary urothelial carcinoma, high grade



# Invasive urothelial carcinoma


- ex flat urothelial ca in situ
- ex non-invasive papillary urothelial ca  
(papillary component commonly present)
- variable grade
- invasion into deep bladder structures  
(muscle layer), adjacent tissues/organs  
(fat, ureters, prostate gland, ...)




# Invasive urothelial carcinoma

- 90% ca in the bladder (rare squamous cell ca or adenocarcinomas), mostly in 50 – 80 yrs old
- etiology: smokers, professional (anilin dyes, plastics industry), analgesic abuse, irradiation. Sq. cell ca in chronic inflammation (schistosomiasis)
- Differentiation grade G1 – G3
- asymptomatic, possible haematuria

# Invasive urothelial carcinoma - staging




# Bladder carcinoma




# Bladder carcinoma




# Invasive urothelial carcinoma




# Carcinoma with glandular transformation




# Mucinous adenocarcinoma



# Squamous cell carcinoma



## Malignant melanoma in the bladder



# MALE GENITAL SYSTEM

# Penis, scrotum

# Congenital lesions

- **Epispadia:** less common, incomplete fusion of urethra, dorsal opening, may be a part of bladder exstrophy
- **Hypospadias:** more common, opening on inferior part of penis (glans, shaft, perineum)
- **Phimosis:** diminished size of prepuce opening, rare inborn, more common acquired – inflammation, scarring, ! ca



# Circulatory disorders

- Edema
- corpora cavernosa thrombosis,
- gangrene (uncommon)



# Balanoposthitis

- Glans + prepuce
- Sexually transmitted diseases – STD: syphilis, gonorrhea, chancroid, herpes
- Non-specific infection: candida, pyogenic bacteria, anaerobic bacteria
- Poor hygiene – repeated infection
- Phimosis, chronic irritation




# Viral infections


- inflammation +/- pseudotumorous lesion (molluscum contagiosum)
- benign tumors: **condyloma accuminatum** – HPV 6, 11) squamous cell papilloma
- preneoplastic lesion/intraepithelial neoplasia: dysplasia mild – moderate – severe – carcinoma in situ
- HPV risk factor




# Penile condyloma



# condyloma accuminatum - HPV





## Molluscum contagiosum

Copyright © 2002, Elsevier Science (USA). All rights reserved.




# Balanitis xerotica obliterans

- chronic inflammatory disorder (= lichen sclerosus)
- epithelial hyperkeratosis, atrophy, inflammatory infiltrate
- possible risky terrain for ca



# Lichen sclerosus



# Tumorous lesions

- Pseudotumors – fibromatosis (Peyronie's disease) – deformation possible
- Fibroepithelial polyp
- Benign tumors – HPV papilloma, adenoma



# Malignant tumors

- Skin tumors (squamous cell carcinoma, melanoma)
- Mucosa tumors (various types of squamous cell carcinoma incl. exophytic verrucous ca)
- Urethral tumors
- Other primary tumors
- Secondary tumors



# Squamous cell ca in situ

**Bowen's disease / erythroplasia of Queyrat on the glans: white / red focus on the mucosa**

**bowenoid papulosis (multiple, HPV 16, non-progressive)**



# Invasive squamous cell ca

geography (Latin America, East Asia)

circumcision - protective factor ( $\downarrow$ HPV,  
carcinogenes in smegma)

risk factor – smoking, occupational (mineral oil,  
tar)

gross – ulcer, non-healing lesion


micro – sq. ca of variable type/grade




# Carcinoma of the penis



# Carcinoma of the penis



# Squamous cell carcinoma of the penis



# Prostate gland

- **infarction** (usually in hyperplastic prostate, repair with sq. cell metaplasia may mimic ca)
- **inflammation** (acute/chronic, a/bacterial, granulomatous)
- **benign nodular hyperplasia** (adenomyomatous) + related lesions
- **precancerous lesions, tumors** (PIN, adenocarcinoma, other)




# Inflammation


- **Acute bacterial:** ~ UTI, intraprostatic reflux of urine, iatrogenic – catheterization, cystoscopy, surgery. Pain, dysuria, fever.
- **Chronic bacterial:** repeated UTI, non-specific symptoms. Difficult to treat.
- **Chronic abacterial:** most common, no UTI, negative bacterial culture, reactive (+ prostatolithiasis), Chlamydia, ureaplasma
- **Granulomatous:** specific: Tb, BCG used for bladder ca; reactive



# Prostatolithiasis





# Chronic prostatitis



# Zonal predisposition of prostate diseases

Prostate zones



	Prostate zone		
	Peripheral	Transition	Central
Focal atrophy	High prevalence	Low prevalence	Medium-high prevalence
Acute inflammation	Low prevalence	None	Medium-high prevalence
Chronic inflammation	High prevalence	Low prevalence	Medium-high prevalence
Benign prostatic hyperplasia	None	High prevalence	Low prevalence
High-grade PIN	High prevalence	Low prevalence	Medium-high prevalence
Carcinoma	High prevalence	Low prevalence	Medium-high prevalence

Legend:

- High prevalence (Red)
- Low prevalence (Yellow)
- Medium-high prevalence (Orange)
- None (White)

- a Central zone
- b Fibromuscular zone
- c Transitional zone
- d Peripheral zone
- e Periurethral gland region

# Benign prostatic hyperplasia

- Common in older men, high incidence > 60 yrs
- Adenomyomatous hyperplasia – stromal (smooth muscle, fibrotic tissue) + glandular, alternating with atrophy, cystic and regressive changes. Two cellular layers – outer myoepithelial, inner secretory
- Gross: enlarged, nodular, firmer consistency, but demarcated.
- Main changes in central (periurethral) region



# Benign prostatic hyperplasia

- **Outcome:** partial → complete urethra obstruction, urinary residuum, risk of infection (+ ascending – pyelonephritis), bladder trabecular hypertrophy, hydronephrosis.
- Benign, but setting for possible preneoplastic changes
- Th: surgery, drugs



# Benign prostatic hyperplasia






FIGURE 17-34  
Growth of the prostate (left) and frequency of nodular hyperplasia (right). By 80 years of age, most men have benign prostatic hyperplasia.

# Benign prostatic hyperplasia - complications




# Benign prostatic hyperplasia




## Benign prostatic hyperplasia




## Benign prostatic hyperplasia




## Benign prostatic hyperplasia



## Parabasal hyperplasia



# Prostate – squamous cell metaplasia





# Prostatic carcinoma

- Adenocarcinoma: usually acinar, less common other types – ductal
- most common male ca (~1:6)
- late middle age – older males
- Highly variable course from clinically latent to extremely aggressive
- Recent studies: screening (PSA) generally not beneficial in asymptomatic males



# Prostatic carcinoma


- Important factors: race, family history, age, hormone level (androgens), environment
- Peripheral part (dorsal) – per rectum
- PIN: prostatic intraepithelial neoplasia – precursor lesion. High grade PIN important + included into pathological report
- Distinctive nucleoli, architectural changes, in PIN myoepithelial layer still present




# Prostatic carcinoma + hyperplasia




# Prostatic carcinoma



# Prostatic carcinoma



# Prostatic carcinoma




# Gleason histologic grading (WHO modification):

- grade of glandular differentiation, growth pattern
- combined score - dominant + secondary pattern in 5-grade system
  - grade 1 similar to normal prostatic tissue (uncommon in ca, in needle biopsy at least Gleason 3 by default)
  - grade 5 with solid, dissociated pattern
  - final combined score, commonly Gleason score 7 (4+3)

Now grade groups are used for grading

- Gleason score up to 6 = grade group 1
- Gleason 7 (3+4) = grade group 2
- Gleason 7 (4+3) = grade group 3
- Gleason 8 = grade group 4
- Gleason 9-10 = grade group 5


Higher grade – worse prognostic sign



# Prostatic carcinoma

- Local spread – into urinary bladder; diff. dg x high grade transitional cell ca, may be concurrent
- Metastatic spread: regional lymph nodes, hematogenous typical into bones – osteoplastic/osteosclerotic meta
- Symptoms – urinary commonly late, more due to prostatic hyperplasia; local spread; metastasis – bone pain, pathologic fracture


# Prostatic carcinoma




# Prostatic carcinoma - spine metastases



# Prostatic carcinoma - spine metastases, X-ray



## Prostatic ca metastasis in bone



# Testis, epididymis, cord


- congenital (cryptorchidism – atrophy, risk of neoplasia)
- regressive changes (atrophy, torsion)
- inflammation (orchitis – nonspecific acute/chronic, STD (gonorrhea, syphilis, chlamydia), mumps, tbc, idiopathic granulomatous)
- tumors

# Cryptorchidism


- Undescended testis
- 1 in 10 newborn males, usually descends during 1st year of life
- remains in inguinal canal or abdominal cavity – surgery necessary before puberty
- atrophy – infertility, germ-cell tumors




# Cryptorchidism



# Partial testicular atrophy



# Testicular atrophy + Leydig cell hypertrophy




# Intrascrotal swelling

- Commonly pathology of epididymis, tunica vaginalis
- Hydrocele – serous fluid in tunica vaginalis
- Haematocele – haemorrhage into tunica vaginalis
- Varicocele – dilated veins
- Spermatocele – epididymitis + cystic dilatation of ducts




# Intrascrotal swelling



# Intrascrotal swelling - hydrocele



# Intrascrotal swelling - varicocele



# Testicular torsion


- Spermatic cord turns around its own axis
- Haemorrhagic necrosis
- Acute severe pain, swelling
- More common in young/teenagers
- Immediate surgery necessary
- Without surgery loss of testicular function



# Torsion



# Testicular infarction



# Testis, epididymis inflammations


- epididymis >>> testis
- usually ascending from urinary tract and/or prostate
- caused by
  - gramnegative bacteria (children)
  - chlamydias, gonococcus (STI, adults)
  - E. coli (older adults)

# Testis, epididymis inflammations

- **Bacterial**
  - purulent → abscess, non-specific orchitis/epididymitis
- **Interstitial non-purulent orchitis**
  - mumps in adults
  - interstitial oedema + lymphocytes, plasma cells, macrophages
- **Granulomatous orchitis**
  - may be posttraumatic, v.s. autoimmune inflammation
  - non-caseating tuberculoid granulomas centered on tubules
  - firmer testicular mass (diff. dg. x tumor)
- **Spermatocytic granuloma**
  - in the head of epididymis due to rupture of tubules
  - reactive tuberculoid granulomas around spermatozoa




# Granulomatous orchitis



Tuberculoid granulomas.

# Testicular tumors



# **WHO pathologic classification of testicular tumors**

## **GERM CELL TUMORS**

**Derived from germ cell neoplasia in situ**

**Tumors of one histologic pattern**

- Seminoma
- Embryonal carcinoma
- Yolk sac tumor (embryonal carcinoma, infantile type)
- Polyembryoma
- Choriocarcinoma



# **GERM CELL TUMORS**

## **(cont.)**

- Teratomas
  - Mature
  - Immature
  - With malignant transformation

**Tumors showing more than one histologic pattern**

- Embryonal carcinoma + teratoma (teratocarcinoma)
- Choriocarcinoma + any other types
- Other combinations




# Germ cell tumors

Unrelated to germ cell neoplasia in situ

- Spermatocytic tumor (formerly spermatocytic seminoma)
- Teratoma, prepubertal type
- Yolk sac tumor, prepubertal type





## **SEX CORD-STROMAL TUMORS**

- **Well-differentiated forms**
- **Mixed forms**
- Leydig cell tumor
- Sertoli cell tumor
- Granulosa cell tumor
- **Incompletely differentiated forms**



# Testicular tumors

- Other types
  - Neuroendocrine tumors
  - Haematopoietic neoplasms
  - Tumors of collecting ducts and rete testis  
(adenoma, carcinoma)
  - Tumors of paratesticular structures  
(mesothelial tumors, tumors of the  
epididymis, ...)

# Testicular tumors

## Clinical features


- painless unilateral enlargement of testis
- secondary hydrocele
- symptoms from metastases
- retroperitoneal mass
- gynaecomastia




# Testicular tumors : histopathological report

- gross picture (incl. size)
- histological type
- presence of vascular / lymphatic propagation
- tumor staging (TNM classification)
- presence of germ cell neoplasia in situ (intratubular, GCNIS)

# Age structure of testicular tumors patients




# Germ cell neoplasia in situ



Copyright © 2002, Elsevier Science (USA). All rights reserved.



## Germ cell neoplasia in situ - IHC



# Seminoma


- most common
- peak in 4th decade, not in infants
- gross: homogenous, whitish/greyish
- micro: large cells, clear cytoplasm, visible cellular membrane, hyperchromatic nucleus w.visible nucleolus
- stroma with lymphocytic reaction, granulomas possible
- meta into lymph nodes
- successfull therapy, good prognosis usual



# Seminoma




## Seminoma




Copyright © 2002, Elsevier Science (USA). All rights reserved.


# Seminoma



# Seminoma



# Seminoma



# Embryonal carcinoma


- mostly 20-30 yrs
- gross: variable, haemorrhage, necrosis
- micro: organoid glandular, trabecular formations
- large anaplastic cells, confluent




## Embryonal carcinoma




# Embryonal carcinoma



# Embryonal carcinoma



# Embryonal carcinoma




# Choriocarcinoma

- similar to gestational choriocarcinoma
- pure very rare, common admixture
- HCG production (! disperse trophoblastic cells possible in seminoma)
- extensive haemorrhage
- cyto- + syncytiotrophoblast



# Choriocarcinoma




# Yolk sac tumor


- Pure: most common testicular tu in children < 3 yrs (not GCNIS associated)
- Common part of mixed germ cell tumors (GCNIS associated)
- AFP production
- variable histology – microcystic, reticular, papillary formation, variable patterns
  - glomeruloid structures (Schiller-Duval bodies)
    - stalk with capillary lined on the surface by layer of tumor cells



# Yolk sac tumor



# Yolk sac tumor – AFP IHC



# Teratoma


- variable components: ecto-, meso- endoderm (intraembryonal)
- commonly glandular + squamous epithelium
- mesenchymal tissues incl. cartilage
- in males usually immature, component of mixed germinal tu (x females)

# Teratoma


- histologic classification
  - differentiated mature t., usually not GCNIS associated
    - completely matured tissues with organoid structure
    - commonly cystic, containing serous fluid, mucus, keratin
  - differentiated immature t. – part of mixed germ cell tumors derived from GCNIS
    - immature tissues of embryonal/fetal appearance (neuroectoderm, mesenchyme)
  - t. with somatic type malignancy
    - sarcoma, carcinoma, PNET




# Teratoma



# Teratoma + embryonal ca



# Teratoma + choriocarcinoma




# Germ cell tumors

- lymphatic spread common (paraaortic LN)
- hematogenous possible, esp. in non-seminomatous (lungs, bones, liver, brain)
- different histology in metastasis possible
- scarring of the primary tumor possible (burn-out tumor) – diff. dg. x other primary localization – extragonadal germ cell tumors (middle line of the body – retroperitoneum, mediastinum, brain)



# Spermatocytic tumor

- unrelated to germ cell neoplasia in situ
- usual age > 65 yrs
- slow growth, almost no metastasis – surgery only
- more pleiomorphic cells – variable stages of differentiation
- no stromal reaction



Mixture of polymorphic tumor cells (~ early stages of spermatogenesis): large cells with lacy chromatin, middle-sized cells with round nuclei, small lymphocyte-like cells.  
Fibrotic septa without lymphocytic infiltrate


# Sex cord-stromal tumors

- less common than germ cell tu
- Leydig cell tumor
  - any age, peak middle age
  - androgen secreting cells – clinical signs incl. precocious puberty
  - benign or malignant, similar histology
- Sertoli cell tumors
  - very uncommon, mostly benign


# Other testicular tumors

- primary malignant lymphoma
  - older males, in this age ML more common than germ cell tu
  - commonly DLBCL
  - may be already systemic


# Testicular DLBCL



# Testicular DLBCL – Ki-67 proliferation rate



# Testicular DLBCL – CD20+ B-cell type



# Epididymis

- nonspecific epididymitis most common
- usually connected to UTI, bacterial
- purulent, abscess formation, necrosis
- progression to orchitis
- healing by repair, fibrosis + cysts possible
- diff. dg. x tumors


# Sperm granuloma

- pathological situation of sperm in direct contact with stroma
- chronic granulomatous inflammation
- diff. dg.
  - macro x tumor
  - micro x other causes of granuloma incl. TB

# Tumors

- most commonly extension from testicular tumors
- primary tumors rare
  - adenomatoid tumor: benign, phenotype mesothelial, possibly from remnants of Müllerian tract

# Adenomatoid tumor



# Adenomatoid tumor IHC

cytokeratine

