

Clinical examination in orthopaedics

Z. Rozkydal

L. Pazourek

Clinical examination

The aim- establish the diagnosis

1. History
2. Objective examination - general
3. Objective examination - local
4. Laboratory tests
5. Imaging methods

History

Family

Personal

Pharmacological

Social

Occupation

Epidemiological

Current problems

Analysis of pain

Family

- Congenital abnormalities
- Important diseases in family (heart, DM, haemophilia, oncological diseases, neurological diseases, TB)
- Birth, miscarriage

Personal

- Important general diseases (hypertension, DM, heart, tumors, lung problems)
- Coagulopathies
- Infections
- Injuries: consequences, complications
- In children- pregnancy, birth
psychomotor development

Current symptoms

Local

Pain, motor function, limping, deformity, ROM, swelling, loss of sensation

General: fever, shivering, cachexia

Cause of the problem

- injury
- overloading
- Infection
- systemic diseases (endocrine, metabolic, inflammation, neurological, haemotological.)

- Development of symptoms
 - Onset, duration
 - Intensity
 - Aleviation, increasing factors
- Management in the past time
 - Examination in the past time
 - Conservative therapy
 - Operative therapy
- Mobility, occupation
- Emotions, psychological condition
- Simulation, dissimulation, aggravation

Analysis of pain

Intensity, frequency, duration

Acute, chronic

Local, irradiating

Visceral

Type- sharp, blunt, burning, stabbing

Neuralgia

Nerve root pain

Phantom pain

Neurogenic claudication

Analysis of pain

Localised, diffuse

Psychological background

During activity or in rest

VAS – visual analogue scale

Scale of ten degrees

0 - no pain

10 - the most severe pain not bearable

Pain 5 or more- change of management

Pharmacology

- Medicines used currently
- Important medicines: warfarin, heparin, other anticoagulants, antiepileptics, cytostatics, immunosuppresives, NSA, corticoids, biological treatment
- Alcohol, smoking, drugs
- Alergy (antibiotics, metal, disinfection liquids, iodine)

Occupation and social

- Occupation, type of work, manual labor
- Rent
- Social situation (living, marriage)

Management after discharge

Gynecological history

- Cycles, gravidity, menopause, current gynecological problems
- Epidemiological history
influenza, viral infections, herpes simplex,
focal infections (UTI, stomatological infections,
ulcers, erysipel)

Objective examination


General examination

General orthopaedic examination

Local orthopaedic examination

Posture and gait

Somatotype




asthenic

pycnic

normosthenic




Gigantisms




Nanisms



Fröhlich syndrom



Achondroplasia



Marfan syndrom

Nutrition

- **Body mass index:** $\frac{\text{weight kg}}{(\text{BMI}) \text{ height}^2 \text{ m}^2}$
- Below 20 - cachexia
- 20-25 - normal weight
- 25-30 - overweight
- 30-35 - obesity
- Over 35 - severe obesity


Body Mass Index- recent data

Stages

těžká podvýživa	$\leq 16,5$
podváha	16,5–18,5
ideální (zdravá) váha	18,5–25
nadváha	25–30
mírná obezita	30–35
střední obezita	35–40
morbidní obezita	> 40

Skin

- Colour



- pigmentation, naevus



- Trophicity, turgor



- Fistulas, ulcers
- Subcutaneus nodes
- nails



- Lymfadenopathy, soft tumors, inflammations

Swelling

- Local
- General
- Decollement




Local signs of inflammations:
readness, swelling, pain, warm, limited function, soft mass, effusion, discharge

Soft mass



- Haematoma
- Lymphonodes
- Tumor







Effusion





Congenital deformity

- 1. Shape, size
- 2. Differential
- 3. Duplicity
- 4. Gigantisms
- 5. Hypoplasia






Malalignment

- varus x valgus
- antecurvature x recurvature
- rotation deformity



Deformity of spine

- Scoliosis
- Hyperkyphosis, hyperlordosis



Hand deformities



Boutonniere deformity

Swan neck deformity

Foot deformities



Talipes
cavus



Talipes
equinus



Talipes
calcaneus



Talipes
valgus



Talipes
equinovalgus



Talipes
calcaneovalgus



Talipes
varus




Talipes
cavovarus



Talipes
calcaneocavus



Bunion



Claw toe



Length of extremity

Lower extremity

- Spinomaleolar distance
- Umbilicomaleolar distance
- Support during standing
- X-ray of the hip, knee, ankle joint



Upper extremity: acromion- 3. finger

- Circumferential measurement

ROM

- Active and passive movements
- Sagital
- Frontal
- Transversal = horizontal
- Rotation

Shoulder





S: extenze - 0 - flexie
50 - 0 - 180

F: abdukce - 0 -
addukce
180 - 0 - 25

T: abdukce - 0 -
addukce
110 - 0 - 30

R: ZR - 0 - VR
90 - 0 - 90


Elbow



S: extenze - 0 -
flexe
10 - 0 - 150

R: supinace - 0 - pronace
90 - 0 - 90

Wrist




F: rad. dukce - 0 - uln.
dukce

20 - 0 - 40

S:extenze (dorz. flexe) - 0 – flexe (palm. flexe)
80 - 0 - 80

Hip




S: extenze - 0 - flexe
15 - 0 - 140


F: abdukce - 0 -
addukce
60 - 0 - 40

T: abdukce - 0 -
addukce
80 - 0 - 30

P: ZP - 0 - VR




Knee



S: extenze - 0 -
flexe
0 - 0 - 140

Ankle




S: extenze (dorziflexe) - 0 - flexe
(plantiflexe)

20 - 0 - 50



Ancylosis

- Extrarticlar
- Intraarticular



Stability of joints

- Stable joint
- Unstable joint
- Instability
 - acute
 - chronic
 - habitual




Deviace


Desaxace

Shoulder



Apprehension test



Drawer sign




Knee



Varus stress test

Valgus stress test




Laxity



- test

Maneuvers

- Maneuvers



Sound phenomena

- Crepitus

Contracture

- Lumbago, torticollis
- Cerebral palsy




Muscles

- Trophicity
- Tonus
- Cramps
- Power

Muscle test


0 - no activity	0 %
1 - trace	10 %
2 - motion without gravity	25 %
3 - motion against gravity	50 %
4 - motion against gravity and slight resistance	75 %
5 - normal activity	100 %

Posture



Gait

- 1. heel strike
- 2. standing
- 3. toe off
- 4. swing phase



Limping

- Antalgic gait
- Shortening of a lower extremity
- Akylosis
- Trendelenburg sign and gait
- Hemiparetic gait
- Spastic gait
- Drop foot gait
- Parkinson gait

Imaging methods

- X-ray, arthrography
- Angiography
- Ultrasonography
- CT, MRI
- Scintigraphy
- DEXA
- Biopsy

X-ray

In two planes

- bone hypertrophy
- bone atrophy
- osteolysis
- osteonecrosis



Kellgren- Lawrence clasification of O.A.

I.


II.

III.

IV.



Fistulography



Arthrography



Angiography

Clasical

CT angiography

MR angiography



Digital subtraction angiography

Ultrasonography

Echogenicity of tissues

Bone, fibrous tissue, muscles, adipous tissue, cartilage, fluid

Anechoogenic structure- black

Hypoechoenic structure- grey


Hyperechoenic structure- white



Soft tissues


Tumors

DDH

Effusion in joints







CT scann


Absorption of X-ray beams

Air – 1000 H.U.

Water 0 H.U.

Bone + 1000 H.U.





Enhancement with a dye



Bone lesions

Bone tumors

CT



4204-134771
2011/2/16
08:16:01



120.0 kV
251.0 mA
30.0 mAs
Velikost pixelu: 0.313 mm
Pozice: 138.3 mm
W: 300 L: 60



P DFOV: 16.00 x 16.00cm



120.0 kV

520.0 mA

21.0 mAs

Velikost pixelu: 0.811 mm

Pozice: -530.8 mm

W: 1000 L: 200

P

DFOV: 41.50 x 41.50cm

MRI

Magnetic field



Hyposignal- dark


Hypersignal - white

Soft tissue tumors

Soft tissue mass




Spine








2

MRI








Scintigraphy

Tc 99

Bone tumors and metastases



Densitometry DEXA

Absorption of X-ray of two energies (70 and 140 kV)


BMD- bone mineral density in g/cm²

T- score - difference from peak bone mass


Z- score - difference in the same age

Change - difference from previous examination

1 SD = 10% of bone mass



WHO definition of osteoporosis




Laboratory tests

- Inflammation: ESR, leu, CRP, differential, ELFO
- Osteopathy: Ca, P, ALP, bone isoenzyme of ALP
osteokalcin, osteonectin, PTH, vitamin D
- Bone markers- no specific in bones
- PSA in carcinoma of prostate

Joint effusion

- Microscopic
- Biochemic
- Bacteriologic
- PCR
- Immunologic
- Cytologic



Biopsy

Histological examination

Biopsy – open
needle
CT, ultrasonography

