

# Physiology of the Heart

## Conduction System

### Cardiac Cellular Electrophysiology

### Electromechanical Coupling

Assoc. Prof. MUDr. Markéta Bébarová, PhD



*Department of Physiology*  
*Faculty of Medicine*  
*Masaryk University*



**This presentation includes only the most important terms and facts. Its content by itself is not a sufficient source of information required to pass the Physiology exam.**


# ORGANIZATION OF CARDIOVASCULAR SYSTEM

## Roles of the Cardiovascular System

- **primary role** - distribution of dissolved gases and other nutrients
- **several secondary roles, for example:**
  - fast chemical signalling to the cells (circulating hormones)
  - thermoregulation (delivery of heat from the core to the surface of the body)
  - immune reaction
- **roles of the heart:**
  - primary role - pumping of blood
  - endocrine organ (natriuretic peptides)


# ORGANIZATION OF CARDIOVASCULAR SYSTEM

## TWO PUMPS INTERCONNECTED IN SERIES



# ORGANIZATION OF CARDIOVASCULAR SYSTEM

## Two Main Phases of the Cardiac Cycle



ONE WAY VALVES	DIASTOLE	SYSTOLE
ATRIOVENTRICULAR (mitral and tricuspid)	open	closed
SEMILUNAR (aortal and pulmonary)	closed	open



# ORGANIZATION OF CARDIOVASCULAR SYSTEM



## Two Major Types of Cardiac Cells

- **cardiomyocytes of the working myocardium** - specialized for contraction (atrial and ventricular myocytes)

### FUNCTIONAL SYNCYTIIUM

- mechanical connections
- electrical connections - **gap junctions**

sarcomere



# ORGANIZATION OF CARDIOVASCULAR SYSTEM


## Two Major Types of Cardiac Cells

- **cardiomyocytes of the working myocardium** - specialized for contraction (atrial and ventricular myocytes)
- **cardiomyocytes of the cardiac conduction system** - specialized for:
  - automatic excitation (pacemaker activity)
  - conduction of excitation

### The cardiac conduction system ensures:

- 1) generation of automatic electrical activity of the heart (pacemaker activity) that initiates its mechanical activity
- 2) optimal timing of the mechanical activity of the heart as a pump

# CARDIAC CONDUCTION SYSTEM






# CARDIAC CONDUCTION SYSTEM

- **SINOATRIAL (SA) NODE**  
**PRIMARY pacemaker** (60-100 impulses/min)

# CARDIAC CONDUCTION SYSTEM

## SA node

### TWO TYPES of the SA-nodal cells



#### SICK SINUS SYNDROME

- *pacemaker P cells are impaired, activity is slowed or stopped*
- *transmission of excitation from P cells to the atrial cells is reduced or interrupted*


# CARDIAC CONDUCTION SYSTEM

- **SINOATRIAL (SA) NODE**  
**PRIMARY pacemaker** (60-100 impulses/min)
- **INTERNODAL PREFERENTIAL PATHWAYS**
- **ATRIOVENTRICULAR (AV) NODE**  
**SECONDARY pacemaker** (40-55 impulses/min)

# CARDIAC CONDUCTION SYSTEM

## AV node

### THREE TYPES of the AV-nodal cells



# CARDIAC CONDUCTION SYSTEM

## AV node

- **SOLE PATHWAY** FOR PROPAGATION OF EXCITATION FROM ATRIA TO VENTRICLES (NH zone merges into the bundle of His)
- **DELAY** IN PROPAGATION OF EXCITATION, ~100 ms (important for adequate timing of atrial and ventricular contractions)
- **SUBSTITUTIVE (SECONDARY) PACEMAKER** (40-55 impulses/min; importance in the case of sick sinus syndrome)
- **FILTER OF SUPRAVENTRICULAR TACHYARRHYTHMIAS**  
atrial excitations are transmitted to the ventricles only up to the limited frequency 180-200 excitations/min (the heart function as a pump is preserved)



# CARDIAC CONDUCTION SYSTEM


- **SINOATRIAL (SA) NODE**  
**PRIMARY pacemaker** (60-100 impulses/min) **0.05 m/s**
- **INTERNODAL PREFERENTIAL PATHWAYS** **1 m/s**
- **ATRIOVENTRICULAR (AV) NODE**  
**SECONDARY pacemaker** (40-55 impulses/min) **0.05 m/s**
- **BUNDLE OF HIS** **1 m/s**
- **BUNDLE BRANCHES (LEFT AND RIGHT)** **1 m/s**
- **PURKINJE FIBRES**  
**TERCIARY pacemaker** (25-40 impulses/min) **4 m/s**

**Conduction velocity in atrial and ventricular muscle: 1 m/s**

# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Ionic Channels



Movement of ions through the open channels  
**down their electrochemical (concentration + electrical) gradients**



$$V_m = \phi_i - \phi_e$$

# CARDIAC CELLULAR ELECTROPHYSIOLOGY


## Ionic Currents Underlying Action Potential Configuration



Despopoulos, Color Atlas of Physiology © 2003



# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Ionic Currents Underlying Action Potential Configuration




# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Ionic Currents Underlying Action Potential Configuration




many subtypes of potassium channels (currents)






# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Ionic Currents Underlying Action Potential Configuration



# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Mechanism of the initial fast depolarization (phase 0)



**regenerative (self restoring) process**

produced by POSITIVE FEEDBACK  
between MEMBRANE VOLTAGE and  
CONDUCTANCE of MEMBRANE  
CHANNELS ( $g_{Na}$ ,  $g_{Ca}$ )

**working myocardium -  $I_{Na}$**

*depolarizing currents*


**↑ depolarization** ⇒ **↑ conductance of  $Na^+$  ( $Ca^{2+}$ ) channels** ⇒ **↑  $I_{Na}$  ( $I_{Ca}$ )**

*(directly proportionate to the fraction of  
 $Na^+$  ( $Ca^{2+}$ ) channels in the open state)*

# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Refractory Period – Suppression of Excitability

action potential



**protection** of the heart against:



- retrograde propagation of excitation (reentry)
- tetanic contraction at higher heart rate

# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Refractory Period – Suppression of Excitability

action potential

contraction



**ABSOLUTE**

**RELATIVE**

**CLINICAL ASPECTS**

**EFFECTIVE REFRACTORY PERIOD**

*(ARP + period of responses that do not propagate)*

**VULNERABLE PERIOD**  
*(propagated responses)*



increased susceptibility to ventricular fibrillation !!!

ECG

T

# CARDIAC CELLULAR ELECTROPHYSIOLOGY



## Refractory Period - Mechanism



**INACTIVATED STATE  
(closed)**


**INACTIVATION  
(at maintained  
membrane  
depolarization)**

**RECOVERY FROM  
INACTIVATION  
(at repolarization)**



**RESTING STATE  
(closed)**

**ACTIVATION  
(depolarization over  
threshold voltage)**





**OPEN STATE**




# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Pacemaker Activity - Mechanism



# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Pacemaker Activity - Mechanism




### FACTORS DETERMINING THE HEART RATE:

- 1) maximal diastolic voltage
- 2) steepness of diastolic depolarization
- 3) threshold voltage for activation of  $I_{Ca-L}$

# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Pacemaker Activity - Mechanism




**COMPLEX PROCESS** resulting from an INTERPLAY between

- **REPOLARIZING CURRENTS**, namely  $I_K$  (including  $I_{K,Ach}$ )
- **DEPOLARIZING CURRENTS**, namely  $I_f$ ,  $I_{Ca-T}$ , and  $I_{NaCa}$

# CARDIAC CELLULAR ELECTROPHYSIOLOGY

## Pacemaker Activity - Mechanism




### SYMPATHETIC STIMULATION

- $\uparrow$  cAMP  $\longrightarrow$   $\uparrow$   $I_f$  and  $I_{Ca-T}$   $\longrightarrow$   $\uparrow$  rate of diastolic depolarization  
 $\longrightarrow$   $\downarrow$  threshold voltage for activation of  $I_{Ca-L}$   
( $\uparrow$  excitability)

# CARDIAC CELLULAR ELECTROPHYSIOLOGY


## Pacemaker Activity - Mechanism




### PARASYMPATHETIC STIMULATION

- $\downarrow$  cAMP  $\longrightarrow$   $\downarrow I_f$  and  $I_{Ca-T}$   $\longrightarrow$   $\downarrow$  rate of diastolic depolarization  
 $\longrightarrow$   $\uparrow$  threshold voltage for activation of  $I_{Ca-L}$   
( $\downarrow$  excitability)
- activation of  $I_{K,Ach}$   $\longrightarrow$   $\downarrow$  maximal diastolic voltage

# SPREADING OF EXCITATION IN THE HEART



electric field  
(magnitude, direction)




# ELECTROMECHANICAL COUPLING

## Excitation-Contraction Coupling

### SARCOTUBULAR SYSTEM


SIMILAR ARRANGEMENT  
IN SKELETAL AND CARDIAC  
MUSCLE CELLS





# ELECTROMECHANICAL COUPLING

## Excitation-Contraction Coupling in Cardiomyocytes



CALSEQUESTRIN  
 $Ca^{2+}$  buffer in SR

$[Ca^{2+}]_i$

200 ms



voltage-dependent  $Ca^{2+}$  channels in the cell membrane  
(both the surface membrane and membrane of t-tubules)



$Ca^{2+}$ -RELEASE channels in SR ( $Ca^{2+}$ -sensitive)




# ELECTROMECHANICAL COUPLING

## Molecular Mechanism of Contraction

### FORMATION OF CROSS BRIDGES BETWEEN ACTIN AND MYOSIN FILAMENTS

MECHANISM IDENTICAL  
IN SKELETAL AND CARDIAC  
MUSCLE CELLS





# ELECTROMECHANICAL COUPLING


## Molecular Mechanism of Contraction

### TROPONIN-TROPOMYOSIN COMPLEX

#### RESTING MUSCLE




#### CONTRACTING MUSCLE



© 2011 Pearson Education, Inc.

# ELECTROMECHANICAL COUPLING

## Molecular Mechanism of Contraction



# ELECTROMECHANICAL COUPLING

## Molecular Mechanism of Contraction

