

THE CRANIAL NERVES(*nervi craniales*)

- I. **Nervus olfactorius – the olfactory nerve**
- II. **Nervus opticus – the optic nerve**
- III. **Nervus oculomotorius – the oculomotor nerve**
- IV. **Nervus trochlearis – the trochlear nerve**
- V. **Nervus trigeminus – the trigeminal nerve**
- VI. **Nervus abducens – the abducens nerve**
- VII. **Nervus facialis – the facial nerve**
- VIII. **Nervus vestibulocochlearis – the vestibulocochlear nerve**
- IX. **Nervus glossopharyngeus – the glossopharyngeal nerve**
- X. **Nervus vagus – the vagus nerve**
- XI. **Nervus accessorius – the accessory nerve**
- XII. **Nervus hypoglossus – the hypoglossal nerve**



THE CRANIAL NERVES (*nervi craniales*)

— sensory fibres
— motor fibres

Optic (II)
sensory: eye



Trochlear (IV)
motor: superior oblique muscle




Olfactory (I)
sensory: nose



Abducent (VI)
motor: external rectus muscle



Oculomotor (III)
motor: all eye muscles except those supplied by IV and VI




Trigeminal (V)
sensory: face, sinuses, teeth, etc.

motor: muscles of mastication

© 2007 Encyclopædia Britannica, Inc.

Facial (VII)
motor: muscles of the face



Hypoglossal (XII)
motor: muscles of the tongue



Intermediate motor:

submaxillary and sublingual gland

sensory:

anterior part of tongue and soft palate

Vestibulocochlear (VIII)
sensory: inner ear




vestibular
cochlear

Glossopharyngeal (IX)
motor:

pharyngeal musculature

sensory:

posterior part of tongue, tonsil, pharynx



Vagus (X)
motor:

heart, lungs, bronchi, gastrointestinal tract

sensory:
heart, lungs, bronchi, trachea, larynx, pharynx, gastrointestinal tract, external ear


Accessory (XI)

motor: sternocleidomastoid and trapezius muscles

The outputs from the brain

- I. – the cerebrum
- II. – the diencephalon
- III.-XII. – the brainstem

IV. – dorsally !!!



Cranial Nerves

Olfactory (I)

Olfactory bulb

Olfactory tract

Optic tract

Vestibulo-cochlear (VIII)

Hypo-glossal (XII)

Vagus (X)

Optic (II)

Oculomotor (III)

Trochlear (IV)


Trigeminal (V)



Abducens (VI)

Facial (VII)


Glossopharyngeal (IX)

Accessory (XI)







The outputs from the skull



I. = The olfactory nerve

- special **sensory** nerve: smell, part of the olfactory tract
- cavitas nasi → lamina cribrosa → cavitas crani anterior
- Superior area of the nasal cavity in range of concha nasalis superior, on the wall, roof and septum




The function of the nerve

sensory



- it carries information from the olfactory receptor into the CNS






II. = The optic nerve


- It hasn't the nuclei— centre is in the occipital lobe
- special **sensory** nerve: vision, part of visual tracts
- orbita → canalis opticus → cavitas cranii media






The function of the nerve **sensory**

- it carries information from visual receptor into the CNS





Optic nerve






The extraocular muscles

- mm. recti (bulbi)
 - superior, inferior, medialis, lateralis
- mm. obliqui (bulbi)
 - *inferior, superior*
- m. levator palpebrae superior
- ***innervation:*** n. III., IV., VI.




Movements of the eyeball





The extraocular muscles innervation:


- Through sinus cavernosus and fissura orbitalis superior, they enter the orbit
- **III. The oculomotor nerve**
 - fossa interpeduncularis (sulcus nervi oculomotorii)
 - motor** – 5 extraocular muscles
 - + parasympathetic fibers for innervation of smooth muscles of the iris - *m. sphincter pupillae* – pupil constriction (miosis) and of corpus ciliare- *m. ciliaris* – accommodation of the lens
- **IV. The trochlear nerve**
 - Posterior surface of the brainstem
 - purely motor** → 1 muscle = ***m. obliquus superior***
- **VI. The abducens nerve**
 - sulcus bulbopontinus
 - purely motor** → 1 muscle - ***m. rectus lateralis***






n.III.










© Patrick J. Lynch, 2006






n.IV.






n.VI.





Nerves passing through superior orbital fissure





Palsy of n. III

- divergent strabismus
- Pupil dilatation- mydriasis
- Accommodation disorder
- falling of the upper eyelid - ptosis
- double vision- diplopia



Palsy of n. IV and n.VI

- convergent strabismus)



V. = The trigeminal nerve

- *ganglion trigeminale* (located within *cavum trigeminale*) - sensory

V1 = n. ophthalmicus = ophthalmic nerve (*sensory*)


V2 = n. maxillaris = maxillary nerve (*sensory*)



V3 = n. mandibularis = mandibular nerve (*mixed*)

Radix motoria = somatomotor branch

for the masticatory muscles and other

4 muscles- fibers only within V3 !!!





n.V.










Illustration : A. Micheau, MD







© 2001 CCND, Winnipeg



N. V


- V1 = N. ophthalmicus**
- V2 = N. maxillaris**
- V3 = N. mandibularis**




V1 = N. ophthalmicus


- fissura orbitalis superior into orbit – to the frontal region
- Skin of the upper part of face (forehead, upper eyelids, dorsum nasi), content of the orbit, part of mucosa of nasal cavity and paranasal sinuses)



- n. frontalis
 - n. lacrimalis
 - n. nasociliaris
- n. supraorbitalis – *sensitivity to palpation*



- n. frontalis: r.supraorbitalis (r. medialis et lateralis), r. supratrochlearis (skin of forehead till interauricular line)
- n. lacrimalis (lacrimal gland, upper eyelid)
- n. nasociliaris-branches: nn. ciliares longi, n. ethmoidalis posterior, n. ethmoidalis anterior, n. infratrochlearis (sinus sphenoidalis, nasal cavity, sensory fibres into the eye)









V2 = N. maxillaris

- foramen rotundum - branches out behind the maxilla- fossa pterygopalatina
- skin of middle part of face (lower eyelid, upper lip, nasal wings and upper half of cheek), **teeth of maxilla**, part of mucosa of nasal and oral cavity




- **n. zygomaticus**: n. zygomaticofacialis, n. zygomaticotemporalis, r. communicans cum nervo zygomatico (skin of cheek)
- **n. infraorbitalis** (teeth of maxilla, sinus maxilaris, mucosa of nasal cavity, palate, nasopharynx)



V3 = N. mandibularis

- foramen ovale – branches out behind ramus mandibulae-fossa infratemporalis
 - skin of inferior part of face (lower lip, inferior half of cheek, chin), teeth of mandibula, part of mucosa of oral cavity
 - masticatory muscles, muscles of neck


FOSSA INFRATEMPORALIS - l.dx.



n. mandibularis

somatotor branches:


- Masticatory muscles
- suprathyoid muscles
- m. tensor veli palatini
- m. tensor tympani



somatosensory branches:

- n. alveolaris inferior (teeth)
- n. lingualis (mucosa of bottom of oral cavity, gingiva, front 2/3 of tongue)
 - chorda tympani from n. VII
- n. buccalis (skin and mucosa of cheek)
- n. auriculotemporalis (auricle and temporal region)
- **parasympathetic ganglion submandibulare + ganglion oticum (small salivary glands of tongue and bottom of oral cavity)**
- n. mentalis – sensitivity to palpation





- Branches for masticatory muscles
- n. buccalis
- n. lingualis (n. sublingualis, rr. linguales)
- n. alveolaris inferior (n. mylohyoideus)
- n. auriculotemporalis (r. meningeus, r. communicans cum ganglio otico)











Illustration : A. Micheau, MD










VII. = The facial nerve

- **mixed nerve**
- **2 basis**
 - n. facialis – motor (facial muscles)
 - n. intermedius - visceromotor + taste + sensory





n. VII.




inside canalis nervi facialis

- n. petrosus major –
parasympathetic into ggl.
pterygopalatinum
- n. stapedius – m. stapedius
- chorda tympani
 - **parasympathetic** into glandula
submandibularis
 - taste → into front 2/3 of
tongue



outside the skull

- n. auricularis posterior –**motor
nerve** to rudimentary muscles of
auricle
- nn. into m. stylohyoideus +
venter post. m. digastrici
- plexus intraparotideus (facial
muscles: rr. temporales, rr.
zygomatici, rr. bucales, r.
marginalis mandibulae, r. colli)



PRŮBĚH CANALIS NERVI FACIALIS V PYRAMIDĚ

Pohled shora





somatosensory – auricle and external auditory canal


sensory – from gustatory cells of front 2/3 of tongue

somatomotor – facial muscles, musculus platysma, musculus stapedius, m.stylohyoideus, m.digastricus- posterior belly

visceromotor – parasympathetic fibres for secretion of lacrimal gland, submandibular and sublingual gland









VIII. – The vestibulocochlear nerve


- *special sensory nerve* – hearing + balance






Common output of



- n.IX., X., XI.



IX. The glossopharyngeal nerve

- Arises from the medulla oblongata
- Passes through foramen jugulare
- Branches out under the base of skull
- Branches: n. tympanicus → n. petrosus minor, rr. pharyngei, tonsillares, linguales






n. IX.






- **Motor in.:** soft palate muscles (except tensor), m. stylopharyngeus, muscles of pharynx
- **Parasympathetic in.:** mucosa of tympanic cavity and glandula parotide (through ganglion oticum)
- **Sensory in.:** pharynx, back third of tongue, tonsil
- **Gustatory in.:** back third of tongue



X. The vagus nerve

- arises from medulla oblongata
- Passes through foramen jugulare
- Descends caudally and passes together with arteria carotis interna (lower arteria carotis communis) and vena jugularis interna, it attaches to the esophagus and along it descends through thoracic cavity till abdominal cavity (in male till scrotum)





n. X.




r. auricularis (sensory branch for wall of ear canal)

rr. pharyngei (muscles of pharynx)

n. laryngeus superior

n. laryngeus recurrens- n.laryngeus inferior

(muscles of larynx)



Organs within thoracic and abdominal cavity:

rr. cardiaci - plexus cardiacus

rr. bronchiales- plexus pulmonalis

rr. oesophagei- plexus oesophagus


rr. gastrici



rr. hepatici



rr. coeliaci - plexus coeliacus


rr. renales

- **sensory in.**: part of skin of the auricle and ear canal
- **viscerosensory in.**: organs of thoracic and most of abdominal cavity
- **somatomotor in.**: striated muscles of pharynx and larynx
- **visceromotor in.**: smooth muscle of respiratory tract and most of alimentary tract, smooth muscle of appropriate vessels and myocardium












XI. The accessory nerve

- arises from the medulla oblongata
- passes through foramen jugulare
- name is abbreviation of the original name „nervus accessorius ad vagum“ (because part of its fibers joins with nervus vagus and below it separates again and innervates muscles of larynx)
- Nerve branches out after leaving the cranial cavity





n. XI.





n.XI.


Radices craniales *Radices spinale*s

Ramus internus- joints n.X.
(muscles of larynx)

Ramus externus


motor – part of **musculus sternocleidomastoideus** and **musculus trapezius**







XII. The hypoglossal nerve

- Arises from the medulla oblongata - sulcus anterolateralis
- Through canalis nervi hypoglossi
- Directs to the tongue, where it branches out







n. XII.




N. hypoglossus

Motor nerve: innervation of striated muscle of the tongue
ansa cervicalis profunda- **infrahyoid muscles (plexus cervicalis)**







Obrázky:

- **Atlas der Anatomie des Menschen/Sobotta.** Putz,R., und Pabst,R. 20. Auflage. München:Urban & Schwarzenberg, 1993
- **Netter: Interactive Atlas of Human Anatomy.**
- **Naňka, Elišková: Přehled anatomie.** Galén, Praha 2009.
- **Čihák: Anatomie I, II, III.**
- **Drake et al: Gray's Anatomy for Students.** 2010

