

Energetic metabolism


Physiology II lecture (aVLFY0422p)

Tibor Stračina

The presentation is copyrighted work created by employees of Masaryk University. Any unauthorized reproduction or distribution of the presentation or individual slides is against the law.

Energetic metabolism


- Energy input (external and internal sources)
 - Energy output
 - Energy stored
- **INPUT = OUTPUT + STORAGE**



Energy input

- Basic substrates: **carbohydrates, fats a proteins**
- Energy is obtained by burning (oxidizing) substrates
 - carbohydrates 4,1 kcal/g
 - fats 9,3 kcal/g
 - proteins 5,3 kcal/g (in the body 4,1 kcal/g)
- Source of substrates: **food intake or mobilization of reserves**

Nutrient burning



Energy output

- **Basal metabolism** – energy expenditure to maintain homeostasis under basal conditions (vital function) – *~75% of AEE in a person sitting at rest*
- **Specific dynamic effect of food** – a small increase in energy expenditure after eating – *~7% of AEE in a person sitting at rest*
- **Thermoregulation**
- **Spontaneous motoric activity** – *~18% of AEE in a person sitting at rest*
- **Physical work (exercise)**

Energy storage and transfers

- Irregular energy intake and output – the need for energy storage
- Ready-to-use stock - macroergic compounds
 - ATP
 - creatin phosphate
 - GTP, CTP, UTP, ITP
- Long-term storage – stock substrates
 - Fat, proteins, glycogen

Adenosine trisphosphate (ATP)

- universal macroergic compound

Synthesis

- circa 63 kg/day (128 mol/day)
- oxidative phosphorylation
- glykolyysis – for short-term production only, production of lactate

Use


- macroergic bond splitting – efficiency is not 100%, **heat release**

Storage substrates

- Triacylglycerols in fat tissue (75% of stores) – up to 2 months
 - Source: FA from food and esterification with α -glycerol phosphate or synthesis of FA from acetyl-CoA from glycolysis (conversion of sugars into a more efficient energy store = fat)
- Proteins in skeletal muscles and blood plasma (25% of stores)
 - Possible conversion to sugars (glukoneogenesis; stimulated by glucocorticoids)
 - Blood plasma proteins – quickly usable; leads to hypoproteinemia, drop of specific immunity
 - Mobilization of muscle proteins leads to sarcopenia
- Carbohydrates in form of glycogen (less than 1% of stores)
 - Important for the CNS and covering energy demands during short-term physical work
 - Glycogen stored in the liver (about 25%) and in the muscles (about 75%)
 - Liver glycogen - glycogenolysis - release of Glc into the blood
 - Muscle glycogen - use only in muscles (glucose-6-phosphatase is missing)

Energy transfers between organs

- Only in the form of substrates (glucose, FA, AA, lactate, ketons, ...)
- Any transfer of substrates consumes some energy (synthesis and splitting of stock substrates, transports, ...)



Measurement of energy expenditure

- Direct calorimetry

- Indirect calorimetry (*PRACTICE!!!*)

- Consumption of O₂ – **energetic equivalent of oxygen** (amount of energy released when consuming 1 liter of O₂)

- carbohydrate: 21,15 kJ/l

- fat: 19,6 kJ/l

- protein: 19,65 kJ/l

- mixed diet: 20,1 kJ/l

- Consumption of O₂ + production of CO₂ – **respiratory quotient** ($RQ = V_{CO_2} / V_{O_2}$)

- carbohydrate: RQ = 1

- fat: RQ = 0,7

- protein: RQ = 0,8 – 0,9

Physiology of Exercise

Physiology II lecture (aVLFY0422p)

Tibor Stračina


Work (physical activity, exercise)



Skeletal muscle

- Contraction: isometric (static work) vs. isotonic (dynamic work)
- Blood flow depends on muscle tension
- Metabolic autoregulation: $\downarrow pO_2$; $\uparrow pCO_2$; $\downarrow pH$; $\uparrow K^+$; \uparrow local temperature
- Metabolism: aerobic vs. anaerobic
- Muscle spindles – muscle tension – afferentation of exercise pressor reflex

Skeletal muscle metabolism



Reaction of the body to exercise

- Sympathetic NS (ergotropic system)
- Cardiovascular changes
- Respiratory changes
- Metabolic changes

- HOMEOSTASIS

Anticipation of exercise

- Reaction of the body (cardiovascular system)
- Prepare the body for the increased metabolism of the exercising skeletal muscles

- Same as the early response to exercise
- Resembling fight-or-flight reaction

Cardiovascular response to exercise


- Increased cardiac output
- Vasoconstriction in inactive skeletal muscles, the GIT, skin, (kidneys)
- Vasodilation in active muscles (metabolic autoregulation)
- Increased venous return
- Histamine release
- Epinephrine release (adrenal medulla)
- Thermoregulation

Increase of cardiac output. Cardiac reserve

- $CO = SV \times HR$ (SNS: positive inotropic and chronotropic effect)
- **Cardiac reserve = maximal CO / resting CO** (4 – 7)
- Coronary reserve = maximal CF / resting CF (~3.5)
- Chronotropic reserve = maximal HR / resting HR (3 – 5)
- Volume reserve = maximal SV / resting SV (~1.5)

CO – cardiac output; CF – coronary flow; HR – heart rate; SV – stroke volume

Cardiac reserve in healthy and failing heart



Changes of arterial blood pressure

PARAMETER	AT REST	DURING EXERCISE	INCREASE (x)
Cardiac output [L/min]	5 – 6	25 (35)	4 – 5 (7) <i>cardiac reserve</i>
Heart rate [1/min]	(45) 60-90	190 – 200 (220) <i>age-dependent</i>	3 – 5 <i>chronotropic reserve</i>
Stroke volume [mL]	75	115	~1.5 <i>volume reserve</i>
Systolic BP [mmHg]	120	<i>static work</i> ↑ <i>dynamic work</i> ↑↑	
Diastolic BP [mmHg]	70	<i>static work</i> ↑↑↑ <i>dynamic work</i> – / ↓	
Mean arterial P (MAP) [mmHg]	~90	<i>static work</i> ↑ <i>dynamic work</i> – / ↑	
Muscle persufion [mL/min/100g]	2 – 4	60 – 120 (180) <i>static vs. dynamic work</i>	30 (10% CO _{max})

Respiratory response to exercise

- Respiratory centre - \uparrow ventilation
 - chemoreceptors: \uparrow pCO₂ + \downarrow pH
 - proprioceptors in lungs
- Sympathetic stimulation (stress – anticipation)

Respiratory response to exercise

PARAMETER	AT REST	DURING EXERCISE	INCREASE (x)
Ventilation [L/min]	6 – 12	90 – 120	15 – 20 <i>respiratory reserve</i>
Breathing frequency [1/min]	12 – 16	40 – 60	4 – 5
Tidal volume (V_T) [mL]	0.5 – 0.75	~2	3 – 4
Pulmonary artery blood flow [mL/min]	5 – 6	25 – 35	4 – 6
O₂ uptake (V_{O_2}) [mL/min]	250 – 300	~3000	10 – 12 (25)
CO₂ production [mL/min]	~200	~8000	~40

Oxygen uptake by lungs

– Spiroergometry

– Resting \dot{V}_{O_2} : **~3.6** mL O₂ / (min x kg)


– $\dot{V}_{O_2 \max}$ – objective index for aerobic power

– untrained middle age person: **30 – 40** mL O₂ / (min x kg)

– elite endurance athletes: **80 – 90** mL O₂ / (min x kg)

– HF / COPD patients: **10 – 20** mL O₂ / (min x kg)

Adopted from:
<https://studentconsult.inkling.com/read/boron-medical-physiology-3e/chapter-60/figure-60-6>




Determinants of $V_{O_2 \max}$

1. Uptake of O_2 by the lungs
 - pulmonary ventilation
2. O_2 delivery to the muscles
 - blood flow (pressure gradient – cardiac output x resistance)
 - haemoglobin concentration
3. Extraction of O_2 from blood by muscle
 - pO_2 gradient: blood-mitochondria

Oxygen consumption during exercise


Adopted from: D.U.Silverthorn:
Human Physiology (An Integrated
Approach)

– Oxygen debt



Energy substrate used by skeletal muscle during exercise


- Low-intensity e.: fats
- High-intensity e.: glucose



Data from G. A. Brooks and J. Mercier, *J App Physiol* 76: 2253–2261, 1994

Adopted from: D.U.Silverthorn:
Human Physiology (An Integrated
Approach)

Energy substrate use – aerobic vs. anaerobic



Adopted from: D.U.Silverthorn:
Human Physiology (An Integrated
Approach)

Testing of fitness

- Spiroergometry
- Standardised workload
 - accurate: in W/kg
 - comparative (simple, inaccurate): in MET
 - metabolic equivalent (actual MR / resting MR)
 - 1 MET = uptake of 3.5 ml O₂/kg.min ≈ 4.31 kJ/kg.h
 - sleeping ≈ 0.9 MET; slow walking ≈ 3-4 MET; fast running ≈ 16 MET

Indexes of fitness

- W_{170} [W/kg]
- $V_{O_2 \max}$ [mL O_2 / (min x kg)]
- Aerobic / anaerobic threshold

- Fatigue
- Training
- Adaptation to exercise

The presentation is copyrighted work created by employees of Masaryk University. Any unauthorised reproduction or distribution of the presentation or individual slides is against the law.