

## Revision of the 3<sup>rd</sup> Declension

### 1) Fill in the table:

Give English translations:		Give Latin translations (+ genitive, gender etc.)	
Margo, inis, m.		Tooth	
Iecur, coris, n.		Part	
Os, ossis, n.		Fever	
Encephalitis, itidis, f.		Cyst	
Radix, dicis, f.		Ear	
Cutis, is, f.		Thumb	
Deciduus, ua, uum		Aperture, opening	
Symptoma, atis, n.		Wound	
Mens, ntis, f.		Acquired	
Frons, ntis, f.		human	

### 2) Fill in missing endings:

Luxatio cox... dextr... cum fractur... aperta pelv... later.. sinistr...  
 Tuberculosis in lob... medi... pulmon... dextr...  
 Carcinoma corpor... uteri et ovari...(pl.)  
 Haemorrhagia postoperativ... in cavitat... abdomin...  
 Ruptura ret... venos... bulb... ocul...  
 Mors in tabul... propter vuln... sclopetari...

### 3) Decline in both singular and plural:

sanatio longa      cutis laesa      rete venosum      sepsis lenta      os fractum

### 4) Translate into Latin:

Fever before operation of gall bladder  
 Extraction<sup>1</sup> of milky teeth without narcosis  
 Full spoons of honey for a sick person against cough  
 Congenital defect of the valves of heart  
 Transversal<sup>2</sup> diameter of the skull

---

<sup>1</sup> extractio

<sup>2</sup> transversus