

Schémata a animace zpracovalo

Servisní středisko pro e-learning na MU

CZ.1.07/2.2.00/28.0041

Centrum interaktivních a multimediálních studijních opor pro inovaci výuky a efektivní učení




INVESTICE DO ROZVOJE VZDĚLÁVÁNÍ


MEMBRANE OF EXCITABLE CELL

ELECTRICAL TRANSMISSION OF INFORMATION




PLASMATIC MEMBRANE




Nexus (gap junction) →



SODIUM-POTASSIUM PUMP



RESTING MEMBRANE VOLTAGE



Nernst equation:

$$E_x = \frac{R \cdot T}{F} \ln \frac{(C_{x_{out}})}{(C_{x_{in}})}$$

$$I_x = g_x \cdot (E - E_x)$$

Equilibrium potential

$$E_{Na} = +40 \text{ mV}$$

$$E_K = -90 \text{ mV}$$


$$E_{Cl} = -70 \text{ mV}$$


$$E_{Ca} = +60 \text{ mV}$$

$$E_r = -85 \text{ mV}$$


I – current, E – voltage, g – specific voltage and time-dependent conductance

ACTION POTENTIAL






ACTION POTENTIAL









- Unit of excitation activity
- „All or nothing“ response
- Propagation without decrement („domino effect“)
- **Refractery**

Local current



Propagation with decrement





Velocity of excitation propagation (depolarisation front) is a function of:

- Intensity of local currents
- Resistance outside the conductor (myelin)
- Resistance of the conductor (indirect relationship)

Nodes of Ranvier, saltatory conduction

Neuron

input section
(coding of inf.)

transmission section
(transmission of inf.)

output section
(decoding of inf.)



SYNAPSES

- excitatory
- inhibitory

Action potential
Calcium ions


Synaptic vesicles (exocytose)





Neurotransmitter (mediator)

Presynaptic membrane

Synaptic cleft

Postsynaptic membrane
(local change of voltage)






SYNAPSIS:

A – axodendritic

B – axosomatic

C - axoaxonal

Sarcomere



Z line

Z line

Thin filaments

actin

Thick filaments


myosin



H zone

I band

A band

I band








Structure and function

	Smooth muscle	Cardiac muscle (striated)	Skeletal muscle (striated)
Motor end-plates	None	None	Yes
Fibers	Fusiform, short (≤ 0.2 mm)	Branched	Cylindrical, long (≤ 15 cm)
Mitochondria	Few	Many	Few (depending on muscle type)
Nucleus per fiber	1	1	Multiple
Sarcomeres	None	Yes, length ≤ 2.6 μm	Yes, length ≤ 3.65 μm
Electr. coupling	Some (single-unit type)	Yes (functional syncytium)	No
Sarcoplasmic reticulum	Little developed	Moderately developed	Highly developed
Ca ²⁺ "switch"	Calmodulin/caldesmon	Troponin	Troponin
Pacemaker	Some spontaneous rhythmic activity (1s^{-1} – 1h^{-1})	Yes (sinus nodes ca. 1s^{-1})	No (requires nerve stimulus)
Response to stimulus	Change in tone or rhythm frequency	All or none	Graded
Tetanizable	Yes	No	Yes
Work range	Length-force curve is variable	In rising length-force curve (see 2.15E)	At peak of length-force curve (see 2.15E)

Response to stimulus

Potential —
Muscle tension —



2 Nerve and Muscle, Physical Work

- **RESTING MEMBRANE POTENTIAL IS A CONDITION OF EXCITABILITY**
- **IT DEPENDS ON HIGH RESTING MEMBRANE CONDUCTIVITY FOR POTASSIUM**

ACTION POTENTIAL IS A PROPAGATED ELECTRICAL SIGNAL GENERATED BY FAST SODIUM CURRENT INTO THE CELL_x

- **ACTION POTENTIAL REPRESENTS UNIT OF INFORMATION**
- **CODING OF INFORMATION IN THIS SYSTEM IS PERFORMED BY CHANGED FREQUENCY OF ACTION POTENTIALS**