


Digestive system

1. **Microscopic anatomy of esophagus, stomach, small and large intestine**
2. Microscopic anatomy of liver, pancreas







Petr Vaňhara, PhD

Department of Histology and Embryology LF MU
PVanhara@med.muni.cz


General architecture of hollow organs incl. gut tube

Four layers







General architecture of hollow organs incl. gut tube



Mucosa (Tunica mucosa)

- inner layer of gut tube
- protective, absorption and resorption
- microscopic structure depending on localization



- **Lamina epithelialis** mucosae
- **Lamina propria** mucosae
- **Lamina muscularis** mucosae



Mucosa (Tunica mucosa)

- **Lamina epithelialis** mucosae

- epithelium type corresponding to function of gut tube
- oral cavity, pharynx, esophagus, anus – **stratified squamous ep.**
- stomach, intestine – **simple columnar**
- **mucus** - secreted by mucosal or submucosal glands (oral cavity, esophagus), secretory epithelium (stomach) or goblet cells (intestine)




- **Lamina propria** mucosae

- Layer of mucosal connective tissue – loose collagen
- Fenestrated blood capillaries – transport of metabolite (intestine)
- mucosal glands in some regions /esophagus)
- innervations, immune system



- **Lamina muscularis** mucosae



- smooth muscles in two layers (inner circular, outer longitudinal)
- small mechanical movements of mucosa facilitating secretion and absorption independently on peristaltic movements.



Submucosa (Tela submucosa)


Submucosal connective tissue

- distinct layer of loose connective tissue
- defines shape of mucosa (rugae, plicae)
- larger blood and lymph veins nourishing mucosa, muscularis externa and serosa
- **innervations** – nerve plexus - **plexus submucosus Meissneri**
 - = groups of multipolar neurons and small ganglions, visceral sensory fibers (sympaticus) and fibers and terminal ganglions of parasympaticus (enteric nerve system)
- glands – different in different regions
 - protective function




Outer muscular layers (Tunica muscularis externa)


- Two concentric, thick layers of smooth muscle, separated by thin layer of connective tissue
- Inner – **circular**, outer – **longitudinal** (spiral)
- Myenteric (Auerbach) plexus
- Peristaltic – passage through the gut tube
- **Local modifications of m.e.**
 - pharyngoesophageal sphincter + external anal sphincter – skeletal muscles
 - stomach – third - oblique - layer
 - taenie coli – thickened part of longitudinal layer in colon



Circular



Longitudinal



Serosa/Adventitia (Tunica serosa/adventitia)


- outermost layer of gut tube

- **Serosa**

- serous membrane of loose connective tissue (Lamina propria serosae) and single layer squamous epithelium (L. epithelialis serosae)
- syn. mesothelium, visceral peritoneum
- continuous with mesenterium
- barrier against various pathogens , antiadhesive properties – intracoelomic movements, immune functions (Ag presentation), ECM production, etc.

- **Adventitia**

- some parts of the tube are not covered with epithelium
- esophagus in thorax, parts of digestive system in peritoneal cavity in walls (duodenum, part of colon, rectum, anal canal)
- connective tissue only continuous with connective tissue of the walls




Esophagus (Oesophagus)

- Mucosa

- nonkeratinized stratified squamous epithelium → mechanically protects esophageal tissue
- L. propria contains cardial glands (tubular mucinous) and diffuse lymphatic tissue

- Submucosa

- loose collagen connective tissue, defines shape of mucosa
- blood and lymph veins, plexus submucosus Meissneri
- submucosal glands (tubular mucinous)
- diffuse lymphatic tissue




Esophagus (Oesophagus)

- Muscularis externa

- inner circular and outer longitudinal layer
- plexus myentericus Auerbachi
- upper third – skeletal muscle, mid third – mixed smooth and skeletal, lower third – smooth muscles only

- Adventitia

- neck and chest – connects esophagus with surrounding tissue
- loose connective tissue
- in peritoneal cavity - serosa





Cardia of stomach – connection with esophagus

Nonkeratinized stratified squamous epithelium → simple columnar epithelium



Stomach (Ventriculus, Gaster)




- general anatomy of hollow tube
- anatomical regions differ also in histologic structure
- rugae gastricae (submucosa)



▲ Light micrograph (LM) of the stomach wall showing four concentric layers at low magnification. A thick mucosa (formed mostly of tightly packed gastric glands) lines the lumen (*). The rectangle indicates a ruga consisting of a submucosal connective tissue core covered by mucosa. A thick layer of mucus secreted by surface cells forms a barrier over the mucosa for protection of tissues from acid and proteolytic enzymes in the lumen. The submucosa (SM) has prominent blood vessels (BV). Serosa covers the muscularis externa (ME) externally. 10x, H&E.

Stomach (Ventriculus, Gaster)


- **Gastric mucosa**
- simple columnar epithelium
- surface epithelium produces mucus
(mucinogenic granules, high content of HCO_3^- , K^+)
= protective function
- **areae gastricae, foveolae gastricae**





Stomach (Ventriculus, Gaster)

- Gastric mucosa
- L. propria contains large amount of glands

- Gl. cardiacae
- Gl. pyloricae
- Gl. gastricae propriae





Mucous
Secretes
gastric
juice



Stomach (Ventriculus, Gaster)

- Gl. gastricae propriae
- glands of fundus and body
- simple tubular or branched
- 2-4 opens to the gastric pits

- four cell types of gl. gastricae propriae




Stomach(Ventriculus, Gaster)

Gl. gastricae propriae


chief

- most abundant, lower part of body and fundus of the gland
- pyramidal shape, basophilic cytoplasm, RER, pepsinogenic granules




parietal

- neck-body junction
- eosinophilic cytoplasm, high numbers of mtch., SER
- complex and dynamic ultrastructure
- intracellular canals in apical part with microvilli – membrane bound enzyme complexes producing H^+ & Cl^- (HCl originates extracellularly)




neck cells

- cubic, mucinous
- capable of regeneration of all cell types in gastric epithelium




Stomach (Ventriculus, Gaster)

Gl. gastricae propriae




(entero)endocrine

- minor, secretion
- granules
- different cell types with different sensitivity to various histological stainings
- secretion of various biologically active compounds
- DNES/APUD
- GIT chemosensing
- see lesson spring semester 2012 - Epithelial tissue




Type	Hormone	Localization/Function
D cells	Somatostatin	- Stomach, intestine, hepatic and pancreatic ducts
EC cells	Serotonin	- Stomach, gallbladder, intestine - Peristaltics
ECL cells	Histamin	- Stomach - HCl secretion
G cells	Gastrin	- Pars pylorica, duodenum - HCl, pepsin secretion
L (EG) cells	Enteroglucagon	- Stomach, intestine - attenuates secretion of pancreatic enzymes and peristaltics

Gastroduodenal junction



General architecture of the intestine




Small intestine – adaptation to efficient resorption

Four basic layers: **mucosa, submucosa, muscularis externa, serosa**

mucosa and submucosa maximise the resorptive area

- **plicae circulares** (Kerckringi) – **mucosa + submucosa**, ca 800, increase **2-3x**, distal region of duodenum



- **villae** (villi intestinales) – **mucosa** (l. propria + epithelium) 0,5-1,5 mm long, 10-40/mm², 4 000 000, increase **5-10x**
- **microvillae** – **apical part of enterocytes** – 1- 2 µm long, 0,1 µm wide, 100 mil./mm², increase **20x**

Intestine: adaptation to effective resorption



plicae circulares (Kerckring)
– 2-3x

villi (villi intestinales)
– 5-10x

microvilli (striated border)
– 20x



Crypts of Lieberkühn

200-600x



Crypts of Lieberkühn (gl. intestinales)

- simple tubular structures of intestinal mucosa, depth 0,3-0,5 mm
- pass through I. propria and open to lumen
- different cell types
 - secretion of digestive enzymes
 - epithelial renewal
 - enteroendocrine cells
 - immune response





Intestinal mucosa

Enterocytes

- tall, columnar cells
- nucleus located in basis of the cell
- apical surface modified- microvilli (3000) + glycocalyx (0,5μm) = *striated border (cuticle)*
- tight intercellular connections, interdigitations




Function:

- digestion – enzymatic complexes on microvilli membrane
- absorption and transport – passive, facilitated or active
- lipid uptake - chylomicrons





▲ EMs of enterocytes at low (Left) and high (Right) magnification. Apical microvilli (MV) make up a striated border and extend from free surfaces of the cells. A fuzzy glycocalyx (Gl) covers them. A terminal web (TW) of actin filaments in the apical cytoplasm reaches into microvilli. Intercellular junctions (circles) are between adjacent cells. The cytoplasm contains mitochondria (Mi), lysosomes (Ly), and smooth (SER) and rough (RER) endoplasmic reticulum. Left: 10,000×; Right: 50,000×.

Microvilli



Microvilli—with core filaments





Transportion and resorption



Intestinal mucosa


Goblet cells

- Cylindrical glandular epithelial cells
- Apical surface – apocrine/merocrine secretion of mucus
- Basal part – RER, GA, nucleus, mitochondria
- Mucinogenic granules
- see lesson spring semester 2015 - Epithelial tissue




Intestinal mucosa

Goblet cells




▲ LM of the colonic mucosa. Surface epithelium containing goblet cells (G) and enterocytes (En) invaginates to form an intestinal crypt. The lamina propria (LP), with capillaries (arrows) and larger blood vessels (BV), is richly cellular. 600 \times , Toluidine blue.



Intestinal mucosa


Paneth cells

- basal part of crypts of Lieberkühn
- basophilic cytoplasm
- GA located above nucleus
- acidophilic (red) granules
- immune system
- secretion granules contain biologically active substances e.g. lysozyme
- influence intestinal microflora



Enteroendocrine cells



- similar to gastric enteroendocrine cells
- regulate pancreatic secretions
- homeostatic axis (brain-intestine-adipose tissue)
- cholecystokinin, secretin, GIP, motilin, neurocrine peptides etc.



Intestinal mucosa

M cells (microfold)


- epithelial cells above Peyer's patches and lymphatic nodules
- no microvilli
- induces immune response
- MHCII
- antigen presentation to dendritic cells and lymphocytes





Intestinal mucosa

Intestinal stem cells

- bottom of crypts of Lieberkühn
- epithelial renewal (4-5 days)
- stem cell niche
- tumour transformation




Nature Reviews | Molecular Cell Biology



Intestinal mucosa

L. propria

- immune system – GALT
- immunologic barrier
- Peyer's patches



Submucose

Brunner's glands

- gl. duodenale Brunneri
- branched tuboalveolar glands, columnar mucinous cells
- connective tissue reduced to thin septa between glandular lobules
- open to crypts of Lieberkühn




Muscularis externa

- two layers of smooth muscle (inner circular, outer longitudinal)
- plexus myentericus Auerbachi

Serosa

- loose collagen connective tissue + simple squamous epithelium (mesothelium)




Colon

- no plicae of Kerckring, villi
- muscularis externa – longitudinal layer forms taenie coli
- surface serosa forms appendices epiploicae (adipose)





Small intestine



Colon


Colon

- absorption of water, electrolytes
- deeper crypts of Lieberkühn, no Paneth cells
- abundant goblet cells
- abundant lymphatic follicles in l. propria (GALT)




Appendix

- develops from and is connected to caecum 8-10 cm (0,5-1cm)
- continuous longitudinal layer of m. externa
- lymphatic follicles reaching submucosa
- irregular crypts of Lieberkühn with Paneth cells





Rectum and anal canal



- Pars pelvina
 - *plicae transversae recti*
 - histological architecture identical to colon
- Canalis analis
 - simple columnar epithelium replaced by stratified squamous epithelium
 - rich venous plexus
 - *columnae rectales, sinus rectales and valvulae rectales*
 - *zona cutanea* – typical skin
- Rectal submucosa – high and loose
(prolapse of mucose)



Rectum and anal canal



Primitive gut development






E7.5
Late gastrula






E8.5
Hindgut



E9.5
Primitive gut tube





Microscopic anatomy of the gut tube

- **General architecture of hollow organs and gut tube:** mucosa (l. epithelialis m., l. propria, l. muscularis m.), submucosa, t. muscularis externa, serosa (l. propria s., l. epith. s.), adventitia
- **Esophagus** - structure, epithelium, mucosal and submucosal glands, differences in t. muscularis ext.
- **Stomach** – anatomical and histological structure, mucosa - areae gastricae, foveolae gastricae, gastric glands (pyloricae vs. propriae), localization, ultrastructure and function of gl. gastricae propriae and its cells (chief, parietal, neck, enteroendocrine)
- **Small and large intestine, appendix** - anatomical and histological structure, mucosa, glands (crypts of Lieberkühn, Brunner's glands), cell types of intestinal mucosa, lymphatic system, modifications of intestinal wall
- **Rectum and anal canal** - anatomical and histological structure, mucosa, epithelium, description of associated structures

Embryonic development

- Development of **primitive gut** and its derivatives, embryonic flexion, differentiation and characteristics of individual regions and associated organs