

Microscopic structure of the sense organs

Aleš Hampl

December 2019

Sense system

It serves to convey stimuli that influence organism from inside and outside

Sensitive nerve endings

(with simple structure)

- Simple sensory endings
- Intraepithelial sensory endings
- Sensory bodies

Complex organs

- Photosensitive organ - Eye
- Organ of hearing and equilibrium - Ear

Photoreceptor organ - Eye

Analyzes the form, light intensity and colour reflected from objects


Eye ball

(three-layered structure)

- tunica externa = fibrosa
- tunica media = vasculosa
- tunica interna = nervosa

Accessory structures

- eye lids
- conjunctiva
- lacrimal apparatus
- muscles



What do we expect from the eye ?


- Ability to sense signals and transfer them to CNS
- Ability to focus on objects
- Enough strength
- Ability to regenerate
- Ability to move with a minimal friction

Enough strength

Eyes sit in the protective environment of the skull, in orbits, surrounded by the fat cussions..

$$\begin{array}{ccc} \text{Cornea} & + & \text{Sclera} \\ 1/6 & + & 5/6 \end{array} = \text{Tunica externa oculi}$$

= 6/6 of the surface



Cornea

- average thickness 0.9-1.0 mm
- colorless
- transparent
- thoroughly avascular
- 5 distinct layers

continues on the next slide

Sclera


- average thickness 0.5 mm (thicker in posterior part)
- bundles of flat collagen I fibers (intersecting in all directions)
- few fibroblasts, minimum ground substance
- relatively avascular
- connected by loose system of collagen fibers with **Tenon's capsule** - **Tenon's space** - allows for free movement of the eye
- **lamina suprachorioidea** - connection to choroid
(loose connective tissue with melanocytes, fibroblasts and elastic fibers)

Enough strength

Eyes sit in the protective environment of the skull, in orbits, surrounded by the fat cussions..

$$\begin{array}{ccc} \text{Cornea} & + & \text{Sclera} \\ 1/6 & + & 5/6 \end{array} = \text{Tunica externa oculi}$$

= 6/6 of the surface




Cornea

- average thickness 0.9-1.0 mm
- colorless
- transparent
- thoroughly avascular
- 5 distinct layers

continues on the next slide


Sclera

- average thickness 0.5 mm (thicker in posterior part)
- bundles of flat collagen I fibers (intersecting in all directions)
- few fibroblasts, minimum ground substance
- relatively avascular
- connected by loose system of collagen fibers with **Tenon's capsule** - **Tenon's space** - allows for free movement of the eye
- **lamina suprachorioidea** - connection to choroid
(loose connective tissue with melanocytes, fibroblasts and elastic fibers)



Cornea

(transversal section)



Anterior

Posterior

Corneal epithelium
Bowman's membrane

Substantia
propria
cornea
= STROMA

Descemet's membrane
Corneal endothelium

- stratified + squamous (5-6 layers)
- nonkeratinizing
- rich in nerve endings
- surface cells equipped with microvilli (protrude into the space with the film of tears)

- = Lamina limitans anterior
- thickness about 7 - 12 μm
 - fine collagen fibers (intersecting in all directions)
 - no cells
 - provides strength

- many layers of collagen fibers (in right angles)
- flat keratocytes in between the collagen lamellae (fibroblast-like cells)
- contains mucoid substance rich in chondroitinsulphate
- properly hydrated

KEY to the TRANSPARENCY

- = Lamina limitans posterior
- fine collagen fibers
 - fibers are organized to 3D network






- simple + squamous
- active in transport to maintain cornea in a proper state
- continues on the frontal part of iris (via spongium anguli iridocornealis)


Ability to regenerate

Limbus - corneoscleral junction

The area of transition of the transparent collagen bundles of cornea into the opaque collagen bundles of sclera.

Highly vascularized - feeds avascular cornea





Enough supply of resources

$$\text{Choroid} + \text{Ciliary body} + \text{Iris} = \text{Tunica media}$$

Choroidea Corpus ciliare Iris

T. vasculosa

Choroid = 4-layered structure

Lamina suprachoroidea

- loose connective tissue
- rich for pigment cells - melanocytes

Lamina vasculosa

- loose connective tissue
- rich for pigment cells - melanocytes
- contains larger vessels and nerves


Lamina choriocapillaris

- loose connective tissue
- network of small vessels

Lamina vitrea = L. basalis = Bruch's membrane

- fibers of collagen and elastin
- overall thickness about 3-4 µm
- links together basal laminae of Lamina choriocapillaris of choroid and pigmented epithelium of retina

Choroid



Sclera


Retina

Ability to focus on objects

Ciliary body - anterior extension of the choroid

Stroma of ciliary body

- loose connective tissue
- contains elastic fibers, vessels and melanocytes
- rich for capillaries (chamber fluid)
- bundles of smooth muscle fibers (anchored to sclera and protrude to the processes of ciliary body - *m. ciliaris*)





Processes of CB (Processus ciliares)

- protrude into posterior chamber
- total number of about 70-80
- rich for capillaries (chamber fluid)
- covered by two-layered epithelium (from the retina - *pars ciliaris retinae*)
- linked to the lens capsula *fibrae suspensoriae lentis* (zonulae)


Triangular on crosssection

Connects to:

- lens + posterior chamber (A)
- sclera (B)
- vitreous body (C)




Ciliary body




Lens accommodation

Closer sight



Longer sight




- muscle contracted
- zonulae loosened


- muscle relaxed
- zonulae stretched

Iris - 1

Anterior continuation of the choroid.
Partially covers the lens.



Iris - 2






Iris = 4-layered structure
Layers from outside:

- 1. Anterior epithelium**
 - continuation of the posterior ep. of the cornea
 - discontinuous layer of flat epithelial cells, fibroblasts & melanocytes
- 2. Anterior border layer**
 - thin layer of connective tissue
 - rich for pigmented cells - melanocytes
 - decides about eye colour**
- 3. Stroma**
 - loose connective tissue
 - large number of radially running vessels
 - concentrically ordered smooth muscle fibers (=musculus sphincter pupillae)
- 4. Pars iridica retinae**
 - 2-layered
 - continues from ciliary body
 - layer facing the stroma contains smooth muscle fibers (=musculus dilatator pupillae)

Lens

Capsule + Epithelium + Fibers

- 10-20 μm
- Collagen IV




Epithelium (cuboidal + low cylindrical) only on the anterior surface.


Fibrae suspensorie lentis are anchored to the equator of the lens.

Ability to sense signals and transfer them to CNS for processing

Retina = Tunica aculi interna - Tunica nervosa




Invagination of prosencephalon creates two-layered **optic cup**.




Neural (optical) retina

minimum 15 different types of neurons with tens of interactions (synapses)




Photoreceptors = Rod and cone cells 1

I. Neurones of the optical path




Photoreceptors = Rod and cone cells 2

I. Neurones of the optical path



Other neurons of the optical path 1



Other neurons of the optical path 2

II. neuron Bipolar cells

Diffuse

- Synapses with two or more receptors

Monosynaptic

- Synapses with only one receptor
- Direct transfer of impulses from some rods

III. neuron Ganglion cells (multipolar)

- Large cells
- Nuclei mainly in one layer
- Dendrites connect to neurites of bipolar and amakrine cells
- Neurites run in 9. layer of the retina and come together to form optic nerve



Associating + integrating neurons

Horizontal cells


- Small
- Multipolar

Amacrine cells

- They don't have neurite




Supporting cells of the retina 1





Supporting cells of the retina 1

Muller cells


= modified glial cells of the CNS






„Does the retina see
the same in all its areas“

Central x Peripheral vision




Fovea centralis of the macula lutea
= the sharpest vision



Audiorceptor system


=

Vestibulocochlear apparatus




External + Middle ear - Organ of hearing



Middle ear - fitted in the cavities of temporal bone along with internal ear - osseous labyrinth.



Extrenal ear - Auricle



Extrenal ear - Auricle - Elastic cartilage



Internal ear


Bony labyrinth

- series of cavities
- petrous portion of temporal bone

Vestibulum

Cochlea

Semicircular
canals



Membranous labyrinth


- series of interconnected tubes and vesicles
- lined by epithelium
- positioned in bony labyrinth

Utricle

Saccule



Semicircular
ducts

Cochlear
duct




Internal ear - Organ of hearing

Cochlea
• 2,5 turns around modiolus
• total length about 35 mm



Modiolus (bony core)
• central axis of cochlear duct
• contains ganglion spirale cochleae, nervus cochlearis and vessels

Internal ear - Detail of cochlear duct







Scala media = Ductus cochlearis

20 μ m


Internal ear - Organ of Corti - 1





Inner hair cells Outer hair cells

20 μ m



Internal ear - Organ of Corti - 2

Secondary receptor cells



- hearing hairs - stereocilia
- in contact with tectorial membrane
- bases wrapped by dendrites of bipolar cells of ganglion spirale

Outer hair cells

- 3-5 rows, ~12 000, no axonema

Inner hair cells

- 1 row, ~3 500, no axon.



Supporting cells

Hensen's cells

Outer phalangeal cells

- support to hair cells, which run through the spaces between ph.


Outer pillar of Corti

Outer pillar of Corti

Inner phalangeal cells

- same as inner ph. cells


Border cells






Hair and phalangeal cells




Spiral ganglion



Inner ear - Principle of hearing



Inner ear - Statokinetic / Vestibular organ - 1



Uniform composition of the wall (vesicles and tubes)
Thin layer of connective tissue + single-layer squamous/cuboidal epithelium.



Unifying concept of the composition of sensing elements
(vesicles - **maculae**; tubes - **cristae ampullares**)

Thickening of the wall with neuroepithelial cells innervated by branches of n. vestibularis.

Inner ear - Statokinetic / Vestibular organ - 2

Sensing of static equilibrium (maculae = static spots)

Deviation from the gravity force (gravity force of otoliths)
max. pressure - max. pull



Position of maculae

Saccule
bottom

Utricle
lateral wall

Kinocilium
(no axonema)

Terminal plexus
of tonofibrils

Stereocilia
(microvilli)



Hair cells

Supporting cell

Afferent innervation


Efferent innervation

Fibers of n. vestibularis
in connective basis
of macula




Inner ear - Statokinetic / Vestibular organ - 3

Sensing of dynamic equilibrium (cristae ampulares)





Reaction on acceleration/deceleration
(movement of endolymph)




Position of crista ampularis


- in ampules of semicircular ducts
- ridges perpendicular to axis of SDs







Membranous labyrinth



Development of sense organs - Overall picture




Sense organs - Reminder - Neural tube




Sense organs - Reminder - Neural crest

Arise from both
dorsal epidermis and neural plate




Sense organs - Cranial neural tube + Placodes



Brain after 4 weeks of development

Placodes: patches of dense columnar epithelium in the epidermis covering the head - their formation is induced by underlying brain and mesenchymal tissue - **develop in week 4**



Dorsolateral placodes

Contribute to:


- eye - lens placode
- ear - otic placode
- nose - nasal placode
- sensory ganglia

Epibranchial placodes

Develop into:

- sensory ganglia of cranial nerves (V, VII, IX, X)

Sense organs - Eye development 1




Brain after 5 weeks of development

Neural plate ectoderm → prosencephalon (forebrain) **eye fields** →

→ neural plate growth carries eye field region forward →

→ eye field invaginates forming **optic grooves (sulci)**

Sense organs - Eye development 2




Lens placode: the ectoderm invaginates in response to signals from the optic cup underneath. It then pinches off as a lens vesicle. Cells elongate to fill the vesicle and start to synthesize crystallins.

Optic cup: forms from the neural tube by invagination. The opening (choroid fissure) closes forming a round optic cup, an extension of the brain.

Optic stalk: connection to the brain that is filled with neurons to form the optic nerve.


Reciprocal interaction: the lens induces the formation of the optic cup and the cup regulates formation of the lens.

Sense organs - Eye development 3




Diencephalon - transversal section - 6 weeks



Sense organs - Eye development 4




Ear development - Inner ear 1



Ear development - Inner ear 2



Apoptosis





Dorsal part

Ventral part



Ear development - Middle ear

Pharyngeal arches 1 + 2




Ossicles

- Mesenchyme of neural crest origin (phar. arches 1+2)
- Embedded in mesenchyme
- Apoptosis late in pregnancy



Ear development - External ear



External auditory meatus - ectoderm

Thank you for your attention !