HEAT, COLD, ELECTRICAL, CHEMICAL INJURIES

I. Čundrle

Department of Anesthesiology and Intensive Care

FNUSA

BURN INJURIES

•





PATHOPHYSIOLOGY

• 45°C

•

- vasoactive mediators are released from damaged tissue loss of plasma into the intersticium hypovolemic shock hypoperfusion organ damage
- Kidney
- GIT
- Lung
- Metabolism

CAUSES

- - •
- •

THERMAL BURNS

- Contact
- Scald
- Flame
 - + possible inhalation trauma
 - + possible intoxication

ELECTRICAL BURNS

- •
- •
- •
- - Electrical damage
 - Electrical arch, lightning






• Acids




Alkali




• Other substances



DAMAGE EXTENT ESTIMATION – RULE OF 9



DEPTH OF BURN INJURIES



First-degree burn

Second-degree burn

Third-degree burn

FIRTS DEGREE

• First degree



SECOND DEGREE

• Second degree

• Ila

• Ilb



THIRD DEGREE

• Third degree




FOURTH DEGREE

• Fourth degree



INHALATION INJURY





SEVERE BURN

- Kids up to 2 years 5 % body surface area
- •
- •
- Kids third degree always
- Adults 20 % body surface area

• + always

SPECIALIZED BURN CLINICS

- •
- •
- •

FIRTS AID

- Remove the source of burn
- vital signs and secure vital functions
- Cooling of the wound

- Anti-shock precautions
- Transport


HEATSTROKE

•

hot and wet environment

FIRST AID

- Lower the body temperature
- Cold fluids



C Healthwise, Incorporated

SUNSTROKE

• Direct effect of sunlight on head

• First to second degree burns

FIRST AID

- •

HYPOTHERMIA

• Symptoms

- - Moderate hypothermia
 - Severe hypothermia

FROSTBITES

FIRST DEGREE

- •



SECOND DEGREE

•



THIRD DEGREE





FIRTS AID

•

- •
- •
- •

• Cause not known

• Acids

Alkali

INGESTION

- •
- •
- - •
- •
- •



ELECTRICAL BURNS

• Low voltage1000 V

• High voltage > 1000 V

• Lightning

THANK YOU FOR YOUR ATTENTION

I BROS

and the second second

AND ADDRESS OF ADDRES