


# The basic principles of gerontology



The modern strategy of health support  
and increasing of independence of seniors  
Brno, October-November 2017

# Gerontology

- ▶ the body of knowledge on ageing, about the problems of aging people and life in old age




# Gerontology subspecialties I

- ▶ **experimental gerontology** - causes and ways of ageing, actually at the cellular and molecular level, neuropsychology of ageing
- ▶ **social gerontology** - relationship between aging people and society, needs of elderly, demography, sociology, economy, law, urbanistics, architecture etc
- ▶ **clinical gerontology** - geriatrics


# Gerontology subspecialties II

► **geriatrics** - summarizes and generalizes across all disciplines main topics of senior´s health and functional status, specific needs, specificities of appearance, symptoms, therapy, prevention and social context of diseases of old age

VĚKOVÁ STRUKTURA OBYVATELSTVA ČR K 31. 12. 2010  
AGE STRUCTURE OF THE CR POPULATION AS AT 31 DECEMBER 2010



Věková skladba obyvatelstva: 2050  
Česká republika




\*1960


\*1975

\*1990

Expected changes of the population  
age-structure 2010-2050



# Life expectancy and infant mortality rate



# Absolute numbers of live births and deaths 1785-2011

# Specific features of diseases in elderly

## Risk of false diagnosis

# Oligosymptomatology

- ▶ expression of less typical symptoms
  - peritonitis without defence musculaire
  - ⇒ pneumonia without fever
  - ⇒ cystitis with polakisuria, but without pain
  - ⇒ tachyfibrillation only in hyperthyreosis

# Microsymptomatology

- uroinfection without fever
- incomplete inflammation symptomatology
- myocardial infarction without typical stenocardia, but with chest tightness only
- florid ulcer disease with dyspepsia, but without typical pain
- inflammation leucocytosis absent



## „Another organ cries“

- ▶ current disease burden most frail organs
  - ⇒ cardiac failure because of pneumonia
  - ⇒ confusion caused by sepsis, urosepsis
  - ⇒ stenocardia more expressed in anemia
  - ⇒ TIA in anemia, cardiac failure, myocardial infarction


# Polymorbidity

- ▶ the number of chronic diseases increases with age
- ▶ 80% of patients above 80 years suffer from more than one chronic disease
- ▶ diseases influence each other - more frequently negatively
- ▶ polypragmasia, compliance, interaction
- ▶ long term recovery
- ▶ risk of immobilization



# Glacier like symptom

- apparent symptomatology is the little part of reality only
- dyspnea in myocardial infarction only
- confusion in cardiac failure
- confusion in acute abdomen
- dementia progression caused by chronic pain



# Interdisciplinary problems

**geriatric giants „4 I“**

**instability**

**cognitive disturbances**

**imobilization**

**incontinentia, skin integrity  
disorders**

# **Specificities and peculiarities of pharmacotherapy in elderly**

**Problem topics**  
**Farmacokinetics**  
**Compliance**

# Problem topics

- pharmacokinetics,  
pharmacodynamics
- compliance
- polymorbidity
- polypragmasia
- medications market
- the patient's wishes
- treatment coordination
- „external“ influences



# Farmakokinetics I

- decrease of gastric acidity
- decrease of gastric motility
- reduced GIT blood flow
- slower resorption

# Farmacokinetics II

- decreased distribution volume for hydrosoluble substances
- increased distribution volume for liposoluble substances
- decreased liver and kidney function
- decreased albumin concentration

# Compliance and its changes in elderly I

- ▶ reciprocal association between compliance and number of medications used
  - 5 medications take exactly 33-44%,
  - 10 medications 10-20% only
- ▶ influence of relatives and caregivers
- ▶ dependence on specialised supervision

# Compliance and its changes in elderly II

- medication price influence
- user's comfort
- medication shape and color
- content of package leaflet



# Polypragmasia? Polypharmacotherapy?

- ▶ tackle fundamental problems
- ▶ improve the quality of life
- ▶ profylactic medications
- ▶ number of medications limitation?
- ▶ respecting of guidelines
- ▶ unwanted symptoms induced by therapy  
express 24-28% patients, 90% of symptoms  
are predictable



# Therapy coordination problems

- ❖ „gate keeping“ x confidence in the knowledge of GP
- ❖ „travelling“ around out-patient clinics
- ❖ addition of recommended treatments
- ❖ lack of communication between GPs and specialists
- ❖ financial limitations of GPs and specialists
- ❖ doubled generics



# Medication at the market

- ▶ many market names of the same generic substance
- ▶ the elderly patient remembers the medication according to shape and colour
- ▶ the influence of advertisement
- ▶ the influence of friends or neighbors „me too“



# Seniors and medications consumption

- ▶ age group 60-75 years creates 15% of population
- consumes 33% prescription medications
- consumes 40% OTC medications

# Creating the medication schedule

- ▶ one coordinator
- ▶ specialist's recommendations
- ▶ substantial medications
- ▶ or to know or to consult

# Ten rules for elderly prescription

- » 1. Define substantial problems to treat
- » 2. Define treatment targets
- » 3. Consider alternative methods including education and non pharmacologicla methods
- » 4. Consider all risks and risk medications already taken
- » 5. Optimal dosage “start low go slow”

# Ten rules for elderly prescription II

- » 6. Select the simplest schedule
- » 7. Consider the risk of cumulation in retarded medications
- » 8. Prepare the table containing recommended medications and ask the patient about understanding
- » 9. Ask the use of OTC or other substances
- » 10. Consider the possibility to stop the taking of some medication

# Non-pharmacological therapy

- ▶ positive alternative to polypragmasia
- ▶ regime measures - sleeping rhythm, to use the bed for sleeping only, regular day and week rhythm
- ▶ reduction of harmful habits
- ▶ change of eating habits - regular warm dishes, care for oral cavity and teeths

# Comprehensive geriatric assessment

# **Comprehensive geriatric assessment (CGA)**

- personality**
- somatic health**
- functional status**
- psychical health**
- social context**

# Personality

- ✓ life situations
- ✓ priorities and decisions - treat/not to treat, reanimate/not to reanimate, decisions in dementia
- ✓ subjective quality of life




# Somatic health

- ✓ diseases - main diseases, other diseases
- ✓ functional burden of diseases
- ✓ syndromological dg (imobilization, incontinentia ...)

# Functional efficiency

- ✓ stability and walking
- ✓ performance and independence
- ✓ physical condition
- ✓ nutrition



# Mental health



- ✓ cognitive and fatic disorders and deliria - active screening and evaluation
- ✓ affective disorders (depression) - active screening and evaluation
- ✓ mental balance, maladaptation, the influence of psychosocial stressors

# Social context

- ✓ social roles and relationships (social network)
- ✓ operation demands and safety of the home environment
- ✓ social needs supplied or claimed



# Evaluation of stability and walking disorders

- ❖ basic neurological assessment
- ❖ getting up from lying to a sitting position and from sitting position to standing
- ❖ spontaneous standing
- ❖ maneuvers in standing - Romberg, pull test, push test
- ❖ spontaneous walking - 10m - base width, lenght of the step, fluidity of movement, start and stop, rotation, obstacles
- ❖ maneuvers in walking - on heels, on tiptoes, with closed eyes, backwards, tandem walking

# Possible pathologies

- ✓ walking of width base with unstable destination
- ✓ polyneuropathic walking - uncertainty, weakness of lower extremities
- ✓ cerebellar walking - like ebrietas
- ✓ choreatic walking
- ✓ short step, stiffness
- ✓ inability to start the step



# Evaluation of physical performance

- anamnestic - comparison with contemporaries, with standards - ADL, IADL
- stress tests - speed evaluation, observation of EKG, blood pressure, heart rate
- selection of tests - izometric, izotonic, treadmill



<i>Činnost</i>	<i>Provedení činnosti</i>	<i>Bodové skóre</i>
1. najedení, napítí	samostatně bez pomoci s pomocí neproveďe	10 5 0
2. oblékání	samostatně bez pomoci s pomocí neproveďe	10 5 0
3. koupání	samostatně nebo s pomocí neproveďe	5 0
4. osobní hygiena	samostatně nebo s pomocí neproveďe	5 0
5. kontinence stolice	plně kontinentní občas inkontinentní trvale inkontinentní	10 5 0
6. kontinence moči	plně kontinentní občas inkontinentní inkontinentní	10 5 0
7. použití WC	samostatně bez pomoci s pomocí neproveďe	10 5 0
8. přesun lůžko - židle	samostatně bez pomoci s malou pomocí vydrží sedět neproveďe	15 10 5 0
9. chůze po rovině	samostatně nad 50 m s pomocí 50 m na vozíku 50 m neproveďe	15 10 5 0
10. chůze po schodech	samostatně bez pomoci s pomocí neproveďe	10 5 0

Hodnocení stupně závislosti v základních výedních činnostech:

0 - 40 bodů: vysoce závislý

45 - 60 bodů: závislost středního stupně

65 - 95 bodů: lehká závislost

100 bodů: nezávislý

<i>Činnost</i>	<i>Hodnocení</i>	<i>Bodové skóre</i>
1. telefonování	vyhledá samostatně číslo, vytvoří je zná několik čísel, odpovídá na zavolání nedokáže použít telefon	10 5 0
2. transport	cestuje samostatně dopravním prostředkem cestuje, je-li doprovázen vyžaduje pomoc druhé osoby, speciálně upravený vůz apod.	10 5 0
3. nakupování	dojde samostatně nakoupit nakoupí s doprovodem a radou druhé osoby neschopen bez podstatné pomoci	10 5 0
4. vaření	uvaří samostatně celé jídlo jídlo ohřeje jídlo musí být připraveno druhou osobou	10 5 0
5. domácí práce	udržuje domácnost s výjimkou těžkých prací provede pouze lehké práce nebo neudrží přiměřenou čistotu potřebuje pomoc při většině prací nebo se práce v domácnosti neúčastní	10 5 0
6. práce kolem domu	provádí samostatně a pravidelně provede pod dohledem vyžaduje pomoc, neprovádí	10 5 0
7. užívání léků	samostatně v určenou dobu správnou dávku, zná názvy léků užívá, jsou-li připraveny a připomenuty léky musí být podány druhou osobou	10 5 0
8. finance	spravuje samostatně, platí účty, zná příjmy a výdaje zvládne drobné výdaje, potřebuje pomoc se složitějšími operacemi neschopen bez pomoci zacházet s penězi	10 5 0

Hodnocení stupně závislosti v instrumentálních všedních činnostech:

0 - 40 bodů: závislý v IADL

45 - 75 bodů: částečně závislý v IADL

80 bodů: nezávislý v IADL

# Cognitive performance evaluation

## ► MMSE

- Mini Mental State Examination 30-27-23-18-13
- clock test
- test connecting numbers and letters

Tab. 4. Test kognitivních funkcí - Mini-Mental State Exam (MMSE)

Oblast hodnocení	Max. skóre
<b>1. Orientace</b> Položte nemocnému 10 otázek. Za každou správnou odpověď započtejte 1 bod. Který je tedy rok? Které je roční období? Můžete mi říci dnešní datum? Který je den v týdnu? Který je tedy měsíc? Ve kterém jsme státeč? Ve které jsme zemi? Ve kterém jsme městě? Jak se jmenuje tato nemocnice (ordinace)? Ve kterém jsme poschodi?	1 1 1 1 1 1 1 1 1 1
<b>2. Vštípivost</b> Výšetřující jmenuje 3 předměty (lopata, šátek, váza) a vyzve pacienta, aby je opakoval. Za každou správnou odpověď bez ohledu na pořadí se započítá jeden bod. Jestliže pacient nereprodukuje všechny 3, opakujte je tak dlouho, dokud se je nenaúčí.	3
<b>3. Pozornost a počítání</b> Vyzvěte nemocného, aby od čísla 100 odečtejte stále číslo 7 (93-86-79-72-65). Skončete po 5 odečtech. Každou správnou odpověď hodnotěte 1 bodem. Alternativou počítání může být hláskování slova "pokrm" pozadu po jednotlivých hláskách.	5
<b>4. Výbavnost</b> Vyzvěte nemocného, aby opakoval 3 slova, která si měl zapamatovat. 1 bod za každou správnou odpověď.	3
<b>5. Řeč, komunikace a konstrukční schopnosti</b> Ukažte nemocnému dva předměty (tužka, hodinky) a vyzvěte ho, aby je pojmenoval. Za správnou odpověď po jednom bodu. Vyzvěte nemocného, aby po Vás opakoval "Žádná ale, jestliže a kdyby". Správné opakování hodnote jedním bodem. Dejte nemocnému třístupňový příkaz (vezměte papír do pravé ruky), (přeložte ho napří) a (položte na podlahu). Za každý správně provedený stupeň započtejte 1 bod. Dejte nemocnému přečíst kartu s nápisem "Zavřete oči". Započtejte 1 bod za zavření očí. Vyzvěte nemocného, aby napsal smysluplnou větu. 1 bod za větu obsahující podmět a příslušek, která dává smysl.	2 1 3 1 1


Vyzvěte nemocného, aby na zvláštní papír nakreslil obrazec podle předlohy. 1 bod, pouze jsou-li zachovány všechny úhly a protnutí vytváří čtyřúhelník.






Celkové skóre

**Hodnocení:** skóre 23 bodů a nižší svědčí pro kognitivní poruchu (delirium nebo demence). Test má senzitivitu 87 % a specifitu 82 %.

# Connecting numbers and letters



# Clock test



# Depression evaluation - geriatric depression scale GDS



Otzávka	Odpověď
Jste v zásadě spokojen(a) se svým životem?	Ano – Ne
Opustil(a) jste mnoho ze svých aktivit a zájmů?	Ano – Ne
Máte pocit, že Váš život je prázdný?	Ano – Ne
Pociťujete často nudu, prázdnnotu?	Ano – Ne
Máte většinou dobrou náladu?	Ano – Ne
Obáváte se, že se vám přihodí něco zlého?	Ano – Ne
Jste většinou šťastný (šťastná)?	Ano – Ne
Cítíte se často bezmocný (bezmocná)?	Ano – Ne
Zůstáváte raději doma, než byste šel (šla) ven a podnikal(a) nové věci?	Ano – Ne
Máte pocit, že máte více problémů s pamětí než ostatní?	Ano – Ne
Myslíte, že je pěkné žít v této době?	Ano – Ne
Tak, jak jste na tom právě nyní, připadáte si bezcenný (bezcenná)?	Ano – Ne
Cítíte se plný (plná) energie?	Ano – Ne
Pociťujete svou situaci jako beznadějnou?	Ano – Ne
Myslíte si, že většina lidí je na tom lépe než vy?	Ano – Ne

Každá tučně označená odpověď = 1 bod. Hodnocení: 0–5 bodů norma, 6 a více bodů suspektní deprese (česká standardizace podle Tošnerové). Původní hodnocení z r. 1999: 0–7 norma, 8–12 mírná deprese, 13–15 těžká deprese.

Poslední 3 měsíce ztráta chuti k jídlu, obtíže GIT, problémy se žvýkáním a polykáním 0 = těžké poruchy 1 = mírné 2 = bez potíží
Ztráta tělesné hmotnosti v posledním měsíci 0 = více než 3 kg 1 = neví 2 = v rozmezí 1-3kg 3 = stabilní hmotnost
Pohyblivost 0 = upoután na lůžko 1 = pohyb v okolí lůžka, po místnosti 3 = vychází ven
Psychický stres v posledních 3 měsících 0 = ano 2 = ne
Neuropsychické problémy 0 = těžká demence, deprese 1 = mírná demence 2 = žádné problémy
Index tělesné hmotnosti BMI 0 = < 19 1 = 19 – 21 2 = 21 – 23 3 = > 23
maximum – 14 bodů
norma - 12 bodů
riziko malnutrice - < 11 bodů

# MNA II

Žije v domácnosti

0 = ne

1 = ano

Užívá více než 3 léky denně

0 = ne

1 = ano

Dekubity

0 = ano

1 = ne

Počet hlavních jídel denně

0 = 1 jídlo

1 = 2 jídla

2 = 3 jídla

Příjem proteinů

1x a vícekrát denně mléčné výrobky

ano      ne

2x a vícekrát denně vejce a luštěninu

ano      ne

maso, ryby denně

ano      ne

0 = 0-1x ano

0,5 = 2x ano

1 = 3x ano

Ovoce a zelenina denně

0 = ne

1 = ano

Příjem tekutin

0 = méně než 3 šálky

0,5 = 3-5 šálků

1 = 5 a více šálků

# MNA III

Způsob příjmu potravy

- 0 = s dopomocí
- 1 = samostatně s obtížemi
- 2 = samostatně bez problémů

Vlastní hodnocení stavu výživy

- 0 = podvyživený
- 1 = nehodnotí
- 2 = nemá nutriční problémy

Hodnocení vlastního zdravotního stavu ve srovnání s vrstevníky

- 0 = nedobrý
- 0,5 = neví
- 1 = stejně dobrý
- 2 = lepší

Střední obvod paže

- 0 = <21cm
- 0,5 = 21-22cm
- 1 = > 22cm

Obvod lýtku

- 0 = < 31 cm
- 1 = 31 cm a více

Zhodnocení – max. 14 bodů

**Celkové hodnocení z obou částí tabulky – 30 bodů**

- maximum

- 17-23,5 bodů** – riziko malnutrice
- < 17 bodů** – malnutrice

# Thank you for your attention

