

M U N I

M E D


M U N I
M E D

5

Somatosensitivity, viscerosensitivity, proprioception and pain I


The role of nervous system

ANTICIPATION




Receptors/sensors

- Energy convertor
 - Signal reception
 - Signal transformation
- Receptor potential
 - Generator potential
- Action potential



<http://www.slideshare.net/CsillaEgri/presentations>


Receptor/generator and action potential



<http://www.slideshare.net/drpsdeb/presentations>

Receptors/sensors

- Energy convertor
 - Signal reception
 - Signal transformation
- Receptor potential
 - Generator potential
- Action potential
- Adequate stimulus
- Non adequate stimulus




<http://www.slideshare.net/CsillaEgri/presentations>

Receptors/sensors

- Energy convertor
 - Signal reception
 - Signal transformation
- Receptor potential
 - Generator potential
- Action potential
- Adequate stimulus
- Non adequate stimulus

✓ **Mechanoreceptors**
✓ **Thermoreceptors**
✓ **Chemoreceptors**
✓ **Fotoreceptors**




<http://www.slideshare.net/CsillaEgri/presentations>

Receptors/sensors

- Energy convertor
 - Signal reception
 - Signal transformation
- Receptor potential
 - Generator potential
- Action potentials
- Adequate stimulus
- Non adequate stimulus

Basic attributes of stimulus
Qualitative
Modality - What?
Localization - Where?



<http://www.slideshare.net/CsillaEgri/presentations>


Receptors/sensors

- Energy convertor
 - Signal reception
 - Signal transformation
- Receptor potential
 - Generator potential
- Action potentials
- Adequate stimulus
- Non adequate stimulus

Basic attributes of stimulus

Qualitative
Modality - What?
Localization - Where?

Quantitative
Intensity - How much?



<http://www.slideshare.net/CsillaEgri/presentations>


Receptors/sensors

- Energy convertor
 - Signal reception
 - Signal transformation
- Receptor potential
 - Generator potential
- Action potentials
- Adequate stimulus
- Non adequate stimulus

Basic attributes of stimulus

Qualitative
Modality - What?
Localization - Where?

Quantitative
Intensity - How much?
Duration




<http://www.slideshare.net/CsillaEgri/presentations>

Intensity coding

How much?

- Amplitude of receptor potential is transduced into the frequency of AP




<http://www.slideshare.net/CsillaEgri/presentations>

Intensity coding

How much?

- In the other words: an increased intensity is associated with increase in frequency of AP
- A high-intensity stimulus may also activate more receptors





<http://neuronresearch.net/neuron/files/neuralcode.htm>

Intensity coding

How much?

- Relation between receptor and action potential is logarithmic



<http://slideplayer.cz/slide/3217923/>


Qualitative information

What?
Where?

- The law of specific nerve energies:

The nature of perception is defined by the pathway over which the sensory information is carried



- Labeled line coding define the information about quality



Qualitative information



What?
Where?

- Labeled line coding
- Receptive field
- Nerve stimulation mimics receptor stimulation




Receptive fields

- Various size and overlap
- Small receptive field – high resolution
- Spatial resolving power increased by lateral inhibition



<http://www.slideshare.net/drpsdeb/presentations>


Lateral inhibition



<http://www.slideshare.net/drpsdeb/presentations>

Receptor adaptation

- The decline of receptor responses in spite of stimulus presence
- Tonic receptors – slow adaptation – presence of stimulus, position
- Phasic receptors – rapid adaptation – change of stimulus



<http://www.slideshare.net/CsillaEgri/presentations>

Receptors

- General
 - Superficial – somatosensors
 - Deep – viscerosensors
 - Muscles, tendons, joints – proprioceptors
- Special
 - Part of sensory organs

Receptors


- ✓ **Mechanoreceptors**
- ✓ **Thermoreceptors**
- ✓ **Chemoreceptors**
- ✓ **Fotoreceptors**

- **General**
 - Superficial – somatosensors
 - Deep – viscerosensors
 - Muscles, tendons, joints – proprioceptors
- **Special**
 - Part of sensory organs

Receptors

- ✓ Mechanoreceptors
- ✓ Thermoreceptors
- ✓ Chemoreceptors
- ✓ Fotoreceptors


- Simple
 - Superficial – somatosensors
 - Deep – viscerosensors
 - Muscles, tendons, joints – proprioceptors
- General
 - Part of sensory organs
- Special
 - Part of sensory organs



<http://www.slideshare.net/CsillaEgri/presentations>

Somato/viscero/ proprio


- Somatosensitivity
 - Pain
 - Temperature
 - Touch
- Viscerosensitivity
 - Pain
- Proprioception
 - Position
 - Movement



Somato/viscero/ proprio


- Somatosensitivity
 - Pain
 - Temperature
 - Touch
- Viscerosensitivity
 - Pain
- Proprioception
 - Position
 - Movement

The majority of information does not reach consciousness




Evolutionary point of view

- The signals indicating potential damage are the most important and the corresponding systems evolved early
 - Pain
 - Temperature



Evolutionary point of view

- The signals indicating potential damage are the most important and the corresponding systems evolved early
 - Pain
 - Temperature
- The touch signals have adaptive value and evolved later




Evolutionary point of view

- The signals indicating potential damage are the most important for the corresponding early evolution involved
 - Temperature
- The touch receptors have an adaptive value and a higher conduction velocity


Immediate survival

Long-term survival




Evolutionary point of view

- The signals indicating potential damage are the most important, the corresponding early responses involved
 - Temperature
- The touch receptors have adaptive value and their evolution is related to
 - Immediate survival
 - Long-term survival
- The structure of the receptor, nerve fibers and pathways reflects the evolution







Free nerve endings

- Non-specialized nerve endings
- Polymodal
 - Nociception
 - Thermoreception
 - Mechanoreception
- A delta fibres
- C fibres



Nerve fibres

| | A α | A β | A δ | C |
|----------------------------|---|---|---|---|
| Axons from skin | | | | |
| Axons from muscles | Group I | II | III | IV |
| |  |  |  |  |
| Diameter (μm) | 13–20 | 6–12 | 1–5 | 0.2–1.5 |
| Speed (m/sec) | 80–120 | 35–75 | 5–30 | 0.5–2 |
| Sensory receptors | Proprioceptors of skeletal muscle | Mechanoreceptors of skin | Pain, temperature | Temperature, pain, itch |

Nociceptors


- Free nerve endings responding to high-intensity stimuli
- Stimulus
 - Mechanical
 - ✓ High pressure
 - ✓ Sharp object
 - Thermal
 - ✓ Above approx. 45°C
 - ✓ Low threshold – variable
 - Chemical
 - ✓ pH
 - ✓ Mediators of inflammation and so on

A delta fibers
– sharp, localised pain

C fibers
– dull, diffuse pain


Thermoreceptors

- Free nerve endings receptive to thermal stimuli
- TRP (transient receptor potential) channels
 - Polymodal receptor (chemoreception, thermoreception)
 - Present also in many cells (including neurons, keratinocytes, mechanoreceptors)



Thermoreceptors

- Perceived temperature is determined by relative activity of cold and warm receptors




Hall: Guyton and Hall Textbook of Medical Physiology, 12th Edition
Copyright © 2011 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

<http://www.slideshare.net/CsillaEgri/presentations>

Thermoreceptors


- Mostly phasic response




<http://www.slideshare.net/CsillaEgri/presentations>

The receptors of the skin


- Simple versus complex




The receptors of the skin



B Location



C Intensity and time course



The receptors of the skin

| Receptor | Type | Sensation | Signals | Adaptation |
|---------------------------|-----------------------------|-----------------------------|---|--------------------------------|
| Meissner corpuscle | Encapsulated & layered | Touch: Flutter & Movement | Frequency/Velocity & Direction | Rapid |
| Pacinian corpuscle | Encapsulated & layered | Touch: Vibration | Frequency: 100-300 Hz | Rapid |
| Ruffini corpuscle | Encapsulated collagen | Touch: Skin Stretch | Direction & Force | Slow |
| Hair follicle | Unencapsulated | Touch: Movement | Direction & Velocity | Rapid |
| Merkel complex | Specialized epithelial cell | Touch, Pressure, Form | Location & Magnitude | Slow |
| Free Nerve Ending | Unencapsulated | Pain, Touch, or Temperature | Tissue damage, Contact, or Temperature change | Depends on information carried |

<http://neuroscience.uth.tmc.edu/s2/chapter02.html>

72. Receptors, receptor potential vs. action potential, receptive field

- ✓ Receptor definition (energy converter)
- ✓ Receptor potential vs. Action potential
 - RP – analogue (amplitude), AP – digital (frequency)
 - RP – various ionic mechanisms, AP - Na-K based
- ✓ Basic attributes of stimulus
 - Modality, localization, intensity, duration
 - The law of specific nerve energies (labeled line coding)
- ✓ Receptive field
- Definition
- Examples of large and small receptive fields, association with resolution
- Lateral inhibition
- Receptor adaptation (tonic and phasic response)
- ✓ Various classifications of receptors
 - Brief overview of the skin receptors

M U N I

M E D