

# LYMPHATIC SYSTEM

Petr Vaňhara  
2023


M U N I  
M E D      Department  
                 of Histology  
                 and Embryology

# LECTURE CONTENT

- Principles of immune response
  - Innate and acquired immunity
  - Humoral and cellular immunity
- Structures essential for functioning of immune system
  - Lymphatic circuitry
  - Lymphatic follicles
  - Lymph nodes
  - Spleen
  - Thymus
  - MALT
- Development of lymphatic system

Useful for understanding

Important for our course





# IMMUNE SYSTEM

## Immunity = self defense


- **Epithelial:** epithelial barriers equipped with antimicrobial substances
- **Innate:** complement, macrophages and neutrophils, natural killers
- **Acquired:** T and B lymphocytes

## Clinical relevance?

- Autoimmune disorders
- Immunodeficiency




# EPITHELIAL IMMUNE RESPONSE



## EPITHELIUM OF RESPIRATORY PASSAGES



# EPITHELIAL INFLAMMATORY RESPONSE



- Chemotaxis and extravasation of leukocytes, mostly neutrophils, monocytes and dendritic cells to the site of inflammation


- Pro-inflammatory cytokines
  - interleukins (e.g. IL-1, IL-8)
  - TNF $\alpha$ , TGF $\beta$
  - interferons
- Other signaling molecules
  - prostaglandins
  - GM-CSF, M-CSF

and many others




# EPITHELIAL IMMUNE RESPONSE


## Damage-associated molecular patterns (DAMPs)



CellCartoons.net





CellCartoons.net




CellCartoons.net

# DENDRITIC CELLS

- „professional“ antigen presentation = activation of immune cells with high efficiency
  - antigen processing – MHC II
  - cytokine production
  - component of monocyte-macrophage system
  - lymphatic organs, epithelia, connective tissue
- 
- **TCR** recognizing antigen presented in **MHC complex** is essential for activation of **T-cells**
  - **Highly regulated mechanism**
  - **MHC restriction**




# DENDRITIC CELLS





# MONOCYTE-MACROPHAGE SYSTEM

- mononuclear phagocytic system, reticuloendothelial system
- originate in bone marrow: monoblasts → monocytes
- after extravasation → macrophages
- irregular surface (hallmark of phagocytosis)
- numerous lysosomes
- Golgi apparatus and rER
- long-living cells (months)
- phagocytosis (large particles)




# MONOCYTE-MACROPHAGE SYSTEM

- monocytes (circulation)
- macrophages (histiocytes) of c.t.
- Kupffer cells (liver)
- osteoclasts (bones)
- microglia (CNS)
- alveolar macrophages (lungs)
- macrophages and dendritic cells (lymphatic organs, epithelia, c.t.)
- Langerhans cells (skin)
- mesangial cells (kidney)



# INNATE AND ACQUIRED IMMUNITY




# LYMPHOCYTES

## Lymphocytes

CD8 T Cell



B Cell



Regulatory T Cell




CD4 T Cell




CellCartoons.net


- „Clusters of differentiation“, CD
- Surface molecules constituting immunophenotype
- Molecular signaling regulating immune response
- Clinically relevant in diagnostics and therapy




# ANTIBODY (HUMORAL) RESPONSE



1. Antigen presentation and activation of B-cells




2. Activated B-cells proliferate and expand




3. Most of B-cells differentiate to plasma cells, some will convert to memory cells

# ANTIBODY (HUMORAL) RESPONSE



# ANTIBODIES

- Immunoglobulins
- Large proteins with defined structure capable of binding antigens
- Variable and constant regions
- Fc region bound by Fc receptor on immune cells




# ANTIBODIES

- Genome rearrangements leading to generation of unique antibody transcripts (**>10 billions**)
- V(D)J recombination during development
- Isotype switching


Genes in heavy chain locus




Removal of unwanted D and J gene segment




Recombination of D and J exons – DJ recombination



Removal of unwanted V and D gene segment




Recombination of V and DJ exons – VDJ recombination




Antibody transcript will also include constant domain gene


Genes in heavy chain locus of an IgM expressing B cell



Removal of DNA segment by enzyme activity between switch regions




Non-homologous end joining of DNA at switch regions



Excised DNA segment


# ANTIBODIES

- Five principal classes
  - IgG: most common (>75%), soluble, stable
  - IgA: in exocrine secretions, mucosa
  - IgM: natural immunity, activator of complement
  - IgE: activator of mast cells
  - IgD: B-cells activators




# ANTIBODIES


- IgE: activator of mast cells




# CELLULAR RESPONSE



1. Antigen presentation and activation of T-cells




2. T-cells differentiate to cytotoxic Tc cells, memory t cells or regulatory cells





3. Cytotoxic Tc eliminate abnormal cells


# CELLULAR RESPONSE



Oh nooo!  
I'm shedding virus particles!  
Does that mean I'm infected?  
What will happen to me?



VIRAL PARTICLES





INFECTED CELL

CellCartoons.net

CD8 Cytotoxic T Cell  
a.k.a 'Killer T cell'

# MHC PROTEINS

- Major histocompatibility complex
  - Activation or attenuation of T-cell response
  - Peptide display
- 
- MHC I: all nucleated cells and platelets
  - MHC II: antigen presenting cells
  - (MHC III: structurally similar to MHC I and II, but with rather unknown function in immune response)




# B- AND T- LYMPHOCYTES IN HUMAN BODY

## T-lymphocytes

- paracortical zone of lymph nodes
- white pulp of spleen (periarteriolar lymphatic sheath, PALS)
- interfollicular regions in other lymphatic organs (tonsils)

## B-lymphocytes


- lymphatic follicles and medullary cords of lymph nodes
- spleen follicles and marginal zone of white pulp
- lymphatic follicles in other organs



# ACQUIRED IMMUNITY

## Histology:

- **Leukocytes**
  - Lymphocytes
  - Antigen presenting cells
- **Lymphatic organs**



# LYMPHATIC ORGANS

## Development of lymphocytes and APC:

### Primary lymphatic organs

- bone marrow
- thymus

### Secondary lymphatic organs


- lymph nodes
- spleen
- MALT including tonsils and appendix

### Tissues

- blood
- lymph
- epithelia
- connective tissues

# LYMPHATIC CIRCULATION

## Lymph vessels



## Function



- Collect interstitial fluid
- Microenvironment for lymphocyte development and maturation
- Lipid transport (chylomicrons)

## Histology

- Lymph capillaries
  - Thin walled, blunt ended vessels with irregular lamina basalis
  - Anchoring filaments, tiny valves
- Lymph vessels
  - T. intima – endothelium and subendothelial c.t.
  - T. media - few layers of smooth muscle cells
  - T. adventitia - collagen c.t.
  - Similar to small veins
  - Valves derived from t. intima
  - Open to d. thoracicus and d. lymphaticus dx. → v. subclavia (at v. jugularis int.)


# LYMPHATIC CIRCULATION

## Lymph capillaries



# LYMPHATIC CIRCULATION

## Lymph flow is unidirectional



The Journal of  
**Physiology**

<https://doi.org/10.1113/JP272088>

## Lymph composition

- Contains similar concentration of ions to plasma, but lower levels of proteins
- Lipid-rich lymph from intestine - chylus
- Immune cells
- Volume in the circulation ca. 1L (2-2.5L new lymph from interstitial fluid per day)


# LYMPHATIC ORGANS

## central:

- thymus
- bone marrow

## peripheral:



- encapsulated – lymph node, spleen
- mucosa associated lymphoid tissue – MALT
  - tonsils (partially encapsulated)
  - lymphatic follicles in mucosa of hollow organs





# LYMPHATIC FOLLICLE

## Folliculus, nodulus lymphaticus


- non-encapsulated aggregates of reticular connective tissue and lymphocytes
- peripheral lymphatic organs
- mucosa of hollow organs (GIT, respiratory, urinary, reproductive system)
- **primary** – prior any contact with antigen
- **secondary** – stimulated by antigen
  - pale germinative center
  - dark mantle zone




# LYMPHATIC (LYMPHORETICULAR) TISSUE



- reticular connective tissue + lymphocytes



Ross, Romrell





Tanaka, Fujita




Mescher

# LYMPH NODE (nodus lymphaticus, lymphonodus)


- C.t. capsule containing *hilus* with vessels
- parenchyma = lymphoreticular tissue (reticular c.t. and lymphocytes)
  - **cortex** (lymphatic follicles and sinuses) (B-cells)
  - **medulla** (cords and sinuses) (B-cells)
- paracortical region (T-cells)
- **sinuses**: subcapsular (marginal), perifollicular (cortical), medullary
- *Littoral cells* – lining of sinuses, phagocytosis




# LYMPH NODE (NODUS LYMPHATICUS, LYMPHONODUS)



# LYMPH NODE CIRCULATION (BLOOD AND LYMPH)





# **LYMPH NODE (NODUS LYMPHATICUS, LYMPHONODUS)**




# LYMPH NODE (NODUS LYMPHATICUS, LYMPHONODUS)

subcapsular sinus

lymphatic follicle




## LYMPH NODE MEDULLA




### Medullary cords and sinuses

Lymph node from lung hilus with dust (carbon) deposits






# LYMPH NODE MEDULLA



Mescher



## Medullary cords and sinuses

- Plasma cells in medullary cords
- High endothelium post-capillary venules – extravasation of leukocytes from blood to lymph node parenchyma




# SPLEEN (LIEN)


- C.t. capsule and trabecules
- Parenchyma = pulp
- **white** (lymphoid)
  - periarteriolar lymphatic sheath - PALS
  - Malpighian bodies - follicles)
- **red** (non-lymphoid)
  - cords of Billroth
  - venous sinuses
- **marginal zone** between lymphoid and non-lymphoid regions in the spleen



# SPLEEN (LIEN)



# SPLEEN (LIEN)





# SPLEEN (LIEN)



Sobotta



# SPLEEN (LIEN)



mesothelium

capsula lienis


trabecule

# SPLEEN (LIEN)

White pulp




a. centralis




# SPLEEN (LIEN)

PALS – periarteriolar lymphatic sheath – around aa.centrales






# SPLEEN (LIEN)



3



# SPLEEN (LIEN)



Mescher




# SPLEEN (LIEN)

- Venous sinuses of red pulp
- Removal of abnormal erythrocytes



# SPLEEN (LIEN)

- Venous sinuses of red pulp
- Removal of abnormal erythrocytes



# SPLEEN (LIEN)



Venous sinus




Reticular cell



Cells in circulation




# SPLEEN BLOOD CIRCULATION



- a. lienalis
- aa. trabeculares
- aa. centrales
- arteriolae penicillatae
  - (*terminal arteriole of the pulp, sheathed arteriole*)
- venous sinuses
- postcapillary venules
- veins of the pulp
- vv. trabeculares
- v. lienalis


# SPLEEN – OPEN AND CLOSED CIRCULATION

Fawcett, Jensh



# SPLEEN – OPEN AND CLOSED CIRCULATION


## Splenic blood flow



Blood dumps into connective tissue  
Captured back into sinusoids (large ~~venules~~)

# SPLEEN AND LYMPH NODE IN ANTIGEN RECOGNITION

Lymph node







Spleen

Lymph filter


Blood filter

# THYMUS

- c.t. capsule
- parenchyma: cortex and medulla
- epithelial reticulum and T-cells
- Hassal bodies




# THYMUS (YOUNG)



# THYMUS (CORTEX)




## Cortex:

- T-cell proliferation, acquisition of **immunocompetence**
- **positive selection** (functional TCR → survival)
- **hemato-thymic barrier** (endothelium + basal lamina + cell of cytoreticulum)
- prevents premature contact with antigens



# THYMUS (MEDULLA)


- negative selection
- prevention of autoimmune reaction
- overall survival 2-3%
- cytoreticulum
- hemato-thymic barrier absent




Hassal bodies in medulla

# T-CELL SELECTION

- positive: CD4+ CD8+
- tissue-restricted antigens (TRAs)





## THYMUS (INVOLUTION)




# TONSILS


- incomplete encapsulation by connective tissue
  - aggregations of lymphatic tissue (follicles) covered by epithelium of crypt
  - crypts – deep and branched invaginations lined by epithelium
  - reticulated epithelium
- 
- **t. palatina** – stratified squamous e.
  - **t. lingualis** – stratified squamous e.
  - **t. pharyngea** – pseudostratified columnar e.
  - **t. tubaria** – pseudostratified columnar e.




# TONSILLA PALATINA



# TONSILLA PALATINA




# TONSILLA PALATINA




# TONSILLA PALATINA

lymphatic follicle


reticulated epithelium




# TONSILLA LINGUALIS



# TONSILLA LINGUALIS




# TONSILLA LINGUALIS



GII. Weberi



1 mm

# TONSILLA PHARYNGEA



# DEVELOPMENT OF LYMPHATIC SYSTEM



- development starts at week 5
- origin unclear, presumably from mesenchyme or as outgrowths of primitive endothelium



- **Week 6-9**, six primary lymph sacs from local dilatations
  1. **Two** jugular (junction of the subclavian veins with the v. precardinales ( $\rightarrow$  v. jugularis int.)
  2. **Two** iliac lymph sacs near the junction of the iliac veins with the v. postcardinales
  3. **Single** retroperitoneal lymph sac
  4. **Single** cisterna chyli dorsal to the retroperitoneal lymph sac

# DEVELOPMENT OF LYMPHATIC SYSTEM

- Lymph vessels grow from lymph sacs
  - jugular: head, neck, thorax, upper limbs
  - iliac: trunk, lower limbs from iliac
  - retroperitoneal and cisterna chyli: intestine
- Development of lymphatic ducts
  - left and right thoracic duct connecting c. chyli and jugular sacs
  - anastomoses
  - D. thoracicus: caudal part of right thoracic duct, cranial part of left thoracic duct
  - D. lymhaticus dx.: cranial part of right thoracic duct



# DEVELOPMENT OF LYMPHATIC SYSTEM

- **Development of lymph nodes**

- lymph sacs (except for c. chyli) are invaded by mesenchymal cells and constitute apparent clusters of lymph nodes
- B-cell compartments (follicles) develop around birth, lack germinative centers (naive)
- lymph nodes develop along lymph vessels by similar mechanism






FIGURE 55. Development of a lymph node.

# DEVELOPMENT OF TONSILS AND THYMUS


- **Tonsilla palatina**
  - second entodermal pharyngeal pouch (fossa)
- **Tonsilla pharyngea, tubaria and lingualis**
  - aggregation of lymph nodules in the nasopharynx, by opening of tuba auditiva or lingual root
- **Thymus**
  - third entodermal pharyngeal pouch

Pharyngeal apparatus






Tonsils and Throat

## DEVELOPING TONSILS AND THYMUS GLAND



# DEVELOPMENT OF THYMUS

- Thymus
  - third pharyngeal pouch



Bone marrow cells colonize thymic anlage in fetus

# DEVELOPMENT OF THYMUS

Vertebral cartilage

trachea



sternum and ribs

thymus

esophagus




Week 8




# DEVELOPMENT OF SPLEEN

- Dorsal mesentery of stomach
- Mesenchymal origin




# DEVELOPMENT OF SPLEEN

- Dorsal mesentery of stomach
- Mesenchymal origin



# DEVELOPMENT OF SPLEEN



# Thank you for attention

Questions? Comments?

[pvanhara@med.muni.cz](mailto:pvanhara@med.muni.cz)



Special thanks to [CellCartoons.net](http://CellCartoons.net)