

The Emerging Hallmarks of Cancer Metabolism

Natalya N. Pavlova¹ and Craig B. Thompson^{1,*}

¹Cancer Biology and Genetics Program, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

*Correspondence: thompsonc@mskcc.org

<http://dx.doi.org/10.1016/j.cmet.2015.12.006>

Tumorigenesis is dependent on the reprogramming of cellular metabolism as both direct and indirect consequence of oncogenic mutations. A common feature of cancer cell metabolism is the ability to acquire necessary nutrients from a frequently nutrient-poor environment and utilize these nutrients to both maintain viability and build new biomass. The alterations in intracellular and extracellular metabolites that can accompany cancer-associated metabolic reprogramming have profound effects on gene expression, cellular differentiation, and the tumor microenvironment. In this Perspective, we have organized known cancer-associated metabolic changes into six hallmarks: (1) deregulated uptake of glucose and amino acids, (2) use of opportunistic modes of nutrient acquisition, (3) use of glycolysis/TCA cycle intermediates for biosynthesis and NADPH production, (4) increased demand for nitrogen, (5) alterations in metabolite-driven gene regulation, and (6) metabolic interactions with the microenvironment. While few tumors display all six hallmarks, most display several. The specific hallmarks exhibited by an individual tumor may ultimately contribute to better tumor classification and aid in directing treatment.

Introduction

While the first observations on metabolic alterations that are characteristic for tumors were first made nearly a century ago, the field of cancer metabolism has become a topic of a renewed interest in the past decade. Aided by new biochemical and molecular biological tools, studies in cancer cell metabolism have expanded our understanding of the mechanisms and functional consequences of tumor-associated metabolic alterations at various stages of tumorigenesis. In particular, it has become evident that tumorigenesis-associated metabolic alterations encompass all stages of cell-metabolite interaction, (1) affecting the metabolite influx through conferring an increased ability to acquire the necessary nutrients, (2) shaping the way the nutrients are preferentially assigned to metabolic pathways that contribute to cellular tumorigenic properties, and (3) exerting long-ranging effects on cellular fate, among which are alterations in differentiation of cancer cells themselves as well as of the components of the tumor microenvironment (Figure 1). In this Perspective, we take a detailed look at distinct hallmarks of tumorigenesis-associated metabolic reprogramming, and examine the functional contribution of these hallmarks to the establishment and maintenance of the tumorigenic state.

Deregulated Uptake of Glucose and Amino Acids

In order to fulfill the biosynthetic demands associated with proliferation, a cell must increase the import of nutrients from the environment. Two principal nutrients that support survival and biosynthesis in mammalian cells are glucose and glutamine. Through the catabolism of glucose and glutamine, a cell maintains pools of diverse carbon intermediates, which are utilized as building blocks for the assembly of various macromolecules. In addition, controlled oxidation of carbon skeletons of glucose and glutamine allows a cell to capture their reducing power either in the form of NADH and FADH₂, which mediate the transfer of electrons to the electron transport chain to fuel ATP generation, or in the form of a related cofactor NADPH, which provides

reducing power for a wide variety of biosynthetic reactions, as well as helps maintain cellular redox capacity.

A markedly increased consumption of glucose by tumors in comparison to the nonproliferating normal tissues was first described more than 90 years ago by the German physiologist Otto Warburg (Warburg and Negelein, 1924; Warburg et al., 1927). This observation has been confirmed in a variety of tumor contexts and shown to correlate with poor tumor prognosis (Som et al., 1980). Positron emission tomography (PET)-based imaging of the uptake of a radioactive fluorine-labeled glucose analog, ¹⁸F-fluorodeoxyglucose (¹⁸F-FDG) has been successfully used in the clinic for tumor diagnosis and staging, as well as for monitoring responsiveness to treatment (Almuhaideb et al., 2011).

Glutamine, a second principal growth-supporting substrate contributes not only carbon but also reduced nitrogen for the de novo biosynthesis of a number of diverse nitrogen-containing compounds. Thus, glutamine provides the nitrogen required for the biosynthesis of purine and pyrimidine nucleotides, glucosamine-6-phosphate, and nonessential amino acids. Glutamine also has been reported to play a role in the uptake of essential amino acids. While nonessential amino acids can be produced by mammalian cells de novo, essential amino acids must be acquired from external sources. Interestingly, the import of an essential amino acid leucine through the plasma membrane-localized neutral amino acid antiporter LAT1 was shown to be coupled to a simultaneous efflux of glutamine (Nicklin et al., 2009). In such a manner, intracellular glutamine may facilitate the import of a broad range of LAT1 substrates, including leucine, isoleucine, valine, methionine, tyrosine, tryptophan, and phenylalanine (Yanagida et al., 2001).

The high demand of proliferating tumor cells for glutamine was first described in the 1950s by the American physiologist Harry Eagle, who demonstrated that the optimal growth of cultured HeLa cells requires a 10- to 100-fold molar excess of glutamine in culture medium relative to other amino acids (Eagle, 1955).




Figure 1. The Emerging Hallmarks of Cancer Metabolism

Cancer cells accumulate metabolic alterations that allow them to gain access to conventional nutrient sources as well as to unconventional nutrient sources, utilize these nutrients toward the creation of new biomass to sustain deregulated proliferation, and take advantage of the ability of select metabolites to affect the fate of cancer cells themselves as well as a variety of normal cell types within the tumor microenvironment. Three layers of cell-metabolite interaction are depicted, all of which become reprogrammed in cancer. On top are the adaptations that involve nutrient uptake (Hallmarks 1 and 2), followed by alterations to intracellular metabolic pathways (Hallmarks 3 and 4) in the middle. Finally, long-ranging effects of metabolic reprogramming on the cancer cell itself (Hallmark 5), as well as on other cells within its microenvironment (Hallmark 6), are depicted at the bottom.

restored by a combined expression of a plasma membrane glucose transporter GLUT1 and the first enzyme of the glycolytic pathway hexokinase (HK) (Rathmell et al., 2003).

In addition to soluble growth factors, interactions of cells with the extracellular matrix play a role in regulating glucose uptake. Culturing mammary epithelial cells under conditions where the cells

Furthermore, glutamine was found to be the most rapidly consumed amino acid by Ehrlich ascites carcinomas as well as by a number of hepatomas and carcinosarcomas proliferating in vivo (Márquez et al., 1989; Sauer et al., 1982). In fact, numerous tumorigenic contexts are associated with the depletion of glutamine from the tumor environment as compared to the corresponding normal tissue (Márquez et al., 1989; Rivera et al., 1988; Roberts and Frankel, 1949; Yuneva et al., 2012). Following the success of the ^{18}F -FDG imaging paradigm, ^{18}F -labeled glutamine tracers have recently shown promise in preclinical and early clinical studies (Lieberman et al., 2011; Venneti et al., 2015). Use of ^{18}F -labeled glutamine as a tracer appears to provide potentially useful tumor information where the use of ^{18}F -fluorodeoxyglucose is not feasible—for instance, in imaging of tumors that are localized to sites of heavy glucose utilization, such as the brain.

What causes tumor cells to internalize high quantities of glucose and glutamine? Normally, despite being surrounded by nutrient-rich plasma and the extracellular fluid, metazoan cells do not import nutrients in a constitutive manner. On the contrary, nutrient uptake is strictly regulated by growth factor signaling (Thompson, 2011) (Figure 2). For instance, when deprived of growth factors, hematopoietic and neuronal cells fail to consume glucose in quantities sufficient to even maintain cellular bioenergetics (Lindsten et al., 2003; Rathmell et al., 2000). This, in turn, negatively affects cell size, mitochondrial potential and ATP generation—all despite the abundance of glucose in the culture medium (Rathmell et al., 2000). However, survival of growth-factor-deprived cells can be readily

are detached from the extracellular matrix compromises glucose uptake and results in a depressed mitochondrial potential and decreased ATP levels (Grassian et al., 2011; Schafer et al., 2009). Taken together, these observations suggest that the influx of glucose into cells is not driven by the immediate bioenergetic needs of a cell but, on the contrary, is rationed strictly as a result of extracellular stimuli.

In contrast to their normal counterparts, which require adhesion- and growth-factor-driven signaling inputs to maintain survival and proliferation, cancer cells accumulate oncogenic alterations that convey to them a significant degree of independence from these external requirements (Hanahan and Weinberg, 2000; Vogelstein and Kinzler, 2004). In particular, genetic alterations that target PI-3 kinase, its negative regulators PTEN and INPP4B, and activating mutations and gene amplifications in a variety of upstream receptor tyrosine kinases result in constitutive glucose uptake and metabolism in diverse cancer types.

A point of convergence of signals from receptor tyrosine kinases as well as from the extracellular matrix, PI3K/Akt signaling acts as a master regulator of glucose uptake. PI3K/Akt signaling promotes both the expression of glucose transporter GLUT1 mRNA and the translocation of GLUT1 protein from the endomembranes to the cell surface (Barthel et al., 1999; Wieman et al., 2007). In addition, Akt potentiates the activity of the HK, which phosphorylates glucose molecules, thus preventing their efflux back into the extracellular space, as well as of the phosphofructokinase (PFK) enzyme, which catalyzes the key irreversible step of glycolysis (Deprez et al., 1997; Gottlob et al., 2001). In fact, the ^{18}F -FDG-PET signal intensity in tumors correlates




Figure 2. Deregulated Uptake of Glucose and Amino Acids

Aberrantly activated oncogenes and loss of tumor suppressors deregulate the import of glucose and amino acids into cancer cells. Solid arrows depict the movement of metabolites or proteins and metabolic reactions. Dashed arrows depict positive and negative regulatory effects of signal transduction components. RTK, receptor tyrosine kinase; GLUT1, glucose transporter 1; ASCT2/SN2, glutamine transporter; LAT1, neutral amino acid transporter; EAA, essential amino acids; GLS1, glutaminase 1; PRPS2, phosphoribosyl pyrophosphate synthetase 2; CAD, carbamoyl-phosphate synthetase 2; HK, hexokinase; ECM, extracellular matrix.

promotes TCA cycle anaplerosis and stimulates uptake of cysteine by acting as an exchange substrate for the cysteine antiporter, xCT (Conrad and Sato, 2012).

closely with the level of PI3K/Akt pathway activity, and is attenuated by PI-3 kinase and receptor tyrosine kinase inhibitors (Benz et al., 2011; Lheureux et al., 2013; Ma et al., 2009). Furthermore, exogenous expression of a constitutively active form of Akt alone can stimulate glycolysis and is sufficient to restore cell size, viability, mitochondrial potential, and ATP levels in growth-factor-deprived cells in a manner that is dependent upon the presence of glucose in the culture medium (Plas et al., 2001; Rathmell et al., 2003). The constitutively active form of Akt also prevents a decrease in ATP levels triggered by the loss of cellular attachment (Schafer et al., 2009). In addition, Akt signaling is central to facilitating glucose uptake in physiological settings of increased biosynthetic demand. For instance, targeted deletion of Akt1 in the mouse mammary gland abrogates a lactation-induced increase in glucose uptake, resulting in insufficient milk production (Boxer et al., 2006).

However, the PI3K/Akt signaling module is not the only oncogenic stimulus that plays a role in facilitating glucose uptake. Other oncogenic signaling proteins—notably, Ras—have been found to upregulate *GLUT1* mRNA expression and increase cellular glucose consumption (Murakami et al., 1992). Taken together, multiple growth signaling nodes that become aberrantly activated in cancer share an ability to facilitate cellular access to glucose, a principal metabolic substrate.

The signaling pathways that regulate glutamine uptake are still being elucidated. The transcription factor *c-myc*, which is upregulated in proliferating cells and is frequently targeted by amplification in various tumor types, is a principal driver of glutamine utilization by proliferating cells (Wang et al., 2011b; Wise et al., 2008). To this end, *c-myc* induces the transcription of glutamine transporters ASCT2 and SN2, and in addition promotes expression of glutamine-utilizing enzymes glutaminase (GLS1), phosphoribosyl pyrophosphate synthetase (PRPS2), and carbamoyl-phosphate synthetase 2 (CAD), which support transporter-facilitated glutamine uptake by converting glutamine to glutamate (Eberhardy and Farnham, 2001; Gao et al., 2009; Mannava et al., 2008). The resulting glutamate cannot exit the cell through glutamine transport, and as it accumulates,

In addition to a positive regulatory input from *c-myc*, glutamine uptake is also subject to negative regulation by the *Rb* tumor suppressor family of proteins. The deletion of *Rb* family proteins has been shown to upregulate the uptake and utilization of glutamine via the E2F-dependent upregulation of ASCT2 and GLS1 (Reynolds et al., 2014). Taken together, *c-myc* and E2F, two critical facilitators of cell division, exert their effects, in part, via enabling the access of cells to glutamine—a metabolic substrate critical for the biosynthetic demands that accompany DNA replication.

For any given cell, an attempt to proliferate in the absence of proper metabolic resources could end in a disaster. To avoid this, aberrantly activated oncogenes and/or loss of tumor suppressors lock cancer cells in a state where they constitutively scavenge available glucose, glutamine, and essential amino acids from the extracellular environment, which, in turn, enables their uncontrolled proliferation.

Use of Opportunistic Modes of Nutrient Acquisition

Despite the avidity with which they take up glucose and amino acids, cancer cells *in vivo* often encounter conditions of nutrient scarcity as a result of the increased rates of nutrient consumption and the inadequacies of the tumor vascular supply (Vaupel et al., 1989). To deal with depleted supplies of the normal anabolic precursors, certain cancers acquire mutations that activate the cellular ability to utilize alternative ways of obtaining necessary nutrients. For instance, plasma and interstitial fluid of tissues are rich in soluble protein, yet extracellular protein is not generally utilized as a source of amino acids. Remarkably, expression of mutant Ras or *c-Src* alleles provides cells with a mechanism that allows them to recover free amino acids through the lysosomal degradation of extracellular proteins (Commissio et al., 2013) (Figure 3A). In contrast to the uptake of low-molecular-weight nutrients, which enter the cell through dedicated transporters, capture of extracellular macromolecules occurs through macropinocytosis, a process in which bulk extracellular fluid is taken up into giant vesicles (Kerr and Teasdale, 2009). Macropinocytosis is stimulated by Ras- and *c-Src*-driven actin




Figure 3. Use of Opportunistic Modes of Nutrient Acquisition

(A–C) When free amino acids are unavailable, cancer cells may recover amino acids from (A) extracellular proteins via macropinosytosis, (B) entosis of living cells, or (C) phagocytosis of apoptotic bodies.

(D) In conditions of oxygen deficiency, which prevents the *de novo* desaturation of stearate (C18:0) into oleate (C18:1), monounsaturated fatty acids such as oleate (C18:1) can be recovered from extracellular lysophospholipids (LPL). MPS, macropinosome; Lys, lysosome; MPC, macropinosytosis. X represents a lipid head group.

cytoskeleton remodeling. Fluid-filled macropinosomes are trafficked into the interior of the cell, where they fuse with lysosomes and the engulfed proteins are subjected to proteolytic degradation, liberating free amino acids. Indeed, supplementation of culture medium with physiological levels (20–30 mg/mL) of serum albumin enables proliferation of KRAS^{G12D}-transformed MEFs in the absence of an essential amino acid leucine (Palm et al., 2015). Albumin supplementation has even been shown to restore the proliferation of a KRAS^{G12D}-driven mouse pancreatic cancer line in the culture medium that was devoid of all free amino acids (Kamphorst et al., 2015). In contrast to the effect of KRAS^{G12D}, PI3K/Akt signaling does not promote cellular utilization of extracellular protein, despite the ability of this pathway to stimulate the uptake of low-molecular-weight nutrients (Palm et al., 2015).

Remarkably, mTORC1 inhibitors greatly increase the cellular ability to recover amino acids from captured extracellular protein and enhance their growth in the absence of essential amino

acids (Palm et al., 2015). Thus, the utilization of extracellular protein as nutrients is repressed by mTORC1 under amino-acid-replete conditions and is only utilized as an emergency source in conditions when free amino acids are insufficient. A spatial difference in the effect of rapamycin treatment in a KRAS^{G12D}-driven mouse model of pancreatic cancer provides a striking illustration of this paradigm. While rapamycin suppresses cancer cell proliferation at the tumor margin where the vascular delivery of nutrients is effective, rapamycin enhances cell proliferation within the poorly vascularized tumor core by promoting enhanced lysosomal degradation of extracellular proteins (Palm et al., 2015).

In addition to soluble extracellular proteins, free amino acids can be recovered from the engulfment and digestion of entire living cells via a process known as entosis (Figure 3B), as well as from phagocytosis of apoptotic cellular corpses (Figure 3C) (Krajcovic et al., 2013; Stolzing and Grune, 2004). In a manner

similar to macropinocytosis, cargo engulfed via entosis or phagocytosis becomes digested within lysosomes, providing a supply of amino acids to support cell survival and proliferation during conditions of amino acid deficit (Krajcovic et al., 2013). Interestingly, cells that harbor a mutant KRAS allele are more likely to perpetrate entosis than to be consumed in this process (Sun et al., 2014). Thus, KRAS mutant cells may not only have a nutritional advantage over their non-mutant neighbors but also actively eliminate the latter. Such a property may contribute to cell-cell competition within tumors, leading to the emergence of more aggressive cell populations.

A second consequence of a lack of vascular nutrient delivery in growing tumors is an emergence of hypoxic areas, which leads to a suppression of a number of biosynthetic reactions that require molecular oxygen as an electron acceptor. For example, hypoxia compromises stearoyl-CoA desaturase 1 (SCD1)-catalyzed introduction of double bonds into the de novo produced fatty acids, creating a deficit of unsaturated fatty acid species. To supplement for the missing fatty acids, hypoxic cells import “ready-made” unsaturated fatty acids from the surrounding environment in a form of single acyl chain-containing lysophospholipids (Kamphorst et al., 2013) (Figure 3D). Accordingly, removal of serum lipids triggers ER stress and apoptosis in certain cancer cells, when cultured under hypoxic conditions (Young et al., 2013).

In other cancer cells, the extracellular liberation of free fatty acids from more complex lipid species is increased. An elevated expression of lipoprotein lipase and monoacylglycerol lipase (MAGL) is detected across diverse cancer subtypes and correlates with invasiveness (Kuemmerle et al., 2011; Nomura et al., 2010). Finally, while some tumors develop adaptations that allow them to recover fatty acids directly from plasma, others induce the release of stored lipids from the neighboring normal cells. For instance, upregulation of a long chain fatty acid-binding protein FABP4 on the surface of metastatic ovarian cancer cells allows them to acquire fatty acids directly from the adipocytes of the omental fat (Nieman et al., 2011).

Taken together, tumor cells utilize opportunistic modes of nutrient acquisition, which allow them to survive and proliferate in metabolically unfavorable conditions. These adaptations involve an ability to access normally inaccessible nutrient sources, as well as to recover premade molecules when their synthesis within the cell has been compromised.

Even in the absence of available extracellular nutrients, cancer cells have been shown to utilize adaptive metabolic pathways to sustain their viability. In particular, cancer cells can withstand long periods of nutrient deprivation via the self-catabolic process of macroautophagy (Boya et al., 2013). During macroautophagy, intracellular macromolecules and whole organelles are enveloped by double-membrane structures and fuse with lysosomes. Once inside a lysosome, the engulfed cargo is degraded by resident proteases and lipases to liberate free amino and fatty acids. Indeed, autophagy allows nutrient- or growth factor-deprived cells to maintain their viability in culture for weeks (Lum et al., 2005). In addition, deletion of a core component of the autophagy machinery, *Atg7*, was shown to dramatically change the nature of lung tumors driven by *Kras*^{G12D} and *Braf*^{V600E} oncogenes from malignant adenocarcinomas to benign oncocytomas (Guo et al., 2013; Strohecker et al., 2013). However, autophagy

cannot supply cells with new biomass and thus cannot support proliferation in nutrient-poor conditions.

Use of Glycolysis/TCA Cycle Intermediates for Biosynthesis and NADPH Production

Cell proliferation not only increases the amount of nutrients a cell requires as outlined above but also actively changes the way nutrients are used. Indeed, when a tumor cell is quiescent, glucose is preferentially utilized for mitochondrial acetyl-CoA generation, which is then subjected to oxidation in the tricarboxylic (TCA) cycle. The electrons extracted from the oxidative reactions of the TCA cycle are shuttled through NAD⁺/NADH and FAD/FADH₂ to the electron transport chain, creating an electrochemical gradient that fuels ATP production.

The carbon economy of a proliferating cell differs dramatically from that of a cell that is quiescent (Vander Heiden et al., 2009) (Figure 4A). The major use of reduced carbon in proliferating cells is for the biosynthesis of a diverse array of biomolecules—among which are fatty acids and cholesterol, pentose and hexose sugar derivatives, glycerol, nucleotides, and nonessential amino acids. To achieve this, a proliferating cell must first transform acquired nutrients into diverse pools of structural intermediates (Figure 4B). These molecules include cytosolic acetyl-CoA, one-carbon-carrying folate cycle units and S-adenosylmethionine (SAM), as well as a battery of glycolytic and TCA cycle intermediates. In addition, many biosynthetic reactions are reductive by nature and thus require a source of reducing power. For instance, generation of palmitic acid consumes 14 reducing equivalents, while building a cholesterol molecule requires 26 (Lunt and Vander Heiden, 2011). A designated donor of reducing equivalents that is used in cellular biosynthetic reactions is NADPH. Generation of NADPH from NADP⁺ is enabled by the controlled oxidation of carbon substrates in pathways distinct from those that generate NADH to support mitochondrial electron transport. Thus, a proliferating cell must allocate a portion of its carbon substrates to be used in NADPH production.

The reprogramming of carbon metabolism by proliferating cells provides part of the explanation for Warburg's original observations concerning cancer metabolism. Experimenting with tumor slices incubated with glucose ex vivo, Warburg demonstrated that despite consuming massive amounts of glucose and being exposed to ambient oxygen, cancer cells did not take advantage of the bioenergetic benefits offered by the coupling of glycolysis to the TCA cycle. On the contrary, when grown in glucose-rich medium, cells converted the surplus pyruvate to lactate, which, in turn, was secreted into the extracellular environment. Working under assumption that the major use of glucose in cells was to generate ATP, Warburg and many subsequent investigators misinterpreted this phenomenon, attributing it to the “irreversible injuring of respiration,” which leaves tumor cells with a considerably less efficient way of deriving ATP from substrate-level phosphorylation reactions of glycolysis (Warburg, 1956).

However, subsequent investigations into the mitochondrial status of tumor cells have shown that, by and large, tumor cells possess functional mitochondria and retain the ability to conduct oxidative phosphorylation. In fact, targeted depletion of mitochondrial DNA reduces the tumorigenic potential of cancer cell




Figure 4. Use of Glycolysis/TCA Cycle Intermediates for Biosynthesis and NADH Production

(A) Differences in the central carbon metabolism of a cell in a quiescent state compared to a proliferating cell; (B) diverse biosynthetic outputs of central carbon metabolism. RTK, receptor tyrosine kinase; GLUT1, glucose transporter 1; PKM2, pyruvate kinase M2; ACS2, acetyl-CoA synthetase 2; LDH-A, lactate dehydrogenase A; PDH, pyruvate dehydrogenase; PDK1, pyruvate dehydrogenase kinase 1; ACLY, ATP-citrate lyase; MCT1, monocarboxylate transporter 1; ASCT2/SN2, glutamine transporter.

lines *in vitro* and *in vivo* (Cavalli et al., 1997; Morais et al., 1994; Tan et al., 2015). Moreover, preferential conversion of glucose into lactate has been later demonstrated in genetically normal proliferating cells, as well as in cells infected by viruses (Brand et al., 1986; Chambers et al., 2010; Noch and Khalili, 2012). These observations suggest that, rather than being an adaptation to a defect in respiration, the Warburg effect is a regulated metabolic state and may, in fact, be beneficial during a time of increased biosynthetic demand.

Why, then, do proliferating cells convert excess pyruvate to lactate rather than transport the excess pyruvate into the mitochondria to maintain oxidative phosphorylation? The answer is just now coming into focus. Surprisingly, proliferating cells have only a modest increase in their consumption of ATP relative to their need for precursor molecules and reducing equivalents in the form of NADPH. Glucose catabolism is a robust provider of these precursors and reducing equivalents. In contrast, TCA cycle activity which generates NADH and ATP is the major negative regulator of glucose metabolism. By converting excess py-

ruvate to lactate, proliferating cells prevent accumulation of cytosolic NADH and reduce ATP production, promoting continued cytosolic glucose metabolism free from feedback repression by excess mitochondrial ATP generation.

This realization has spurred further exploration of advantages that the decoupling of glycolysis from oxidative phosphorylation might offer to a proliferating cell. Though glycolysis is classically depicted as a single chain of molecular events that leads to the generation of pyruvate, a number of glycolytic intermediates can be diverted into branching pathways, generating diverse biosynthetic precursors. First in the series of these branching pathways is the pentose phosphate pathway (PPP), in which glucose-6-phosphate becomes partially oxidized to generate NADPH and ribose-5-phosphate—a structural component of nucleotides. Utilization of PPP is frequently elevated in tumorigenesis. The key enzymes in the pathway, transketolase-like 1 (TKTL1) and transaldolase (TALDO), are frequently overexpressed in cancer (Wang et al., 2011a; Xu et al., 2009b). Both oncogenes and tumor suppressors have been shown to regulate

PPP activity. For instance, Ras-driven transformation induces transcriptional upregulation of enzymes that mediate ribose-5-phosphate biosynthesis (Ying et al., 2012). Furthermore, wild-type but not the mutant p53 inhibits PPP by directly binding and inactivating the rate-limiting PPP enzyme glucose-6-phosphate dehydrogenase (G6PD) (Jiang et al., 2011).

Fructose-6-phosphate, which follows glucose-6-phosphate in the glycolytic pathway, can also leave glycolysis and become utilized as a substrate for hexosamine biosynthesis. A first committed step of hexosamine biosynthesis is a glutamine fructose-6-phosphate aminotransferase 1 (GFPT1)-catalyzed production of glucosamine-6-phosphate from fructose-6-phosphate and glutamine. Through the generation of N-acetylglucosamine (GlcNAc), a substrate for N- and O-linked glycosylation, the hexosamine pathway provides substrates for cellular glycosylation reactions as well as heparan sulfate and hyaluronic acid biosynthesis, potentiates receptor-mediated signaling, and regulates the stability of select proteins, including *c-myc* (Itkonen et al., 2013; Spiro, 2002; Wellen et al., 2010).

The next glycolytic intermediate that is utilized in macromolecular synthesis is dihydroxyacetone phosphate (DHAP). DHAP is converted by glycerol-3-phosphate dehydrogenase 1 (GPD1) into glycerol-3-phosphate, which is utilized in the biosynthesis of diverse phospholipids, a major structural component of cellular membranes.

Perhaps the most intensely studied growth-promoting mechanism that shunts metabolites out of the glycolytic pathway is the use of 3-phosphoglycerate as a precursor for the biosynthesis of serine, glycine, and as a means to generate methyl donor groups and NADPH. As metabolic flux studies reveal, cancer cells may utilize as much as 50% of glucose-derived carbon in serine biosynthesis and its subsequent catabolism (Locasale et al., 2011). A rate-limiting serine biosynthesis enzyme, 3-phosphoglycerate dehydrogenase (PHGDH), is frequently focally amplified in breast cancer and melanoma and is required for the growth of PHGDH-amplified cells in vitro as well as in vivo (Locasale et al., 2011; Possemato et al., 2011).

Serine has a unique metabolic role in the cell as a major substrate for the so-called one-carbon, or folate, cycle. To this end, its γ -carbon can be transferred to a carrier molecule, tetrahydrofolate (THF), in a reaction catalyzed by serine hydroxymethyltransferase 2 (SHMT2) in the mitochondria, and SHMT1 in the cytosol, generating 5, 10-methylene-THF and glycine. Then 5, 10-methylene-THF undergoes a series of oxidative-reductive transformations, creating a battery of one-carbon-THF species (Tibbetts and Appling, 2010). One-carbon-THF species are utilized as substrates for the biosynthesis of purines, thymidine, as well as in the production of S-adenosylmethionine, a principal substrate for cellular methylation reactions. In addition, stepwise oxidation of one-carbon-THF species was recently shown to produce up to 50% of all cellular NADPH (Fan et al., 2014). Accordingly, hypoxia-driven induction of SHMT2 expression was shown to protect cells from hypoxia-associated oxidative stress (Ye et al., 2014). Thus, one-carbon pathway metabolites contribute to a number of cellular biosynthetic and regulatory processes. Finally, methylene tetrahydrofolate dehydrogenase 2 (MTHFD2), a component of the mitochondrial arm of the one-carbon pathway, was found to be among the top three most frequently overexpressed metabolic enzymes in cancer, sug-

gesting that alterations in the one-carbon pathway may be universally selected for in tumorigenesis (Nilsson et al., 2014).

In conclusion, glycolytic intermediates leave glycolysis to take part in diverse biosynthetic reactions; accordingly, the rate-limiting enzymes within branching pathways of glycolysis are frequently upregulated in tumors. To balance biosynthetic outputs of glycolysis with its role in providing pyruvate that supports TCA cycle activity, proliferating cells have evolved a novel mechanism by which to regulate the last step of glycolysis. This step is regulated by pyruvate kinase (PK), an enzyme that converts phosphoenolpyruvate to the final product of glycolysis, pyruvate. With the exception of liver and kidney, which express tissue-specific PK isoforms PKL and PKR, most tissues express the PKM (muscle) form of PK. PKM exists in two splice variants (Noguchi et al., 1986). While PKM1 is more efficient at producing pyruvate, the majority of proliferating cells and essentially all cancer cells express primarily the PKM2 variant (Christofk et al., 2008a).

Unlike PKM1, PKM2's activity is highly regulated. Thus, PKM2 is inhibited by tyrosine phosphorylation and activated by serine (Chaneton et al., 2012; Christofk et al., 2008b; Ye et al., 2012). In such a manner, growth-factor-dependent signal transduction inhibits PKM2, leading to accumulation of glycolytic intermediates until the growing cell's need for a catabolizable pool of free serine is saturated. In support of this paradigm, a reintroduction of the PKM1 isoform compromises cancer cell proliferation and tumorigenesis in vivo, in part by negatively affecting serine-dependent biosynthesis (Ye et al., 2012). In addition to serine, PKM2 can also be allosterically activated by excess accumulation of other byproducts of glucose metabolism including succinylaminoimidazolecarboxamide (SAICAR), an intermediate product of ribose-5-phosphate biosynthesis (Keller et al., 2012).

The emerging evidence favors the hypothesis that glycolysis is utilized by proliferating cells as a versatile production line that generates metabolic intermediates for numerous biosynthetic processes. Any excess glycolytic flux not utilized for biosynthesis is preferentially converted to lactate to help preserve a sufficient pool of NAD^+ to sustain glycolysis and avoid flooding the mitochondria with a supply of NADH that would suppress the TCA cycle. Excess NADH generated by glycolysis is not imported into mitochondria of proliferating cells solely through chemical shuttles but is also utilized to convert the glycolytic intermediate DHAP into glycerol-3-phosphate that donates electrons directly into the electron transport chain utilizing glycerol-3-phosphate dehydrogenase (mGPDH) localized to the outer mitochondrial membrane. By using FAD as an electron acceptor, the mGPDH-catalyzed reaction reduces net mitochondrial electrical chemical potential from NADH generated via glycolysis. In fact, increased activity of GPDH has been detected in prostate cancer cells, insulinomas, and carcinoid tumors (Chowdhury et al., 2005; MacDonald et al., 1990).

Mitochondria are protected further from excess glycolysis because any pyruvate that does enter the TCA cycle is channeled into citrate that is secreted into the cytosol through the mitochondrial tricarboxylate carrier and broken down into acetyl-CoA and oxaloacetate. Oxaloacetate is converted to malate and reimported into mitochondria to maintain anaplerosis, while the acetyl-CoA serves as a precursor for lipid biosynthesis and protein acetylation. The electrons entering the mitochondrial

electron transport chain through glycerol phosphate dehydrogenase activity and Complex I as a result of mitochondrial conversion of malate to oxaloacetate are sufficient to maintain both mitochondrial integrity and ATP production despite reduced catabolic TCA cycle activity.

A variety of oncogenes have been shown to contribute to the metabolic adaptations of proliferating cells outlined above. For instance, *c-myc* coordinately increases the expression of PDK1; lactate dehydrogenase A (LDH-A), an enzyme which catalyzes the reductive conversion of pyruvate to lactate; and monocarboxylate transporter (MCT1), which facilitates the efflux of lactate into the extracellular space (Wahlström and Henriksen, 2015). In addition to *c-myc*, β -catenin/TCF signaling has also been shown to upregulate MCT1 and PDK1 transcription (Pate et al., 2014). Finally, stabilization of HIF1 α by hypoxia or in various oncogenic contexts also triggers coordinated transcriptional upregulation of LDH-A and PDK1 (Kim et al., 2006; Papandreou et al., 2006).

Even with these adaptations, proliferating cells often accumulate electron transport flux that exceeds the capacity of the ATP synthase, resulting in the formation of excess reactive oxygen species (ROS). In fact, the damaging consequences of the overproduction of ROS from the overloaded electron transport chain may underlie the phenomenon of oncogene-induced cellular senescence (OIS). Triggered by the introduction of a mutant BRAF or RAS alleles, OIS is associated with a profound oxidative damage and irreversible growth arrest (Courtois-Cox et al., 2008). Interestingly, attenuation of the activity of pyruvate dehydrogenase (PDH), a gatekeeper enzyme that converts pyruvate into acetyl-CoA, via the ectopic expression of its negative regulator pyruvate dehydrogenase kinase 1 (PDK1) or an RNAi-mediated depletion of a PDH phosphatase PDP2, was found to bypass mutant BRAF^{V600E}-induced OIS and potentiate tumorigenesis (Kaplon et al., 2013).

Whereas overproduction of ROS is detrimental to cell growth and survival, moderate levels of ROS, which are generated either during respiration or in a targeted manner by a class of NADPH oxidases, constitute an important signaling input that contributes to the maintenance of the tumorigenic state (Sullivan and Chandel, 2014). To this end, ROS act as inhibitors of protein phosphatases such as PTEN and PTP1B, as well as activators of Src family kinases and MAPK (Denu and Tanner, 1998; Kamata et al., 2005; Lee et al., 2002; Yan and Berton, 1996). Furthermore, increased ROS levels facilitate the activation of HIF1 α and NRF2 transcription factors, thus promoting transcriptional programs that further contribute to tumorigenesis (Bertout et al., 2008; Sporn and Liby, 2012).

In contrast to the proliferating bulk of a tumor, accumulating evidence suggests that quiescent subpopulations of tumor cells are markedly less glycolytic and exhibit higher dependence on oxidative phosphorylation, as well as elevated expression of mitochondrial respiratory components. Among such populations are stem-like subclones that emerge following the oncogene ablation in vivo, as well as circulating tumor cells (CTCs) (LeBleu et al., 2014; Viale et al., 2014). Such apparent dichotomy in utilization of carbon for either predominantly biosynthetic or predominantly bioenergetic purposes may explain why tumors retain the capacity for oxidative phosphorylation, despite the original hypothesis put forth by Otto Warburg.

In addition to glycolytic intermediates, growth factor signaling promotes the use of select TCA cycle intermediates in the generation of biosynthetic precursors. In particular, PI3K/Akt activation allows a cell to expand its cytosolic pool of acetyl-CoA, a substrate for the de novo biosynthesis of fatty acids. To this end, Akt activates the ATP-citrate lyase (ACLY) enzyme, which catalyzes cytosol-localized cleavage of the TCA cycle-derived citrate into acetyl-CoA and oxaloacetate (Bauer et al., 2005; Berwick et al., 2002).

While the de novo biosynthesis of fatty acids is low in normal adult tissues—with the exception of lipogenic tissues such as liver, adipose tissue, or mammary epithelium in the period of lactation, tumorigenesis is associated with a dramatic increase in lipid production (Li and Cheng, 2014; Menendez and Lupu, 2007). An increased capacity for producing lipids de novo not only facilitates the formation of lipid bilayers but also enables a cell to alter its membrane composition in favor of oxidative damage-resistant saturated fatty acids as means of adapting to oxidative stress (Rysman et al., 2010). The biosynthesis of fatty acid chains begins with the carboxylation of cytosolic acetyl-CoA by acetyl-CoA carboxylase (ACC) to produce malonyl-CoA, which is further assembled into long fatty acid chains by fatty acid synthase (FASN). In fact, all three major components involved in fatty acyl chain biosynthesis—ACLY, ACC, and FASN—are frequently upregulated in transformed cells; furthermore, their inhibition reduces tumor cell growth in vitro and in vivo (Chajès et al., 2006; Flavin et al., 2010; Migita et al., 2008; Milgraum et al., 1997). In addition to fatty acid production, the de novo biosynthesis of cholesterol from malonyl-CoA plays a role in tumorigenesis as well. Thus, interference with cholesterol biosynthesis by inhibiting a rate-limiting enzyme, hydroxymethylglutaryl coenzyme A reductase (HMGCR), restores the normal acinar morphology of breast cancer cells, suggesting that changes in membrane composition and fluidity may have an effect on tissue architecture and anchorage-independent growth (Freed-Pastor et al., 2012).

In addition to supplying citrate, the TCA cycle also provides metabolic precursors for the biosynthesis of nonessential amino acids including aspartate and asparagine. In fact, aspartate biosynthesis has recently been shown to be critically dependent upon a cell's ability to conduct oxidative phosphorylation (Birsoy et al., 2015; Sullivan et al., 2015). To maintain the ability to synthesize these amino acids, their production must be carefully balanced with an anaplerotic influx into the cycle. The major anaplerotic substrate in growing cells is glutamine. Glutamine can be captured in mitochondria by glutaminase and the resulting glutamate converted into α -ketoglutarate by either glutamate dehydrogenase or amino acid transaminases. Most proliferating cells depend on a continuous supply of glutamine to maintain the integrity of TCA cycle intermediates (DeBerardinis et al., 2007). In *c-myc*-transformed cells in particular, glutamine deprivation leads to the collapse of the TCA cycle and triggers cell death, which can be rescued by the addition of oxaloacetate or a membrane-permeable form of α -ketoglutarate (Wise et al., 2008; Yuneva et al., 2007). Oxidation of glutamine-derived α -ketoglutarate into oxaloacetate does not only help to maintain the ability of cells to synthesize citrate. Thus, the resulting oxaloacetate can be converted to malate, which, in turn, can be further oxidized by the malic enzyme (ME1) to pyruvate in a reaction

that produces NADPH in a manner independent of glucose (Son et al., 2013). Finally, α -ketoglutarate may provide a source of citrate for the de novo lipogenesis in a setting in which hypoxia or certain oncogenic contexts create a deficit of glucose-derived acetyl-CoA. In a process referred to as reductive carboxylation, a section of the TCA cycle reverses itself, converting α -ketoglutarate into citrate, which can be then used to generate a source of cytosolic acetyl-CoA (Metallo et al., 2012; Wise et al., 2011).

Glutamine-derived α -ketoglutarate is not the only anaplerotic substrate that can be exploited by proliferating cells. In fact, tracing the fate of glucose-derived carbons in human patients and in mouse models of glioblastoma and non-small cell lung carcinoma (NSCLC) reveals that at least in these tumorigenic contexts, the principal source for the TCA cycle intermediates is not glutamine, but glucose (Maher et al., 2012; Marin-Valencia et al., 2012; Sellers et al., 2015). Anaplerotic entry of glucose-derived carbons into the TCA cycle takes place via carboxylation of pyruvate into the TCA cycle intermediate oxaloacetate, catalyzed by pyruvate carboxylase (PC). Indeed, survival of glutamine-deprivation-resistant derivatives of a glioblastoma cell line in culture was found to be dependent on pyruvate carboxylase (Cheng et al., 2011). Moreover, PC inhibition in NSCLC-derived cell lines suppresses fatty acid biosynthesis and dramatically reduces proliferation even in the presence of ample glutamine (Sellers et al., 2015). These reports highlight that the cell lineage context, as well as potential secondary effects associated with the adaptation of cells to tissue culture conditions may play an important role in the choice of an anaplerotic substrate.

Finally, extracellular acetate has recently been shown to be utilized by some cancers as a source of acetyl-CoA for biosynthesis. Specifically, a variety of tumor types localized to the brain, including primary glioblastomas as well as brain metastases that originated from other tissues, have been shown to assimilate exogenous acetate and incorporate acetate-derived carbons into de novo synthesized fatty acids (Comerford et al., 2014; Mashimo et al., 2014; Schug et al., 2015). Imported acetate is converted to acetyl-CoA in a reaction catalyzed by acetyl-CoA synthetase 2 (ACSS2), which is a target for amplification in breast cancer. Thus, hypoxia-associated deficit of acetyl-CoA can be augmented not only through the reductive carboxylation of α -ketoglutarate, but also by increasing the utilization of free acetate from plasma and interstitial fluid. Furthermore, upregulation of ACSS2 may potentially allow cells to recycle acetate released from deacetylation of histones and other cellular proteins as well.

Increased Demand for Nitrogen

In addition to increasing the consumption of carbon in biosynthetic pathways, growth signaling concomitantly elevates the cellular demand for reduced nitrogen. Indeed, a proliferating cell must synthesize a number of nitrogen-containing molecules de novo, including nucleotides, nonessential amino acids, and polyamines. As discussed above, the transcriptional effects of *c-myc* and E2F result in enhanced cellular uptake of glutamine. A nonessential amino acid containing two atoms of reduced nitrogen, glutamine serves as the principal way in which reduced nitrogen is transported between cells in metazoan organisms. The amide group of glutamine is an indispensable donor of nitro-

gen for the biosynthesis of purine and pyrimidine bases (Figure 5). Thus, one glutamine molecule is used in the production of uracil and thymine, cytosine and adenine each require two, and building a guanine base costs a cell three molecules of glutamine. In addition, the assembly of both pyrimidine and purine rings utilizes aspartate, which is derived from the transamination of the TCA cycle metabolite oxaloacetate and glutamic acid, both of which are catabolites of glutamine. Thus, glutamine is a key structural building block in the biosynthesis of nucleotides. Accordingly, glutamine levels have been shown to be rate-limiting for cell cycle progression, and glutamine deprivation leads to cell cycle arrest in S phase in certain cellular contexts (Fontenelle and Henderson, 1969; Gaglio et al., 2009).

The oncogene *c-myc* orchestrates nucleotide biosynthesis by upregulating the expression of a number of nucleotide biosynthesis enzymes. Among the *c-myc*-regulated targets are phosphoribosyl pyrophosphate synthetase 2 (PRPS2), which catalyzes a first step of purine biosynthesis, as well as carbamoyl phosphate synthetase II (CAD), which initiates the pyrimidine ring-building cascade (Cunningham et al., 2014; Eberhardy and Farnham, 2001; Mannava et al., 2008). Other targets of *c-myc* that are involved in nucleotide biosynthesis are thymidylate synthase (TS) and inosine monophosphate dehydrogenase 1 (IMPDH1) and 2 (IMPDH2) (Liu et al., 2008; Mannava et al., 2008). Thus, *c-myc* not only facilitates glutamine uptake but also promotes its utilization in the biogenesis of purine and pyrimidine bases. Besides the regulatory inputs from *c-myc*, mutant p53 alleles have also been shown to facilitate expression of nucleotide biosynthesis genes, among which are IMPDH1 and IMPDH2, GMP synthetase (GMPS), and nucleoside salvage enzymes deoxycytidine kinase (DCK) and thymidine kinase (TK1) (Kollareddy et al., 2015). Furthermore, activity of CAD enzyme is regulated via phosphorylation by MAPK and by mTORC1-dependent S6 kinase (Ben-Sahra et al., 2013; Graves et al., 2000; Robitaille et al., 2013). In this manner, mTORC1-driven CAD activation may enable the cell to regulate pyrimidine biogenesis in response to the levels of intracellular glutamine, as the latter contributes to mTORC1 activation (Durán et al., 2012; Jewell et al., 2015).

In addition to being utilized for nucleotide biosynthesis, glutamine can be directly deamidated to glutamate by glutaminase which is frequently upregulated in cancer cells in a *c-myc*-dependent manner. Glutamine-derived glutamate serves as a donor of nitrogen for the production of a number of nonessential amino acids via transamination. In contrast, biosynthesis of asparagine from aspartate, catalyzed by asparagine synthetase (ASNS), utilizes the amide nitrogen of glutamine. Though structurally similar to glutamine, asparagine is the only amino acid that mammalian cells do not catabolize. Yet despite this, asparagine plays a crucial regulatory role in conditions of glutamine deprivation (Zhang et al., 2014). Exactly how asparagine supports cell survival and adaptation to glutamine depletion remains to be fully elucidated. ASNS is frequently upregulated in tumors and is associated with poor prognosis (Sircar et al., 2012; Zhang et al., 2014). In contrast, acute lymphoblastic leukemia cells lack ASNS expression, leading to asparagine auxotrophy. Depletion of plasma asparagine by a recombinant form of bacterial L-Asparaginase is an effective anticancer intervention for this cancer type (Hill et al., 1967; Pieters et al., 2011).




Figure 5. Increased Demand for Nitrogen

Sources of nitrogen atoms and regulation of nucleotide and polyamine biosynthesis. PRPP, phosphoribosyl pyrophosphate; PRPS2, phosphoribosyl pyrophosphate synthetase 2; CAD, carbamoyl-phosphate synthetase 2; Gln, glutamine, Glu, glutamate; Gly, glycine; Asp, aspartate; Fum, fumarate; IMPDH1/2, inosine-5-monophosphate dehydrogenase; TS, thymidylate synthase; GMPS, guanosine monophosphate synthetase; DCK, deoxycytidine kinase; TK, thymidine kinase, ARG1, arginase 1; ODC, ornithine decarboxylase; CAT1/2, cationic amino acid transporter.

Despite the existence of a glutamine biosynthetic pathway in mammalian cells, the majority of proliferating cells in culture require an exogenous supply of glutamine. Interestingly, some cell types, such as embryonic stem cells and luminal breast cancer cells, are capable of proliferating in the absence of glutamine in the culture medium, indicating that in some cellular contexts, glutamine can be produced *de novo* (Carey et al., 2015; Kung et al., 2011). In addition, human glioblastoma samples implanted directly into the brains of immunocompromised mice were found to accumulate glutamine relative to the normal brain tissue (Marin-Valencia et al., 2012); furthermore, a similar accumulation of glucose-derived glutamine within the tumor tissue was observed in a mouse model of c-Met-driven liver cancer (Yuneva et al., 2012). Glutamine synthetase (GS) has been found to be overexpressed in some cancers (Osada et al., 2000), yet its role and its mode of activation remain to be fully understood.

Arginine, despite being a nonessential amino acid, also becomes conditionally essential in some tumorigenic contexts (Figure 5). A carrier of four nitrogen atoms, arginine serves as a precursor for a wide variety of nitrogenous compounds, among which are polyamines, creatine, agmatine, and pyrroline-5-

carboxylate, a precursor for the biosynthesis of proline. The *de novo* biosynthesis of arginine is an integral part of the urea cycle, where it is derived from argininosuccinate in a cleavage reaction catalyzed by argininosuccinate lyase (ASL). Argininosuccinate, in turn, is produced by the enzyme argininosuccinate synthase (ASS1) from citrulline and aspartate. Intriguingly, both ASS1 and ASL enzymes are frequently epigenetically silenced in melanoma, renal cell carcinoma, and hepatocellular carcinoma (Bowles et al., 2008; Ensor et al., 2002; Yoon et al., 2007). Reduced expression of ASS1 and ASL are associated with poor prognosis and resistance to chemotherapy (McAlpine et al., 2014; Nicholson et al., 2009). Accordingly, the auxotrophy of tumors for arginine is being exploited in the development of anticancer therapies (Delage et al., 2010).

Why do tumors opt out of the *de novo* production of arginine and instead begin relying on its exogenous influx? One possibility is that the inactivation of ASL and ASS1 allows cancer cells to accumulate ornithine, which is then utilized in the production of polyamines, a class of nitrogen-containing polycationic aliphatic carbon molecules. Polyamine levels increase in proliferating cells; moreover, polyamines have been shown to inhibit




Figure 6. Alterations in Metabolite-Driven Gene Regulation

Diverse metabolites serve as cofactors or substrates for enzymes involved in deposition and removal of epigenetic marks. HAT, histone acetyltransferase enzymes; Ac, an acetyl mark; SAM, S-adenosylmethionine; SAH, S-adenosylhomocysteine; DNMT, DNA methyltransferase enzymes; HMT, histone methyltransferase enzymes; Me, a methyl mark; LSD1, lysine-specific histone demethylase 1; JHDM, Jumonji domain-containing histone demethylase enzymes; Cyt, cytosine; ^{5me}Cyt, 5-methylcytosine; ^{5hm}Cyt, 5-hydroxymethylcytosine; TET1/2, ten-eleven translocation methylcytosine dioxygenase 1/2; α -KG, α -ketoglutarate; SDH, succinate dehydrogenase; FH, fumarate hydratase; IDH1/2, isocitrate dehydrogenase 1/2.

metabolism to enable increased nutrient acquisition and biosynthesis. However, metabolic networks themselves are not merely passive recipients of growth signals, but quite the contrary, directly transmit the information about the cellular metabolic state to a diverse array of regulatory enzymes, among which are those

apoptosis and promote tumor growth and invasion (Casero and Marton, 2007; Gerner and Meyskens, 2004). Another possible reason for arginine auxotrophy in cancer may be related to the fact that the suppression of ASS1-driven argininosuccinate production leads to an accumulation of its substrate aspartate, which can then be applied toward nucleotide biosynthesis.

In addition to its effects on glutamine metabolism, *c-myc*-orchestrated remodeling of nonessential amino acid metabolism involves the metabolism of proline (Phang et al., 2008). Proline can be produced either from glutamate or from arginine-derived ornithine via a common intermediate, pyrroline-5-carboxylate. The expression of the principal enzyme in proline biosynthesis, pyrroline-5-carboxylate reductase (PYCR1), is upregulated by *c-myc* (Liu et al., 2012). In fact, a meta-analysis of metabolic enzyme expression across diverse tumor types identified PYCR1 as one of the most commonly overexpressed genes in tumors (Nilsson et al., 2014). Reciprocally, proline oxidase (POX), which mediates proline degradation, is negatively controlled by *c-myc* via *miR-23b*, and is a p53 target as well (Donald et al., 2001; Liu et al., 2012). Expression of POX has been shown to inhibit tumor cell growth by triggering cell cycle arrest in the G2 phase (Liu et al., 2009). The contribution of altered proline metabolism to tumorigenesis remains to be elucidated. One explanation suggests that the elevated proline pool may facilitate production of collagen and new extracellular matrix deposition, facilitating tumor invasion. Taken together, while this area of cancer metabolism remains under active investigation, it is clear that, similarly to carbon, the metabolism of nitrogen undergoes complex reprogramming in tumorigenesis.

Alterations in Metabolite-Driven Gene Regulation

Aberrantly activated growth and survival signals that drive tumorigenesis facilitate the reprogramming of cancer cell

that mediate the deposition and removal of epigenetic marks from chromatin (Katada et al., 2012) (Figure 6).

A key metabolite that builds up when cells metabolize more glucose than needed for bioenergetic support is cytosolic acetyl-CoA. Cytosolic acetyl-CoA is the obligate substrate for enzymes that acetylate histones and other proteins. The deposition of acetyl marks on histones is associated with the increased accessibility of the genomic DNA for the assembly of transcriptional complexes, and has a rapid turnover rate. Histone acetylation is exquisitely sensitive to alterations in the cellular nutritional and signaling status (Cai et al., 2011; Shi and Tu, 2013). Indeed, withdrawal and readdition of glucose, as well as activation of oncogenic signaling via introduction of an oncogenic KRAS mutant or a constitutively active form of Akt, increase total histone acetylation, which, in turn, promotes the enhanced and broader gene expression (Lee et al., 2014). Immunohistochemical analysis of glioblastoma and prostate tumor samples has shown that Akt activation levels closely correlate with global histone acetylation status (Lee et al., 2014). Activated Akt expands the extramitochondrial pool of acetyl-CoA by activating ACLY, which converts cytosolic citrate into acetyl-CoA and oxaloacetate. Furthermore, studies have shown that both the glucose- and oncogene-driven increases in global histone acetylation are blocked by ACLY inhibition (Lee et al., 2014; Wellen et al., 2009).

In addition to acetyl-CoA, a p300 histone acetyltransferase has recently been shown to utilize crotonyl-CoA as a substrate as well. Notably, deposition of crotonyl marks on select amino acid residues within histone tails was found to activate gene expression even more potently than acetyl marks (Sabari et al., 2015). Crotonyl-CoA can be produced from the catabolism of lysine and tryptophan, as well as of a short-chain fatty acid butyrate. In fact, a large variety of novel histone marks, such as formylation, propionylation, butyrylation, malonylation, and

succinylation, have recently been identified by tandem mass spectrometry (Chen et al., 2007; Wisniewski et al., 2008; Xie et al., 2012). Further investigation of these marks has potential to dramatically expand the number of metabolic inputs into the regulation of global patterns of gene expression.

Numerous methylation reactions in the cell, including the deposition of methyl marks on histone tails, cytosine methylation on DNA, and adenosine methylation on mRNA utilize S-adenosylmethionine (SAM) as a donor of methyl groups. SAM is a product of the one-carbon metabolic pathway and is fueled by serine catabolism as described above. A number of recent reports demonstrate that histone and DNA methylation is sensitive to alterations in SAM levels (Chiang et al., 2009; Shyh-Chang et al., 2013; Towbin et al., 2012).

The removal of acetyl and methyl marks is also guided by the cellular metabolic state. For instance, sirtuins, a class of deacetylases that catalyze the removal of acetyl marks from histone and nonhistone proteins, utilize NAD⁺ as a cofactor, while FAD serves as a cofactor for a lysine-specific demethylase LSD1 (Imai et al., 2000; Shi et al., 2004). Sensitive to changes in NAD⁺ or FAD availability, these enzymes orchestrate global posttranslational and epigenetic changes that promote energy conservation (Hino et al., 2012; Schwer and Verdin, 2008).

A wide variety of posttranslational modifications in the cell are carried out by the members of a large class of α -ketoglutarate-dependent dioxygenases. Among α -ketoglutarate-dependent dioxygenases are a TET family of DNA demethylases, Jumonji C family of histone demethylases, mRNA demethylases FTO and ALKBH5, and a family of prolyl hydroxylase (PHD) enzymes, which, among other processes, regulate HIF1 α levels in response to oxygen availability and oxidative stress. The reaction mechanism of α -ketoglutarate-dependent dioxygenases involves a concurrent oxidation of a cosubstrate α -ketoglutarate into succinate. Accordingly, intracellular levels of α -ketoglutarate can directly influence the activity of these enzymes. In addition, α -ketoglutarate-dependent dioxygenases are prone to the inhibition by their reaction product, succinate, as well as by fumarate, the downstream product of succinate degradation in the TCA cycle. For instance, a biallelic loss of succinate dehydrogenase (SDH), an alteration found in familial paragangliomas and pheochromocytomas, as well as in a subset of sporadic gastrointestinal stromal tumors, creates a buildup of succinate (Astuti et al., 2001; Janeway et al., 2011). Similarly, loss of a fumarate-metabolizing enzyme fumarate hydratase (FH) and accumulation of fumarate is found in the familial cancer syndrome HLRCC (hereditary leiomyomatosis and renal cell cancer) and in a subset of paragangliomas and pheochromocytomas as well (Tomlinson et al., 2002). Tumors that exhibit SDH and FH loss share a number of phenotypic features that are consistent with dioxygenase inhibition, among which is a characteristic global increase in DNA methylation (Killian et al., 2013; Letouzé et al., 2013; Xiao et al., 2012), as well as elevated levels of HIF1 α (Selak et al., 2005). In fact, the latter, at least in the context of SDH loss, can be reversed by the addition of a membrane-permeable form of α -ketoglutarate (MacKenzie et al., 2007).

Another prominent class of tumor-associated genetic alterations that modulate the activity of α -ketoglutarate-dependent dioxygenases are gain-of-function mutations of isocitrate dehydrogenase 1 (IDH1) and isocitrate dehydrogenase 2 (IDH2). IDH1

and IDH2 mutations have been identified in low-grade glioma, where they represent a likely initiator lesion, as well as in chondrosarcoma, cholangiocarcinoma, and acute myeloid leukemia (AML) (Balss et al., 2008; Borger et al., 2012; Cohen et al., 2013; Mardis et al., 2009; Parsons et al., 2008; Paschka et al., 2010). Mutant alleles of IDH1 and IDH2 exhibit an unusual neomorphic enzymatic function. In contrast to wild-type isocitrate dehydrogenases, which convert the TCA cycle metabolite isocitrate to α -ketoglutarate, a mutant form of IDH prefers to use α -ketoglutarate as a substrate, catalyzing its conversion to the D-enantiomer of 2-hydroxyglutarate (2-HG) (Dang et al., 2010; Ward et al., 2010). Owing to its structural similarity to α -ketoglutarate, 2-HG acts as a competitive inhibitor of α -ketoglutarate-dependent dioxygenases (Xu et al., 2011). Indeed, IDH-driven gliomas, leukemias, and chondrosarcomas display a prominent CpG island hypermethylation, reminiscent of the hypermethylator phenotype seen in SDH- and FH-deficient cancers (Figuera et al., 2010; Lu et al., 2013; Turcan et al., 2012). IDH mutations in AML are mutually exclusive with inactivating mutations in TET2 methylcytosine hydroxylases, further implicating mutant IDH as a bona fide driver of epigenetic remodeling (Figuera et al., 2010). Furthermore, a knockin of a mutant IDH1 allele into normal hematopoietic cells and into chondrocytes in vivo triggers an aberrant expansion of targeted cell lineages (Hirata et al., 2015; Sasaki et al., 2012). In various cellular settings, ectopic expression of a mutant IDH allele or treatment with exogenous D-2-HG is sufficient to induce DNA and histone hypermethylation and block cellular differentiation (Losman et al., 2013; Lu et al., 2012). Conversely, small molecule-mediated inhibition of mutant IDH1 in glioma and IDH2 in leukemia markedly promoted differentiation in these contexts (Rohle et al., 2013; Wang et al., 2013). The abundance of 2-HG has been utilized for ¹H magnetic resonance spectroscopy-based imaging of IDH mutant gliomas (Choi et al., 2012). Moreover, small molecule inhibitors of IDH1 and IDH2 have been reported to induce remissions in phase I clinical trials for treatment of AML patients with the respective mutations.

Notably, some tumor types display elevations in 2-HG even in the absence of IDH mutations. For instance, elevated 2-HG levels have been detected in triple-negative breast cancer (Terunuma et al., 2014). In addition, a specific increase in the L enantiomer of 2-HG, which was shown to exert similar, but not identical, inhibitory effects on α -ketoglutarate-dependent dioxygenases, was found in clear cell renal cell carcinoma, where it correlates with increased methylcytosine levels (Shim et al., 2014). Furthermore, L-2-HG levels were found to increase in response to hypoxia as a result of promiscuous substrate use by LDH-A and malate dehydrogenase (MDH) (Intlekofer et al., 2015), and two recent studies implicate L-2-HG as a metabolic mediator of the cellular stress response to hypoxia (Intlekofer et al., 2015; Oldham et al., 2015).

Metabolic Interactions with the Microenvironment

The information about a metabolic state of a cell affects not only its own long-term decision-making but also has the potential to influence the fate of other cells in its vicinity. Indeed, a variety of genetically stable cell types, among which are tumor-associated fibroblasts, endothelial cells, and innate and adaptive immune system components, are known to undergo characteristic




Figure 7. Metabolic Interactions with the Microenvironment

Cancer cells alter the chemical composition of the extracellular milieu, which exerts pleiotropic effects on the phenotypes of normal cells that reside in the vicinity of the tumor, as well as the extracellular matrix. Reciprocally, the microenvironment affects the metabolism and signaling responses of cancer cells themselves. ECM, extracellular matrix; Treg, regulatory T cells; HA, hyaluronic acid; MMPs, matrix metalloproteinases; MCT1, monocarboxylate transporter 1; CAIX, carbonic anhydrase IX; IDO1, indoleamine-2, 3-dioxygenase 1; TDO2, tryptophan-2, 3-dioxygenase 2; Kyn, kynurenine; AhR, aryl hydrocarbon receptor.

phenotypic changes as a consequence of residing in the vicinity of a growing tumor (Hanahan and Coussens, 2012). How cancer cells reprogram their microenvironment to assist tumor growth and dissemination is an area of intense investigation, but it is clear that such reprogramming encompasses multiple strategies, among which are secreted growth factors and alterations to the extracellular matrix and cell-cell interactions.

In addition to these inputs, proliferating cancer cells alter the metabolic composition of the extracellular milieu around them as well (Figure 7). The high utilization of extracellular glucose and glutamine by cancer cells results in the accumulation of extracellular lactate, which was shown to affect a number of cell types within the tumor microenvironment. Increased lactate levels promote the emergence of an immune-permissive microenvironment by attenuating dendritic and T cell activation and monocyte migration (Fischer et al., 2007; Goetze et al., 2011; Gottfried et al., 2006). In addition, lactate stimulates the polariza-

tion of resident macrophages to a so-called M2 state, which plays a role in immunosuppression and wound healing (Carmona-Fontaine et al., 2013; Colegio et al., 2014). Furthermore, lactate accumulation is instrumental for the promotion of angiogenesis. Thus, lactate promotes the stabilization of HIF1 α and activates NF- κ B and PI-3 kinase signaling in endothelial cells, as well as induces secretion of a proangiogenic factor VEGF from tumor-associated stromal cells (Constant et al., 2000; Schmid et al., 2007; Sonveaux et al., 2012; Végran et al., 2011). Increased levels of lactate also stimulate hyaluronic acid production by fibroblasts, which may contribute to tumor invasiveness (Stern et al., 2002).

The secretion of lactate into the extracellular space via the monocarboxylate transporter MCT1 is coupled to the cotransport of H⁺, which causes acidification of the cellular microenvironment. In addition, the surplus of CO₂ generated in mitochondrial decarboxylation reactions contributes to the extracellular

acidification as well. To this end, CO₂ diffuses into the extracellular space, where it becomes converted to H⁺ and HCO₃⁻ by a class of extracellular carbonic anhydrases (Swietach et al., 2007). Expression of carbonic anhydrases, in particular the CAIX isoform, is elevated during hypoxia (Svastová et al., 2004). Increased extracellular acidification stimulates the proteolytic activity of matrix metalloproteinases (MMPs) and cathepsins, promoting the degradation of the extracellular matrix components and enhancing tumor invasion (Martínez-Zaguilán et al., 1996; Rothberg et al., 2013).

Whereas lactate accumulation and concomitant acidification of the extracellular space may be seen as collateral effects of cancer-specific metabolic reprogramming, some tumors employ a distinct strategy as means of promoting the emergence of an immune-permissive microenvironment around them. Specifically, numerous solid tumor types overexpress tryptophan-degrading dioxygenases indoleamine-2, 3-dioxygenase (IDO1) and tryptophan-2, 3-dioxygenase (TDO2), which catalyze the conversion of an essential amino acid, tryptophan, into its derivative, kynurenine (Munn and Mellor, 2007). As a consequence, tryptophan depletion triggers amino acid deprivation-associated apoptosis of effector T cells (Fallarino et al., 2002). Furthermore, accumulated kynurenine acts as a ligand for aryl hydrocarbon receptor (AhR) (Opitz et al., 2011). In a manner that is dependent on AhR, kynurenine promotes regulatory T cell phenotype, further contributing to the suppression of antitumor immune responses (Fallarino et al., 2006). Finally, kynurenine potentiates autocrine signaling through AhR on cancer cells themselves, promoting the degradation of the extracellular matrix and invasion (Opitz et al., 2011). Small molecule inhibitors of IDO1 are currently being tested in clinical trials (Vacchelli et al., 2014).

Reciprocally, the conditions within the tumor microenvironment have profound effects on the metabolism of a cancer cell. As discussed earlier, tumors are often faced with nutrient- and oxygen-poor surroundings and develop various nutrient-scavenging strategies to bypass these limitations. Furthermore, hypoxia impedes the ability of cells to carry out oxidative phosphorylation and other reactions that require oxygen, and disrupts the redox balance, affecting cellular signaling and transcriptional programs. Taken together, reciprocal interactions between cancer cells and their microenvironment impose a selective pressure that further shapes cancer cell metabolism and actively contributes to the emergence of a more aggressive state.

Remaining Questions

Oncogene-driven metabolic reprogramming allows cancer cells to maintain deregulated proliferation, withstand metabolic challenges that are associated with oxygen and nutrient limitations, maintain a dedifferentiated state through alterations in global patterns of gene expression, and corrupt the surrounding microenvironment into actively assisting tumor growth and dissemination. While most of the studies to this point have been focused on alterations in the metabolism of glucose and glutamine, cancer cells utilize a great variety of other nutrients, among which are sulfur-containing amino acids cysteine and methionine, essential fatty acids, choline, trace metals, and vitamins. We are only beginning to understand the extent to which these nutrients contribute to tumorigenesis.

For instance, multiple lines of evidence indicate that the metabolism of sulfur may undergo reprogramming in cancer. Thus, cancer cells upregulate the uptake of cysteine via the transcriptional upregulation of the xCT transporter, as well as secondary to the increased rate of glutamine utilization by cells (Lo et al., 2008). In fact, no less than 30% of imported glutamine was shown to leave the cell through the xCT transporter as glutamate, while being exchanged for cysteine (Collins et al., 1998). Consistent with its heavy utilization by cancer cells, cysteine was found to be the second most depleted amino acid in pancreatic tumors in comparison to normal pancreatic tissue (Kamphorst et al., 2015). Cysteine has several metabolic fates in proliferating cells, among which are the biosynthesis of glutathione and iron-sulfur clusters, as well as of hydrogen sulfide (H₂S), a gasotransmitter with complex and not yet fully defined functions in cellular physiology, among which are protection from oxidative stress, increased mitochondrial respiration, protection from apoptosis, and facilitation of angiogenesis (Wu et al., 2015).

Alterations in serine metabolism of tumors have received considerable attention in the past 4 years. Notably, in addition to the de novo-produced serine, the inputs from the exogenous one-carbon donors may be affecting one-carbon metabolism and SAM levels as well. Indeed, dietary serine and glycine deficiency has been shown to impair xenograft growth of colon cancer cells in mice (Maddocks et al., 2013). Two other dietary compounds that can be utilized as donors of methyl groups for SAM production are choline and its derivative betaine. Increased choline uptake is observed in breast and prostate cancer; in fact, radioactive ¹⁸F- and ¹¹C-choline is used in the clinic for PET-based tumor imaging (Krause et al., 2013; Kwee et al., 2007). High dietary intake of choline has been linked to a significantly increased risk of lethal prostate cancer (Richman et al., 2012). In contrast, a diet high in choline and betaine was associated with a decreased breast and lung cancer risk (Xu et al., 2009a; Ying et al., 2013; Zhang et al., 2013). Furthermore, diets deficient in choline and betaine were shown to directly affect histone and DNA methylation in a number of animal models (Du et al., 2009; Mehedint et al., 2010; Niculescu et al., 2006), suggesting that alterations in exogenous methyl donor availability may have strong effects on cellular epigenetic state and thus contribute to tumorigenesis.

The effect of various micronutrients—namely, vitamins and trace metals—on tumor growth is a promising area of research. Micronutrients are utilized as cofactors for various enzymes and play complex and not yet fully defined roles in regulating cellular metabolic circuits, as well as signal transduction. For instance, ascorbic acid acts as a cofactor for the α -ketoglutarate-dependent dioxygenases. Notably, ascorbic acid treatment was shown to affect DNA methylation in embryonic stem cells and mouse embryonic fibroblasts in a manner that is dependent on TET2 (Blaschke et al., 2013; Minor et al., 2013). As many cancers are characterized by increased methylation of CpG islands, alterations in ascorbic acid import may thus affect their epigenetic state as well. Furthermore, levels of select trace metals, such as zinc and copper, are altered in cancer cells compared to the normal tissue. Copper levels in particular can be elevated as much as 2- to 3-fold in breast and ovarian tumors, as well as in leukemia (Gupte and Mumper, 2009). Interestingly, copper

was recently shown to be required for BRAF^{V600E}-driven ERK activation in melanoma cells, and the effect of copper depletion could be rescued by the constitutive activation of ERK (Brady et al., 2014).

Finally, the contribution of a broad spectrum of metabolites produced by the body's "forgotten organ," i.e., microbiota, to tumor initiation and progression is only beginning to be elucidated (Garrett, 2015). Taken together, the concept of metabolic alterations of cancer cells, first called to attention nearly a century ago, continues to uncover new connections between nutrient utilization and tumorigenic state. Furthermore, the study of cancer metabolism is not only shedding light on carcinogenesis but is also revealing new principles of how the biochemistry of anabolic metabolism is orchestrated to support normal cell growth, proliferation, and differentiation.

ACKNOWLEDGMENTS

We thank the members of the Thompson laboratory, particularly Dr. Tullia Lindsten, for critical reading of the manuscript, and Leeza Menon for help with manuscript preparation. C.B.T. is supported by grants from the NIH and NCI, and is a founder of Agios Pharmaceuticals and a member of its scientific advisory board. C.B.T. also serves on the board of directors of Merck and Charles River Laboratories. N.N.P. is supported by the Terri Brodeur Breast Cancer Foundation postdoctoral fellowship.

REFERENCES

Almuhaideb, A., Papanthasiou, N., and Bomanji, J. (2011). 18F-FDG PET/CT imaging in oncology. *Ann. Saudi Med.* 31, 3–13.

Astuti, D., Latif, F., Dallol, A., Dahia, P.L., Douglas, F., George, E., Sköldböck, F., Husebye, E.S., Eng, C., and Maher, E.R. (2001). Gene mutations in the succinate dehydrogenase subunit SDHB cause susceptibility to familial pheochromocytoma and to familial paraganglioma. *Am. J. Hum. Genet.* 69, 49–54.

Balss, J., Meyer, J., Mueller, W., Korshunov, A., Hartmann, C., and von Deimling, A. (2008). Analysis of the IDH1 codon 132 mutation in brain tumors. *Acta Neuropathol.* 116, 597–602.

Barthel, A., Okino, S.T., Liao, J., Nakatani, K., Li, J., Whitlock, J.P., Jr., and Roth, R.A. (1999). Regulation of GLUT1 gene transcription by the serine/threonine kinase Akt1. *J. Biol. Chem.* 274, 20281–20286.

Bauer, D.E., Hatzivassiliou, G., Zhao, F., Andreadis, C., and Thompson, C.B. (2005). ATP citrate lyase is an important component of cell growth and transformation. *Oncogene* 24, 6314–6322.

Ben-Sahra, I., Howell, J.J., Asara, J.M., and Manning, B.D. (2013). Stimulation of de novo pyrimidine synthesis by growth signaling through mTOR and S6K1. *Science* 339, 1323–1328.

Benz, M.R., Herrmann, K., Walter, F., Garon, E.B., Reckamp, K.L., Figlin, R., Phelps, M.E., Weber, W.A., Czernin, J., and Allen-Auerbach, M.S. (2011). (18)F-FDG PET/CT for monitoring treatment responses to the epidermal growth factor receptor inhibitor erlotinib. *J. Nucl. Med.* 52, 1684–1689.

Bertout, J.A., Patel, S.A., and Simon, M.C. (2008). The impact of O₂ availability on human cancer. *Nat. Rev. Cancer* 8, 967–975.

Berwick, D.C., Hers, I., Heesom, K.J., Moule, S.K., and Tavaré, J.M. (2002). The identification of ATP-citrate lyase as a protein kinase B (Akt) substrate in primary adipocytes. *J. Biol. Chem.* 277, 33895–33900.

Birsoy, K., Wang, T., Chen, W.W., Freinkman, E., Abu-Remaileh, M., and Sabatini, D.M. (2015). An Essential Role of the Mitochondrial Electron Transport Chain in Cell Proliferation Is to Enable Aspartate Synthesis. *Cell* 162, 540–551.

Blaschke, K., Ebata, K.T., Karimi, M.M., Zepeda-Martínez, J.A., Goyal, P., Mahapatra, S., Tam, A., Laird, D.J., Hirst, M., Rao, A., et al. (2013). Vitamin C induces Tet-dependent DNA demethylation and a blastocyst-like state in ES cells. *Nature* 500, 222–226.

Borger, D.R., Tanabe, K.K., Fan, K.C., Lopez, H.U., Fantin, V.R., Straley, K.S., Schenkein, D.P., Hezel, A.F., Ancukiewicz, M., Liebman, H.M., et al. (2012). Frequent mutation of isocitrate dehydrogenase (IDH)1 and IDH2 in cholangiocarcinoma identified through broad-based tumor genotyping. *Oncologist* 17, 72–79.

Bowles, T.L., Kim, R., Galante, J., Parsons, C.M., Virudachalam, S., Kung, H.J., and Bold, R.J. (2008). Pancreatic cancer cell lines deficient in argininosuccinate synthetase are sensitive to arginine deprivation by arginine deiminase. *Int. J. Cancer* 123, 1950–1955.

Boxer, R.B., Stairs, D.B., Dugan, K.D., Notarfrancesco, K.L., Portocarrero, C.P., Keister, B.A., Belka, G.K., Cho, H., Rathmell, J.C., Thompson, C.B., et al. (2006). Isoform-specific requirement for Akt1 in the developmental regulation of cellular metabolism during lactation. *Cell Metab.* 4, 475–490.

Boya, P., Reggiori, F., and Codogno, P. (2013). Emerging regulation and functions of autophagy. *Nat. Cell Biol.* 15, 713–720.

Brady, D.C., Crowe, M.S., Turski, M.L., Hobbs, G.A., Yao, X., Chaikuad, A., Knapp, S., Xiao, K., Campbell, S.L., Thiele, D.J., and Counter, C.M. (2014). Copper is required for oncogenic BRAF signalling and tumorigenesis. *Nature* 509, 492–496.

Brand, K., Leibold, W., Lippa, P., Schoerner, C., and Schulz, A. (1986). Metabolic alterations associated with proliferation of mitogen-activated lymphocytes and of lymphoblastoid cell lines: evaluation of glucose and glutamine metabolism. *Immunobiology* 173, 23–34.

Cai, L., Sutter, B.M., Li, B., and Tu, B.P. (2011). Acetyl-CoA induces cell growth and proliferation by promoting the acetylation of histones at growth genes. *Mol. Cell* 42, 426–437.

Carey, B.W., Finley, L.W., Cross, J.R., Allis, C.D., and Thompson, C.B. (2015). Intracellular α -ketoglutarate maintains the pluripotency of embryonic stem cells. *Nature* 518, 413–416.

Carmona-Fontaine, C., Bucci, V., Akkari, L., Deforet, M., Joyce, J.A., and Xavier, J.B. (2013). Emergence of spatial structure in the tumor microenvironment due to the Warburg effect. *Proc. Natl. Acad. Sci. USA* 110, 19402–19407.

Casero, R.A., Jr., and Marton, L.J. (2007). Targeting polyamine metabolism and function in cancer and other hyperproliferative diseases. *Nat. Rev. Drug Discov.* 6, 373–390.

Cavalli, L.R., Varella-Garcia, M., and Liang, B.C. (1997). Diminished tumorigenic phenotype after depletion of mitochondrial DNA. *Cell Growth Differ.* 8, 1189–1198.

Chajès, V., Cambot, M., Moreau, K., Lenoir, G.M., and Joulin, V. (2006). Acetyl-CoA carboxylase alpha is essential to breast cancer cell survival. *Cancer Res.* 66, 5287–5294.

Chambers, J.W., Maguire, T.G., and Alwine, J.C. (2010). Glutamine metabolism is essential for human cytomegalovirus infection. *J. Virol.* 84, 1867–1873.

Chaneton, B., Hillmann, P., Zheng, L., Martin, A.C., Maddocks, O.D., Chokkathukalam, A., Coyle, J.E., Jankevics, A., Holding, F.P., Vousden, K.H., et al. (2012). Serine is a natural ligand and allosteric activator of pyruvate kinase M2. *Nature* 491, 458–462.

Chen, Y., Sprung, R., Tang, Y., Ball, H., Sangras, B., Kim, S.C., Falck, J.R., Peng, J., Gu, W., and Zhao, Y. (2007). Lysine propionylation and butyrylation are novel post-translational modifications in histones. *Mol. Cell. Proteomics* 6, 812–819.

Cheng, T., Sudderth, J., Yang, C., Mullen, A.R., Jin, E.S., Matés, J.M., and DeBerardinis, R.J. (2011). Pyruvate carboxylase is required for glutamine-independent growth of tumor cells. *Proc. Natl. Acad. Sci. USA* 108, 8674–8679.

Chiang, E.P., Wang, Y.C., Chen, W.W., and Tang, F.Y. (2009). Effects of insulin and glucose on cellular metabolic fluxes in homocysteine transsulfuration, remethylation, S-adenosylmethionine synthesis, and global deoxyribonucleic acid methylation. *J. Clin. Endocrinol. Metab.* 94, 1017–1025.

Choi, C., Ganji, S.K., DeBerardinis, R.J., Hatanpaa, K.J., Rakheja, D., Kovacs, Z., Yang, X.L., Mashimo, T., Raisanen, J.M., Marin-Valencia, I., et al. (2012). 2-hydroxyglutarate detection by magnetic resonance spectroscopy in IDH-mutated patients with gliomas. *Nat. Med.* 18, 624–629.

Chowdhury, S.K., Gemin, A., and Singh, G. (2005). High activity of mitochondrial glycerophosphate dehydrogenase and glycerophosphate-dependent

- ROS production in prostate cancer cell lines. *Biochem. Biophys. Res. Commun.* 333, 1139–1145.
- Christofk, H.R., Vander Heiden, M.G., Harris, M.H., Ramanathan, A., Gerszten, R.E., Wei, R., Fleming, M.D., Schreiber, S.L., and Cantley, L.C. (2008a). The M2 splice isoform of pyruvate kinase is important for cancer metabolism and tumour growth. *Nature* 452, 230–233.
- Christofk, H.R., Vander Heiden, M.G., Wu, N., Asara, J.M., and Cantley, L.C. (2008b). Pyruvate kinase M2 is a phosphotyrosine-binding protein. *Nature* 452, 181–186.
- Cohen, A.L., Holmen, S.L., and Colman, H. (2013). IDH1 and IDH2 mutations in gliomas. *Curr. Neurol. Neurosci. Rep.* 13, 345.
- Colegio, O.R., Chu, N.Q., Szabo, A.L., Chu, T., Rhebergen, A.M., Jairam, V., Cyrus, N., Brokowski, C.E., Eisenbarth, S.C., Phillips, G.M., et al. (2014). Functional polarization of tumour-associated macrophages by tumour-derived lactic acid. *Nature* 513, 559–563.
- Collins, C.L., Wasa, M., Souba, W.W., and Abcouwer, S.F. (1998). Determinants of glutamine dependence and utilization by normal and tumor-derived breast cell lines. *J. Cell. Physiol.* 176, 166–178.
- Comerford, S.A., Huang, Z., Du, X., Wang, Y., Cai, L., Witkiewicz, A.K., Walters, H., Tantawy, M.N., Fu, A., Manning, H.C., et al. (2014). Acetate dependence of tumors. *Cell* 159, 1591–1602.
- Commisso, C., Davidson, S.M., Soydaner-Azeloglu, R.G., Parker, S.J., Kamphorst, J.J., Hackett, S., Grabocka, E., Nofal, M., Drebin, J.A., Thompson, C.B., et al. (2013). Macropinocytosis of protein is an amino acid supply route in Ras-transformed cells. *Nature* 497, 633–637.
- Conrad, M., and Sato, H. (2012). The oxidative stress-inducible cystine/glutamate antiporter, system x (c) (-): cystine supplier and beyond. *Amino Acids* 42, 231–246.
- Constant, J.S., Feng, J.J., Zabel, D.D., Yuan, H., Suh, D.Y., Scheuenstuhl, H., Hunt, T.K., and Hussain, M.Z. (2000). Lactate elicits vascular endothelial growth factor from macrophages: a possible alternative to hypoxia. *Wound Repair Regen.* 8, 353–360.
- Courtois-Cox, S., Jones, S.L., and Cichowski, K. (2008). Many roads lead to oncogene-induced senescence. *Oncogene* 27, 2801–2809.
- Cunningham, J.T., Moreno, M.V., Lodi, A., Ronen, S.M., and Ruggero, D. (2014). Protein and nucleotide biosynthesis are coupled by a single rate-limiting enzyme, PRPS2, to drive cancer. *Cell* 157, 1088–1103.
- Dang, L., White, D.W., Gross, S., Bennett, B.D., Bittinger, M.A., Driggers, E.M., Fantin, V.R., Jang, H.G., Jin, S., Keenan, M.C., et al. (2010). Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature* 465, 966.
- DeBerardinis, R.J., Mancuso, A., Daikhin, E., Nissim, I., Yudkoff, M., Wehri, S., and Thompson, C.B. (2007). Beyond aerobic glycolysis: transformed cells can engage in glutamine metabolism that exceeds the requirement for protein and nucleotide synthesis. *Proc. Natl. Acad. Sci. USA* 104, 19345–19350.
- Delage, B., Fennell, D.A., Nicholson, L., McNeish, I., Lemoine, N.R., Crook, T., and Szlosarek, P.W. (2010). Arginine deprivation and argininosuccinate synthetase expression in the treatment of cancer. *Int. J. Cancer* 126, 2762–2772.
- Denu, J.M., and Tanner, K.G. (1998). Specific and reversible inactivation of protein tyrosine phosphatases by hydrogen peroxide: evidence for a sulfenic acid intermediate and implications for redox regulation. *Biochemistry* 37, 5633–5642.
- Deprez, J., Vertommen, D., Alessi, D.R., Hue, L., and Rider, M.H. (1997). Phosphorylation and activation of heart 6-phosphofructo-2-kinase by protein kinase B and other protein kinases of the insulin signaling cascades. *J. Biol. Chem.* 272, 17269–17275.
- Donald, S.P., Sun, X.Y., Hu, C.A., Yu, J., Mei, J.M., Valle, D., and Phang, J.M. (2001). Proline oxidase, encoded by p53-induced gene-6, catalyzes the generation of proline-dependent reactive oxygen species. *Cancer Res.* 61, 1810–1815.
- Du, Y.P., Peng, J.S., Sun, A., Tang, Z.H., Ling, W.H., and Zhu, H.L. (2009). Assessment of the effect of betaine on p16 and c-myc DNA methylation and mRNA expression in a chemical induced rat liver cancer model. *BMC Cancer* 9, 261.
- Durán, R.V., Oppliger, W., Robitaille, A.M., Heiserich, L., Skendaj, R., Gottlieb, E., and Hall, M.N. (2012). Glutaminolysis activates Rag-mTORC1 signaling. *Mol. Cell* 47, 349–358.
- Eagle, H. (1955). The minimum vitamin requirements of the L and HeLa cells in tissue culture, the production of specific vitamin deficiencies, and their cure. *J. Exp. Med.* 102, 595–600.
- Eberhardy, S.R., and Farnham, P.J. (2001). c-Myc mediates activation of the cad promoter via a post-RNA polymerase II recruitment mechanism. *J. Biol. Chem.* 276, 48562–48571.
- Ensor, C.M., Holtsberg, F.W., Bomalaski, J.S., and Clark, M.A. (2002). Pegylated arginine deiminase (ADI-SS PEG20,000 mw) inhibits human melanomas and hepatocellular carcinomas in vitro and in vivo. *Cancer Res.* 62, 5443–5450.
- Fallarino, F., Grohmann, U., Vacca, C., Bianchi, R., Orabona, C., Spreca, A., Fioretti, M.C., and Puccetti, P. (2002). T cell apoptosis by tryptophan catabolism. *Cell Death Differ.* 9, 1069–1077.
- Fallarino, F., Grohmann, U., You, S., McGrath, B.C., Cavener, D.R., Vacca, C., Orabona, C., Bianchi, R., Belladonna, M.L., Volpi, C., et al. (2006). The combined effects of tryptophan starvation and tryptophan catabolites down-regulate T cell receptor zeta-chain and induce a regulatory phenotype in naive T cells. *J. Immunol.* 176, 6752–6761.
- Fan, J., Ye, J., Kamphorst, J.J., Shlomi, T., Thompson, C.B., and Rabinowitz, J.D. (2014). Quantitative flux analysis reveals folate-dependent NADPH production. *Nature* 510, 298–302.
- Figuerola, M.E., Abdel-Wahab, O., Lu, C., Ward, P.S., Patel, J., Shih, A., Li, Y., Bhagwat, N., Vasanthakumar, A., Fernandez, H.F., et al. (2010). Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell* 18, 553–567.
- Fischer, K., Hoffmann, P., Voelkl, S., Meidenbauer, N., Ammer, J., Edinger, M., Gottfried, E., Schwarz, S., Rothe, G., Hoves, S., et al. (2007). Inhibitory effect of tumor cell-derived lactic acid on human T cells. *Blood* 109, 3812–3819.
- Flavin, R., Peluso, S., Nguyen, P.L., and Loda, M. (2010). Fatty acid synthase as a potential therapeutic target in cancer. *Future Oncol.* 6, 551–562.
- Fontenelle, L.J., and Henderson, J.F. (1969). Sources of nitrogen as rate-limiting factors for purine biosynthesis de novo in Ehrlich ascites tumor cells. *Biochim. Biophys. Acta* 177, 88–93.
- Freed-Pastor, W.A., Mizuno, H., Zhao, X., Langerød, A., Moon, S.H., Rodriguez-Barrueco, R., Barsotti, A., Chicas, A., Li, W., Polotskaia, A., et al. (2012). Mutant p53 disrupts mammary tissue architecture via the mevalonate pathway. *Cell* 148, 244–258.
- Gaglio, D., Soldati, C., Vanoni, M., Alberghina, L., and Chiaradonna, F. (2009). Glutamine deprivation induces abortive s-phase rescued by deoxyribonucleotides in k-ras transformed fibroblasts. *PLoS ONE* 4, e4715.
- Gao, P., Tchernyshyov, I., Chang, T.C., Lee, Y.S., Kita, K., Ochi, T., Zeller, K.I., De Marzo, A.M., Van Eyk, J.E., Mendell, J.T., and Dang, C.V. (2009). c-Myc suppression of miR-23a/b enhances mitochondrial glutaminase expression and glutamine metabolism. *Nature* 458, 762–765.
- Garrett, W.S. (2015). Cancer and the microbiota. *Science* 348, 80–86.
- Gerner, E.W., and Meyskens, F.L., Jr. (2004). Polyamines and cancer: old molecules, new understanding. *Nat. Rev. Cancer* 4, 781–792.
- Goetze, K., Walenta, S., Ksiazkiewicz, M., Kunz-Schughart, L.A., and Mueller-Klieser, W. (2011). Lactate enhances motility of tumor cells and inhibits monocyte migration and cytokine release. *Int. J. Oncol.* 39, 453–463.
- Gottfried, E., Kunz-Schughart, L.A., Ebner, S., Mueller-Klieser, W., Hoves, S., Andreesen, R., Mackensen, A., and Kreutz, M. (2006). Tumor-derived lactic acid modulates dendritic cell activation and antigen expression. *Blood* 107, 2013–2021.
- Gottlob, K., Majewski, N., Kennedy, S., Kandel, E., Robey, R.B., and Hay, N. (2001). Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. *Genes Dev.* 15, 1406–1418.
- Grassian, A.R., Coloff, J.L., and Brugge, J.S. (2011). Extracellular matrix regulation of metabolism and implications for tumorigenesis. *Cold Spring Harb. Symp. Quant. Biol.* 76, 313–324.

- Graves, L.M., Guy, H.I., Kozlowski, P., Huang, M., Lazarowski, E., Pope, R.M., Collins, M.A., Dahlstrand, E.N., Earp, H.S., 3rd, and Evans, D.R. (2000). Regulation of carbamoyl phosphate synthetase by MAP kinase. *Nature* **403**, 328–332.
- Guo, J.Y., Karsli-Uzunbas, G., Mathew, R., Aisner, S.C., Kamphorst, J.J., Strohecker, A.M., Chen, G., Price, S., Lu, W., Teng, X., et al. (2013). Autophagy suppresses progression of K-ras-induced lung tumors to oncocytomas and maintains lipid homeostasis. *Genes Dev.* **27**, 1447–1461.
- Gupte, A., and Mumper, R.J. (2009). Elevated copper and oxidative stress in cancer cells as a target for cancer treatment. *Cancer Treat. Rev.* **35**, 32–46.
- Hanahan, D., and Coussens, L.M. (2012). Accessories to the crime: functions of cells recruited to the tumor microenvironment. *Cancer Cell* **21**, 309–322.
- Hanahan, D., and Weinberg, R.A. (2000). The hallmarks of cancer. *Cell* **100**, 57–70.
- Hill, J.M., Roberts, J., Loeb, E., Khan, A., MacLellan, A., and Hill, R.W. (1967). L-asparaginase therapy for leukemia and other malignant neoplasms. Remission in human leukemia. *JAMA* **202**, 882–888.
- Hino, S., Sakamoto, A., Nagaoka, K., Anan, K., Wang, Y., Mimasu, S., Ume-hara, T., Yokoyama, S., Kosai, K., and Nakao, M. (2012). FAD-dependent lysine-specific demethylase-1 regulates cellular energy expenditure. *Nat. Commun.* **3**, 758.
- Hirata, M., Sasaki, M., Cairns, R.A., Inoue, S., Puvindran, V., Li, W.Y., Snow, B.E., Jones, L.D., Wei, Q., Sato, S., et al. (2015). Mutant IDH is sufficient to initiate enchondromatosis in mice. *Proc. Natl. Acad. Sci. USA* **112**, 2829–2834.
- Imai, S., Armstrong, C.M., Kaeberlein, M., and Guarente, L. (2000). Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* **403**, 795–800.
- Intlekofer, A.M., Demattee, R.G., Venneti, S., Finley, L.W., Lu, C., Judkins, A.R., Rustenburg, A.S., Grinaway, P.B., Chodera, J.D., Cross, J.R., et al. (2015). Hypoxia induces production of L-2-hydroxyglutarate. *Cell Metab.* **22**, 304–311.
- Itkonen, H.M., Minner, S., Guldvik, I.J., Sandmann, M.J., Tsourlakis, M.C., Berge, V., Svindland, A., Schlomm, T., and Mills, I.G. (2013). O-GlcNAc transferase integrates metabolic pathways to regulate the stability of c-MYC in human prostate cancer cells. *Cancer Res.* **73**, 5277–5287.
- Janeway, K.A., Kim, S.Y., Lodish, M., Nosé, V., Rustin, P., Gaal, J., Dahia, P.L., Liegl, B., Ball, E.R., Raygada, M., et al.; NIH Pediatric and Wild-Type GIST Clinic (2011). Defects in succinate dehydrogenase in gastrointestinal stromal tumors lacking KIT and PDGFRA mutations. *Proc. Natl. Acad. Sci. USA* **108**, 314–318.
- Jewell, J.L., Kim, Y.C., Russell, R.C., Yu, F.X., Park, H.W., Plouffe, S.W., Tagliabracci, V.S., and Guan, K.L. (2015). Metabolism. Differential regulation of mTORC1 by leucine and glutamine. *Science* **347**, 194–198.
- Jiang, P., Du, W., Wang, X., Mancuso, A., Gao, X., Wu, M., and Yang, X. (2011). p53 regulates biosynthesis through direct inactivation of glucose-6-phosphate dehydrogenase. *Nat. Cell Biol.* **13**, 310–316.
- Kamata, H., Honda, S., Maeda, S., Chang, L., Hirata, H., and Karin, M. (2005). Reactive oxygen species promote TNF α -induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* **120**, 649–661.
- Kamphorst, J.J., Cross, J.R., Fan, J., de Stanchina, E., Mathew, R., White, E.P., Thompson, C.B., and Rabinowitz, J.D. (2013). Hypoxic and Ras-transformed cells support growth by scavenging unsaturated fatty acids from lysophospholipids. *Proc. Natl. Acad. Sci. USA* **110**, 8882–8887.
- Kamphorst, J.J., Nofal, M., Commisso, C., Hackett, S.R., Lu, W., Grabocka, E., Vander Heiden, M.G., Miller, G., Drebin, J.A., Bar-Sagi, D., et al. (2015). Human pancreatic cancer tumors are nutrient poor and tumor cells actively scavenge extracellular protein. *Cancer Res.* **75**, 544–553.
- Kaplon, J., Zheng, L., Meissl, K., Chaneton, B., Selivanov, V.A., Mackay, G., van der Burg, S.H., Verdegaaal, E.M., Cascante, M., Shlomi, T., et al. (2013). A key role for mitochondrial gatekeeper pyruvate dehydrogenase in oncogene-induced senescence. *Nature* **498**, 109–112.
- Katada, S., Imhof, A., and Sassone-Corsi, P. (2012). Connecting threads: epigenetics and metabolism. *Cell* **148**, 24–28.
- Keller, K.E., Tan, I.S., and Lee, Y.S. (2012). SAICAR stimulates pyruvate kinase isoform M2 and promotes cancer cell survival in glucose-limited conditions. *Science* **338**, 1069–1072.
- Kerr, M.C., and Teasdale, R.D. (2009). Defining macropinocytosis. *Traffic* **10**, 364–371.
- Killian, J.K., Kim, S.Y., Miettinen, M., Smith, C., Merino, M., Tsokos, M., Quezado, M., Smith, W.I., Jr., Jahromi, M.S., Xekouki, P., et al. (2013). Succinate dehydrogenase mutation underlies global epigenomic divergence in gastrointestinal stromal tumor. *Cancer Discov.* **3**, 648–657.
- Kim, J.W., Tchernyshyov, I., Semenza, G.L., and Dang, C.V. (2006). HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab.* **3**, 177–185.
- Kollareddy, M., Dimitrova, E., Vallabhaneni, K.C., Chan, A., Le, T., Chauhan, K.M., Carrero, Z.I., Ramakrishnan, G., Watabe, K., Haupt, Y., et al. (2015). Regulation of nucleotide metabolism by mutant p53 contributes to its gain-of-function activities. *Nat. Commun.* **6**, 7389.
- Krajcovic, M., Krishna, S., Akkari, L., Joyce, J.A., and Overholtzer, M. (2013). mTOR regulates phagosome and entotic vacuole fission. *Mol. Biol. Cell* **24**, 3736–3745.
- Krause, B.J., Souvatzoglou, M., and Treiber, U. (2013). Imaging of prostate cancer with PET/CT and radioactively labeled choline derivatives. *Urol. Oncol.* **31**, 427–435.
- Kuemmerle, N.B., Rysman, E., Lombardo, P.S., Flanagan, A.J., Lipe, B.C., Wells, W.A., Pettus, J.R., Froehlich, H.M., Memoli, V.A., Morganelli, P.M., et al. (2011). Lipoprotein lipase links dietary fat to solid tumor cell proliferation. *Mol. Cancer Ther.* **10**, 427–436.
- Kung, H.N., Marks, J.R., and Chi, J.T. (2011). Glutamine synthetase is a genetic determinant of cell type-specific glutamine independence in breast epithelia. *PLoS Genet.* **7**, e1002229.
- Kwee, S.A., DeGrado, T.R., Talbot, J.N., Gutman, F., and Coel, M.N. (2007). Cancer imaging with fluorine-18-labeled choline derivatives. *Semin. Nucl. Med.* **37**, 420–428.
- LeBleu, V.S., O’Connell, J.T., Gonzalez Herrera, K.N., Wikman, H., Pantel, K., Haigis, M.C., de Carvalho, F.M., Damascena, A., Domingos Chinen, L.T., Rocha, R.M., et al. (2014). PGC-1 α mediates mitochondrial biogenesis and oxidative phosphorylation in cancer cells to promote metastasis. *Nat. Cell Biol.* **16**, 992–1003, 1001–1015.
- Lee, S.R., Yang, K.S., Kwon, J., Lee, C., Jeong, W., and Rhee, S.G. (2002). Reversible inactivation of the tumor suppressor PTEN by H₂O₂. *J. Biol. Chem.* **277**, 20336–20342.
- Lee, J.V., Carrer, A., Shah, S., Snyder, N.W., Wei, S., Venneti, S., Worth, A.J., Yuan, Z.F., Lim, H.W., Liu, S., et al. (2014). Akt-dependent metabolic reprogramming regulates tumor cell histone acetylation. *Cell Metab.* **20**, 306–319.
- Letouzé, E., Martinelli, C., Lorient, C., Burnichon, N., Abermil, N., Ottolenghi, C., Janin, M., Menara, M., Nguyen, A.T., Benit, P., et al. (2013). SDH mutations establish a hypermethylator phenotype in paraganglioma. *Cancer Cell* **23**, 739–752.
- Lheureux, S., Lecerf, C., Briand, M., Louis, M.H., Dutoit, S., Jebahi, A., Giffard, F., Fournier, C.B., Batalla, A., Poulain, L., and Aide, N. (2013). (18)F-FDG is a surrogate marker of therapy response and tumor recovery after drug withdrawal during treatment with a dual PI3K/mTOR inhibitor in a preclinical model of cisplatin-resistant ovarian cancer. *Transl. Oncol.* **6**, 586–595.
- Li, J., and Cheng, J.X. (2014). Direct visualization of de novo lipogenesis in single living cells. *Sci. Rep.* **4**, 6807.
- Lieberman, B.P., Ploessl, K., Wang, L., Qu, W., Zha, Z., Wise, D.R., Chodosh, L.A., Belka, G., Thompson, C.B., and Kung, H.F. (2011). PET imaging of glutaminolysis in tumors by 18F-(2S,4R)4-fluoroglutamine. *J. Nucl. Med.* **52**, 1947–1955.
- Lindsten, T., Golden, J.A., Zong, W.X., Minarcik, J., Harris, M.H., and Thompson, C.B. (2003). The proapoptotic activities of Bax and Bak limit the size of the neural stem cell pool. *J. Neurosci.* **23**, 11112–11119.
- Liu, Y.C., Li, F., Handler, J., Huang, C.R., Xiang, Y., Neretti, N., Sedivy, J.M., Zeller, K.I., and Dang, C.V. (2008). Global regulation of nucleotide biosynthetic genes by c-Myc. *PLoS ONE* **3**, e2722.

- Liu, Y., Borchert, G.L., Donald, S.P., Diwan, B.A., Anver, M., and Phang, J.M. (2009). Proline oxidase functions as a mitochondrial tumor suppressor in human cancers. *Cancer Res.* *69*, 6414–6422.
- Liu, W., Le, A., Hancock, C., Lane, A.N., Dang, C.V., Fan, T.W., and Phang, J.M. (2012). Reprogramming of proline and glutamine metabolism contributes to the proliferative and metabolic responses regulated by oncogenic transcription factor c-MYC. *Proc. Natl. Acad. Sci. USA* *109*, 8983–8988.
- Lo, M., Wang, Y.Z., and Gout, P.W. (2008). The x(c)-cystine/glutamate antiporter: a potential target for therapy of cancer and other diseases. *J. Cell. Physiol.* *215*, 593–602.
- Locasale, J.W., Grassian, A.R., Melman, T., Lyssiotis, C.A., Mattaini, K.R., Bass, A.J., Heffron, G., Metallo, C.M., Muranen, T., Sharfi, H., et al. (2011). Phosphoglycerate dehydrogenase diverts glycolytic flux and contributes to oncogenesis. *Nat. Genet.* *43*, 869–874.
- Losman, J.A., Looper, R.E., Koivunen, P., Lee, S., Schneider, R.K., McMahon, C., Cowley, G.S., Root, D.E., Ebert, B.L., and Kaelin, W.G., Jr. (2013). (R)-2-hydroxyglutarate is sufficient to promote leukemogenesis and its effects are reversible. *Science* *339*, 1621–1625.
- Lu, C., Ward, P.S., Kapoor, G.S., Rohle, D., Turcan, S., Abdel-Wahab, O., Edwards, C.R., Khanin, R., Figueroa, M.E., Melnick, A., et al. (2012). IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature* *483*, 474–478.
- Lu, C., Venneti, S., Akalin, A., Fang, F., Ward, P.S., Dematteo, R.G., Intlekofer, A.M., Chen, C., Ye, J., Hameed, M., et al. (2013). Induction of sarcomas by mutant IDH2. *Genes Dev.* *27*, 1986–1998.
- Lum, J.J., Bauer, D.E., Kong, M., Harris, M.H., Li, C., Lindsten, T., and Thompson, C.B. (2005). Growth factor regulation of autophagy and cell survival in the absence of apoptosis. *Cell* *120*, 237–248.
- Lunt, S.Y., and Vander Heiden, M.G. (2011). Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annu. Rev. Cell Dev. Biol.* *27*, 441–464.
- Ma, W.W., Jacene, H., Song, D., Vilardell, F., Messersmith, W.A., Laheru, D., Wahl, R., Endres, C., Jimeno, A., Pomper, M.G., and Hidalgo, M. (2009). [18F]fluorodeoxyglucose positron emission tomography correlates with Akt pathway activity but is not predictive of clinical outcome during mTOR inhibitor therapy. *J. Clin. Oncol.* *27*, 2697–2704.
- MacDonald, M.J., Warner, T.F., and Mertz, R.J. (1990). High activity of mitochondrial glycerol phosphate dehydrogenase in insulinomas and carcinoid and other tumors of the amine precursor uptake decarboxylation system. *Cancer Res.* *50*, 7203–7205.
- MacKenzie, E.D., Selak, M.A., Tennant, D.A., Payne, L.J., Crosby, S., Frederiksen, C.M., Watson, D.G., and Gottlieb, E. (2007). Cell-permeating alpha-ketoglutarate derivatives alleviate pseudohypoxia in succinate dehydrogenase-deficient cells. *Mol. Cell. Biol.* *27*, 3282–3289.
- Maddocks, O.D., Berkers, C.R., Mason, S.M., Zheng, L., Blyth, K., Gottlieb, E., and Vousden, K.H. (2013). Serine starvation induces stress and p53-dependent metabolic remodeling in cancer cells. *Nature* *493*, 542–546.
- Maher, E.A., Marin-Valencia, I., Bachoo, R.M., Mashimo, T., Raisanen, J., Hatanpaa, K.J., Jindal, A., Jeffrey, F.M., Choi, C., Madden, C., et al. (2012). Metabolism of [U-13 C]glucose in human brain tumors in vivo. *NMR Biomed.* *25*, 1234–1244.
- Mannava, S., Grachtchouk, V., Wheeler, L.J., Im, M., Zhuang, D., Slavina, E.G., Mathews, C.K., Shewach, D.S., and Nikiforov, M.A. (2008). Direct role of nucleotide metabolism in C-MYC-dependent proliferation of melanoma cells. *Cell Cycle* *7*, 2392–2400.
- Mardis, E.R., Ding, L., Dooling, D.J., Larson, D.E., McLellan, M.D., Chen, K., Koboldt, D.C., Fulton, R.S., Delehaunty, K.D., McGrath, S.D., et al. (2009). Recurring mutations found by sequencing an acute myeloid leukemia genome. *N. Engl. J. Med.* *361*, 1058–1066.
- Marin-Valencia, I., Yang, C., Mashimo, T., Cho, S., Baek, H., Yang, X.L., Rajagopalan, K.N., Maddie, M., Vemireddy, V., Zhao, Z., et al. (2012). Analysis of tumor metabolism reveals mitochondrial glucose oxidation in genetically diverse human glioblastomas in the mouse brain in vivo. *Cell Metab.* *15*, 827–837.
- Márquez, J., Sánchez-Jiménez, F., Medina, M.A., Quesada, A.R., and Núñez de Castro, I. (1989). Nitrogen metabolism in tumor bearing mice. *Arch. Biochem. Biophys.* *268*, 667–675.
- Martínez-Zaguilán, R., Seftor, E.A., Seftor, R.E., Chu, Y.W., Gillies, R.J., and Hendrix, M.J. (1996). Acidic pH enhances the invasive behavior of human melanoma cells. *Clin. Exp. Metastasis* *14*, 176–186.
- Mashimo, T., Pichumani, K., Vemireddy, V., Hatanpaa, K.J., Singh, D.K., Sira-sanagandla, S., Nannepaga, S., Piccirillo, S.G., Kovacs, Z., Foong, C., et al. (2014). Acetate is a bioenergetic substrate for human glioblastoma and brain metastases. *Cell* *159*, 1603–1614.
- McAlpine, J.A., Lu, H.T., Wu, K.C., Knowles, S.K., and Thomson, J.A. (2014). Down-regulation of argininosuccinate synthetase is associated with cisplatin resistance in hepatocellular carcinoma cell lines: implications for PEGylated arginine deiminase combination therapy. *BMC Cancer* *14*, 621.
- Mehedint, M.G., Niculescu, M.D., Craciunescu, C.N., and Zeisel, S.H. (2010). Choline deficiency alters global histone methylation and epigenetic marking at the Re1 site of the calbindin 1 gene. *FASEB J.* *24*, 184–195.
- Menendez, J.A., and Lupu, R. (2007). Fatty acid synthase and the lipogenic phenotype in cancer pathogenesis. *Nat. Rev. Cancer* *7*, 763–777.
- Metallo, C.M., Gameiro, P.A., Bell, E.L., Mattaini, K.R., Yang, J., Hiller, K., Jewell, C.M., Johnson, Z.R., Irvine, D.J., Guarente, L., et al. (2012). Reductive glutamine metabolism by IDH1 mediates lipogenesis under hypoxia. *Nature* *481*, 380–384.
- Migita, T., Narita, T., Nomura, K., Miyagi, E., Inazuka, F., Matsuura, M., Ushijima, M., Mashima, T., Seimiya, H., Satoh, Y., et al. (2008). ATP citrate lyase: activation and therapeutic implications in non-small cell lung cancer. *Cancer Res.* *68*, 8547–8554.
- Milgram, L.Z., Witters, L.A., Pasternack, G.R., and Kuhajda, F.P. (1997). Enzymes of the fatty acid synthesis pathway are highly expressed in situ breast carcinoma. *Clin. Cancer Res.* *3*, 2115–2120.
- Minor, E.A., Court, B.L., Young, J.I., and Wang, G. (2013). Ascorbate induces ten-eleven translocation (Tet) methylcytosine dioxygenase-mediated generation of 5-hydroxymethylcytosine. *J. Biol. Chem.* *288*, 13669–13674.
- Morais, R., Zinkewich-Péotti, K., Parent, M., Wang, H., Babai, F., and Zollinger, M. (1994). Tumor-forming ability in athymic nude mice of human cell lines devoid of mitochondrial DNA. *Cancer Res.* *54*, 3889–3896.
- Munn, D.H., and Mellor, A.L. (2007). Indoleamine 2,3-dioxygenase and tumor-induced tolerance. *J. Clin. Invest.* *117*, 1147–1154.
- Murakami, T., Nishiyama, T., Shirotani, T., Shinohara, Y., Kan, M., Ishii, K., Kanai, F., Nakazuru, S., and Ebina, Y. (1992). Identification of two enhancer elements in the gene encoding the type 1 glucose transporter from the mouse which are responsive to serum, growth factor, and oncogenes. *J. Biol. Chem.* *267*, 9300–9306.
- Nicholson, L.J., Smith, P.R., Hiller, L., Szlosarek, P.W., Kimberley, C., Sehouli, J., Koensgen, D., Mustea, A., Schmid, P., and Crook, T. (2009). Epigenetic silencing of argininosuccinate synthetase confers resistance to platinum-induced cell death but collateral sensitivity to arginine auxotrophy in ovarian cancer. *Int. J. Cancer* *125*, 1454–1463.
- Nicklin, P., Bergman, P., Zhang, B., Triantafellow, E., Wang, H., Nyfeler, B., Yang, H., Hild, M., Kung, C., Wilson, C., et al. (2009). Bidirectional transport of amino acids regulates mTOR and autophagy. *Cell* *136*, 521–534.
- Niculescu, M.D., Craciunescu, C.N., and Zeisel, S.H. (2006). Dietary choline deficiency alters global and gene-specific DNA methylation in the developing hippocampus of mouse fetal brains. *FASEB J.* *20*, 43–49.
- Nieman, K.M., Kenny, H.A., Penicka, C.V., Ladanyi, A., Buell-Gutbrod, R., Zillhardt, M.R., Romero, I.L., Carey, M.S., Mills, G.B., Hotamisligil, G.S., et al. (2011). Adipocytes promote ovarian cancer metastasis and provide energy for rapid tumor growth. *Nat. Med.* *17*, 1498–1503.
- Nilsson, R., Jain, M., Madhusudhan, N., Sheppard, N.G., Strittmatter, L., Kampf, C., Huang, J., Asplund, A., and Mootha, V.K. (2014). Metabolic enzyme expression highlights a key role for MTHFD2 and the mitochondrial folate pathway in cancer. *Nat. Commun.* *5*, 3128.
- Noch, E., and Khalil, K. (2012). Oncogenic viruses and tumor glucose metabolism: like kids in a candy store. *Mol. Cancer Ther.* *11*, 14–23.

- Noguchi, T., Inoue, H., and Tanaka, T. (1986). The M1- and M2-type isozymes of rat pyruvate kinase are produced from the same gene by alternative RNA splicing. *J. Biol. Chem.* **261**, 13807–13812.
- Nomura, D.K., Long, J.Z., Niessen, S., Hoover, H.S., Ng, S.W., and Cravatt, B.F. (2010). Monoacylglycerol lipase regulates a fatty acid network that promotes cancer pathogenesis. *Cell* **140**, 49–61.
- Oldham, W.M., Clish, C.B., Yang, Y., and Loscalzo, J. (2015). Hypoxia-mediated increases in l-2-hydroxyglutarate coordinate the metabolic response to reductive stress. *Cell Metab.* **22**, 291–303.
- Opitz, C.A., Litztenburger, U.M., Sahn, F., Ott, M., Tritschler, I., Trump, S., Schumacher, T., Jestaedt, L., Schrenk, D., Weller, M., et al. (2011). An endogenous tumour-promoting ligand of the human aryl hydrocarbon receptor. *Nature* **478**, 197–203.
- Osada, T., Nagashima, I., Tsuno, N.H., Kitayama, J., and Nagawa, H. (2000). Prognostic significance of glutamine synthetase expression in unifocal advanced hepatocellular carcinoma. *J. Hepatol.* **33**, 247–253.
- Palm, W., Park, Y., Wright, K., Pavlova, N.N., Tuveson, D.A., and Thompson, C.B. (2015). The utilization of extracellular proteins as nutrients is suppressed by mTORC1. *Cell* **162**, 259–270.
- Papandreou, I., Cairns, R.A., Fontana, L., Lim, A.L., and Denko, N.C. (2006). HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab.* **3**, 187–197.
- Parsons, D.W., Jones, S., Zhang, X., Lin, J.C., Leary, R.J., Angenendt, P., Manjoo, P., Carter, H., Siu, I.M., Gallia, G.L., et al. (2008). An integrated genomic analysis of human glioblastoma multiforme. *Science* **321**, 1807–1812.
- Paschka, P., Schlenk, R.F., Gaidzik, V.I., Habdank, M., Krönke, J., Bullinger, L., Späth, D., Kayser, S., Zucknick, M., Götze, K., et al. (2010). IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* **28**, 3636–3643.
- Pate, K.T., Stringari, C., Sprowl-Tanio, S., Wang, K., TeSlaa, T., Hoverter, N.P., McQuade, M.M., Garner, C., Digman, M.A., Teitell, M.A., et al. (2014). Wnt signaling directs a metabolic program of glycolysis and angiogenesis in colon cancer. *EMBO J.* **33**, 1454–1473.
- Phang, J.M., Donald, S.P., Pandhare, J., and Liu, Y. (2008). The metabolism of proline, a stress substrate, modulates carcinogenic pathways. *Amino Acids* **35**, 681–690.
- Pieters, R., Hunger, S.P., Boos, J., Rizzari, C., Silverman, L., Baruchel, A., Goekbuget, N., Schrappe, M., and Pui, C.H. (2011). L-asparaginase treatment in acute lymphoblastic leukemia: a focus on *Erwinia* asparaginase. *Cancer* **117**, 238–249.
- Plas, D.R., Talapatra, S., Edinger, A.L., Rathmell, J.C., and Thompson, C.B. (2001). Akt and Bcl-xL promote growth factor-independent survival through distinct effects on mitochondrial physiology. *J. Biol. Chem.* **276**, 12041–12048.
- Possemato, R., Marks, K.M., Shaul, Y.D., Pacold, M.E., Kim, D., Birsoy, K., Sethumadhavan, S., Woo, H.K., Jang, H.G., Jha, A.K., et al. (2011). Functional genomics reveal that the serine synthesis pathway is essential in breast cancer. *Nature* **476**, 346–350.
- Rathmell, J.C., Vander Heiden, M.G., Harris, M.H., Frauwirth, K.A., and Thompson, C.B. (2000). In the absence of extrinsic signals, nutrient utilization by lymphocytes is insufficient to maintain either cell size or viability. *Mol. Cell* **6**, 683–692.
- Rathmell, J.C., Fox, C.J., Plas, D.R., Hammerman, P.S., Cinalli, R.M., and Thompson, C.B. (2003). Akt-directed glucose metabolism can prevent Bax conformation change and promote growth factor-independent survival. *Mol. Cell Biol.* **23**, 7315–7328.
- Reynolds, M.R., Lane, A.N., Robertson, B., Kemp, S., Liu, Y., Hill, B.G., Dean, D.C., and Clem, B.F. (2014). Control of glutamine metabolism by the tumor suppressor Rb. *Oncogene* **33**, 556–566.
- Richman, E.L., Kenfield, S.A., Stampfer, M.J., Giovannucci, E.L., Zeisel, S.H., Willett, W.C., and Chan, J.M. (2012). Choline intake and risk of lethal prostate cancer: incidence and survival. *Am. J. Clin. Nutr.* **96**, 855–863.
- Rivera, S., Azcón-Bieto, J., López-Soriano, F.J., Miralpeix, M., and Argilés, J.M. (1988). Amino acid metabolism in tumour-bearing mice. *Biochem. J.* **249**, 443–449.
- Roberts, E., and Frankel, S. (1949). Free amino acids in normal and neoplastic tissues of mice as studied by paper chromatography. *Cancer Res.* **9**, 645–648, 643 pl.
- Robitaille, A.M., Christen, S., Shimobayashi, M., Cornu, M., Fava, L.L., Moes, S., Prescianotto-Baschong, C., Sauer, U., Jenoe, P., and Hall, M.N. (2013). Quantitative phosphoproteomics reveal mTORC1 activates de novo pyrimidine synthesis. *Science* **339**, 1320–1323.
- Rohle, D., Popovici-Muller, J., Palaskas, N., Turcan, S., Grommes, C., Campos, C., Tsoi, J., Clark, O., Oldrini, B., Komisopoulou, E., et al. (2013). An inhibitor of mutant IDH1 delays growth and promotes differentiation of glioma cells. *Science* **340**, 626–630.
- Rothberg, J.M., Bailey, K.M., Wojtkowiak, J.W., Ben-Nun, Y., Bogoy, M., Weber, E., Moin, K., Blum, G., Mattingly, R.R., Gillies, R.J., and Sloane, B.F. (2013). Acid-mediated tumor proteolysis: contribution of cysteine cathepsins. *Neoplasia* **15**, 1125–1137.
- Rysman, E., Brusselmans, K., Scheys, K., Timmermans, L., Derua, R., Munck, S., Van Veldhoven, P.P., Waltregny, D., Daniëls, V.W., Machiels, J., et al. (2010). De novo lipogenesis protects cancer cells from free radicals and chemotherapeutics by promoting membrane lipid saturation. *Cancer Res.* **70**, 8117–8126.
- Sabari, B.R., Tang, Z., Huang, H., Yong-Gonzalez, V., Molina, H., Kong, H.E., Dai, L., Shimada, M., Cross, J.R., Zhao, Y., et al. (2015). Intracellular crotonyl-CoA stimulates transcription through p300-catalyzed histone crotonylation. *Mol. Cell* **58**, 203–215.
- Sasaki, M., Knobbe, C.B., Munger, J.C., Lind, E.F., Brenner, D., Brüstle, A., Harris, I.S., Holmes, R., Wakeham, A., Haight, J., et al. (2012). IDH1(R132H) mutation increases murine hematopoietic progenitors and alters epigenetics. *Nature* **488**, 656–659.
- Sauer, L.A., Stayman, J.W., 3rd, and Dauchy, R.T. (1982). Amino acid, glucose, and lactic acid utilization in vivo by rat tumors. *Cancer Res.* **42**, 4090–4097.
- Schafer, Z.T., Grassian, A.R., Song, L., Jiang, Z., Gerhart-Hines, Z., Irie, H.Y., Gao, S., Puigserver, P., and Brugge, J.S. (2009). Antioxidant and oncogene rescue of metabolic defects caused by loss of matrix attachment. *Nature* **461**, 109–113.
- Schmid, S.A., Gaumann, A., Wondrak, M., Eckermann, C., Schulte, S., Mueller-Klieser, W., Wheatley, D.N., and Kunz-Schughart, L.A. (2007). Lactate adversely affects the in vitro formation of endothelial cell tubular structures through the action of TGF-beta1. *Exp. Cell Res.* **313**, 2531–2549.
- Schug, Z.T., Peck, B., Jones, D.T., Zhang, Q., Grosskurth, S., Alam, I.S., Goodwin, L.M., Smethurst, E., Mason, S., Blyth, K., et al. (2015). Acetyl-CoA synthetase 2 promotes acetate utilization and maintains cancer cell growth under metabolic stress. *Cancer Cell* **27**, 57–71.
- Schwer, B., and Verdin, E. (2008). Conserved metabolic regulatory functions of sirtuins. *Cell Metab.* **7**, 104–112.
- Selak, M.A., Armour, S.M., MacKenzie, E.D., Boulahbel, H., Watson, D.G., Mansfield, K.D., Pan, Y., Simon, M.C., Thompson, C.B., and Gottlieb, E. (2005). Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase. *Cancer Cell* **7**, 77–85.
- Sellers, K., Fox, M.P., Bousamra, M., 2nd, Slone, S.P., Higashi, R.M., Miller, D.M., Wang, Y., Yan, J., Yuneva, M.O., Deshpande, R., et al. (2015). Pyruvate carboxylase is critical for non-small-cell lung cancer proliferation. *J. Clin. Invest.* **125**, 687–698.
- Shi, L., and Tu, B.P. (2013). Acetyl-CoA induces transcription of the key G1 cyclin CLN3 to promote entry into the cell division cycle in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **110**, 7318–7323.
- Shi, Y., Lan, F., Matson, C., Mulligan, P., Whetstone, J.R., Cole, P.A., Casero, R.A., and Shi, Y. (2004). Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* **119**, 941–953.
- Shim, E.H., Livi, C.B., Rakheja, D., Tan, J., Benson, D., Parekh, V., Kho, E.Y., Ghosh, A.P., Kirkman, R., Velu, S., et al. (2014). L-2-Hydroxyglutarate: an epigenetic modifier and putative oncometabolite in renal cancer. *Cancer Discov.* **4**, 1290–1298.

- Shyh-Chang, N., Locasale, J.W., Lyssiotis, C.A., Zheng, Y., Teo, R.Y., Ratanasirintrao, S., Zhang, J., Onder, T., Unternaehrer, J.J., Zhu, H., et al. (2013). Influence of threonine metabolism on S-adenosylmethionine and histone methylation. *Science* 339, 222–226.
- Sircar, K., Huang, H., Hu, L., Cogdell, D., Dhillon, J., Tzelepi, V., Efstathiou, E., Koumakpayi, I.H., Saad, F., Luo, D., et al. (2012). Integrative molecular profiling reveals asparagine synthetase is a target in castration-resistant prostate cancer. *Am. J. Pathol.* 180, 895–903.
- Som, P., Atkins, H.L., Bandoyadhyay, D., Fowler, J.S., MacGregor, R.R., Matsui, K., Oster, Z.H., Sacker, D.F., Shiue, C.Y., Turner, H., et al. (1980). A fluorinated glucose analog, 2-fluoro-2-deoxy-D-glucose (F-18): nontoxic tracer for rapid tumor detection. *J. Nucl. Med.* 21, 670–675.
- Son, J., Lyssiotis, C.A., Ying, H., Wang, X., Hua, S., Ligorio, M., Perera, R.M., Ferrone, C.R., Mullarky, E., Shyh-Chang, N., et al. (2013). Glutamine supports pancreatic cancer growth through a KRAS-regulated metabolic pathway. *Nature* 496, 101–105.
- Sonveaux, P., Copetti, T., De Saedeleer, C.J., Végran, F., Verrax, J., Kennedy, K.M., Moon, E.J., Dhup, S., Danhier, P., Frérart, F., et al. (2012). Targeting the lactate transporter MCT1 in endothelial cells inhibits lactate-induced HIF-1 activation and tumor angiogenesis. *PLoS ONE* 7, e33418.
- Spiro, R.G. (2002). Protein glycosylation: nature, distribution, enzymatic formation, and disease implications of glycopeptide bonds. *Glycobiology* 12, 43R–56R.
- Sporn, M.B., and Liby, K.T. (2012). NRF2 and cancer: the good, the bad and the importance of context. *Nat. Rev. Cancer* 12, 564–571.
- Stern, R., Shuster, S., Neudecker, B.A., and Formby, B. (2002). Lactate stimulates fibroblast expression of hyaluronan and CD44: the Warburg effect revisited. *Exp. Cell Res.* 276, 24–31.
- Stolzinger, A., and Grune, T. (2004). Neuronal apoptotic bodies: phagocytosis and degradation by primary microglial cells. *FASEB J.* 18, 743–745.
- Strohecker, A.M., Guo, J.Y., Karsli-Uzunbas, G., Price, S.M., Chen, G.J., Mathew, R., McMahon, M., and White, E. (2013). Autophagy sustains mitochondrial glutamine metabolism and growth of BRAFV600E-driven lung tumors. *Cancer Discov.* 3, 1272–1285.
- Sullivan, L.B., and Chandel, N.S. (2014). Mitochondrial reactive oxygen species and cancer. *Cancer Metab.* 2, 17.
- Sullivan, L.B., Gui, D.Y., Hosios, A.M., Bush, L.N., Freinkman, E., and Vander Heiden, M.G. (2015). Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferating Cells. *Cell* 162, 552–563.
- Sun, Q., Luo, T., Ren, Y., Florey, O., Shirasawa, S., Sasazuki, T., Robinson, D.N., and Overholtzer, M. (2014). Competition between human cells by entosis. *Cell Res.* 24, 1299–1310.
- Svastová, E., Hulíková, A., Rafajová, M., Zát'ovicová, M., Gibadulinová, A., Casini, A., Cecchi, A., Scozzafava, A., Supuran, C.T., Pastorek, J., and Pastorková, S. (2004). Hypoxia activates the capacity of tumor-associated carbonic anhydrase IX to acidify extracellular pH. *FEBS Lett.* 577, 439–445.
- Swietach, P., Vaughan-Jones, R.D., and Harris, A.L. (2007). Regulation of tumor pH and the role of carbonic anhydrase 9. *Cancer Metastasis Rev.* 26, 299–310.
- Tan, A.S., Baty, J.W., Dong, L.F., Bezawork-Geleta, A., Endaya, B., Goodwin, J., Bajzikova, M., Kovarova, J., Peterka, M., Yan, B., et al. (2015). Mitochondrial genome acquisition restores respiratory function and tumorigenic potential of cancer cells without mitochondrial DNA. *Cell Metab.* 21, 81–94.
- Terunuma, A., Putluri, N., Mishra, P., Mathé, E.A., Dorsey, T.H., Yi, M., Wallace, T.A., Issaq, H.J., Zhou, M., Killian, J.K., et al. (2014). MYC-driven accumulation of 2-hydroxyglutarate is associated with breast cancer prognosis. *J. Clin. Invest.* 124, 398–412.
- Thompson, C.B. (2011). Rethinking the regulation of cellular metabolism. *Cold Spring Harb. Symp. Quant. Biol.* 76, 23–29.
- Tibbetts, A.S., and Appling, D.R. (2010). Compartmentalization of Mammalian folate-mediated one-carbon metabolism. *Annu. Rev. Nutr.* 30, 57–81.
- Tomlinson, I.P., Alam, N.A., Rowan, A.J., Barclay, E., Jaeger, E.E., Kelsell, D., Leigh, I., Gorman, P., Lamlium, H., Rahman, S., et al.; Multiple Leiomyoma Consortium (2002). Germline mutations in FH predispose to dominantly inherited uterine fibroids, skin leiomyomata and papillary renal cell cancer. *Nat. Genet.* 30, 406–410.
- Towbin, B.D., González-Aguilera, C., Sack, R., Gaidatzis, D., Kalck, V., Meister, P., Askjaer, P., and Gasser, S.M. (2012). Step-wise methylation of histone H3K9 positions heterochromatin at the nuclear periphery. *Cell* 150, 934–947.
- Turcan, S., Rohle, D., Goenka, A., Walsh, L.A., Fang, F., Yilmaz, E., Campos, C., Fabis, A.W., Lu, C., Ward, P.S., et al. (2012). IDH1 mutation is sufficient to establish the glioma hypermethylator phenotype. *Nature* 483, 479–483.
- Vacchelli, E., Aranda, F., Eggermont, A., Sautès-Fridman, C., Tartour, E., Kennedy, E.P., Platten, M., Zitvogel, L., Kroemer, G., and Galluzzi, L. (2014). Trial watch: IDO inhibitors in cancer therapy. *Oncolimmunology* 3, e957994.
- Vander Heiden, M.G., Cantley, L.C., and Thompson, C.B. (2009). Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 324, 1029–1033.
- Vaupel, P., Kallinowski, F., and Okunieff, P. (1989). Blood flow, oxygen and nutrient supply, and metabolic microenvironment of human tumors: a review. *Cancer Res.* 49, 6449–6465.
- Végran, F., Boidot, R., Michiels, C., Sonveaux, P., and Feron, O. (2011). Lactate influx through the endothelial cell monocarboxylate transporter MCT1 supports an NF- κ B/IL-8 pathway that drives tumor angiogenesis. *Cancer Res.* 71, 2550–2560.
- Venneti, S., Dunphy, M.P., Zhang, H., Pitter, K.L., Zanzonico, P., Campos, C., Carlin, S.D., La Rocca, G., Lyashchenko, S., Ploessl, K., et al. (2015). Glutamine-based PET imaging facilitates enhanced metabolic evaluation of gliomas in vivo. *Sci. Transl. Med.* 7, 274ra17.
- Viale, A., Pettazzoni, P., Lyssiotis, C.A., Ying, H., Sánchez, N., Marchesini, M., Carugo, A., Green, T., Seth, S., Giuliani, V., et al. (2014). Oncogene ablation-resistant pancreatic cancer cells depend on mitochondrial function. *Nature* 514, 628–632.
- Vogelstein, B., and Kinzler, K.W. (2004). Cancer genes and the pathways they control. *Nat. Med.* 10, 789–799.
- Wahlström, T., and Henriksson, M.A. (2015). Impact of MYC in regulation of tumor cell metabolism. *Biochim. Biophys. Acta* 1849, 563–569.
- Wang, C., Guo, K., Gao, D., Kang, X., Jiang, K., Li, Y., Sun, L., Zhang, S., Sun, C., Liu, X., et al. (2011a). Identification of transaldolase as a novel serum biomarker for hepatocellular carcinoma metastasis using xenografted mouse model and clinic samples. *Cancer Lett.* 313, 154–166.
- Wang, R., Dillon, C.P., Shi, L.Z., Milasta, S., Carter, R., Finkelstein, D., McCormick, L.L., Fitzgerald, P., Chi, H., Munger, J., and Green, D.R. (2011b). The transcription factor Myc controls metabolic reprogramming upon T lymphocyte activation. *Immunity* 35, 871–882.
- Wang, F., Travins, J., DeLaBarre, B., Penard-Lacronique, V., Schalm, S., Hansen, E., Straley, K., Kernysky, A., Liu, W., Gliser, C., et al. (2013). Targeted inhibition of mutant IDH2 in leukemia cells induces cellular differentiation. *Science* 340, 622–626.
- Warburg, O. (1956). On the origin of cancer cells. *Science* 123, 309–314.
- Warburg, O.P.K., and Negelein, E. (1924). Über den Stoffwechsel der Carcinomzelle. *Biochem Zeitschr* 152, 309–344.
- Warburg, O., Wind, F., and Negelein, E. (1927). The metabolism of tumors in the body. *J. Gen. Physiol.* 8, 519–530.
- Ward, P.S., Patel, J., Wise, D.R., Abdel-Wahab, O., Bennett, B.D., Collier, H.A., Cross, J.R., Fantin, V.R., Hedvat, C.V., Perl, A.E., et al. (2010). The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell* 17, 225–234.
- Wellen, K.E., Hatzivassiliou, G., Sachdeva, U.M., Bui, T.V., Cross, J.R., and Thompson, C.B. (2009). ATP-citrate lyase links cellular metabolism to histone acetylation. *Science* 324, 1076–1080.
- Wellen, K.E., Lu, C., Mancuso, A., Lemons, J.M., Ryzcko, M., Dennis, J.W., Rabinowitz, J.D., Collier, H.A., and Thompson, C.B. (2010). The hexosamine biosynthetic pathway couples growth factor-induced glutamine uptake to glucose metabolism. *Genes Dev.* 24, 2784–2799.

Wieman, H.L., Wofford, J.A., and Rathmell, J.C. (2007). Cytokine stimulation promotes glucose uptake via phosphatidylinositol-3 kinase/Akt regulation of Glut1 activity and trafficking. *Mol. Biol. Cell* **18**, 1437–1446.

Wise, D.R., DeBerardinis, R.J., Mancuso, A., Sayed, N., Zhang, X.Y., Pfeiffer, H.K., Nissim, I., Daikhin, E., Yudkoff, M., McMahon, S.B., and Thompson, C.B. (2008). Myc regulates a transcriptional program that stimulates mitochondrial glutaminolysis and leads to glutamine addiction. *Proc. Natl. Acad. Sci. USA* **105**, 18782–18787.

Wise, D.R., Ward, P.S., Shay, J.E., Cross, J.R., Gruber, J.J., Sachdeva, U.M., Platt, J.M., DeMatteo, R.G., Simon, M.C., and Thompson, C.B. (2011). Hypoxia promotes isocitrate dehydrogenase-dependent carboxylation of α -ketoglutarate to citrate to support cell growth and viability. *Proc. Natl. Acad. Sci. USA* **108**, 19611–19616.

Wisniewski, J.R., Zougman, A., and Mann, M. (2008). Nepsilon-formylation of lysine is a widespread post-translational modification of nuclear proteins occurring at residues involved in regulation of chromatin function. *Nucleic Acids Res.* **36**, 570–577.

Wu, D., Si, W., Wang, M., Lv, S., Ji, A., and Li, Y. (2015). Hydrogen sulfide in cancer: friend or foe? *Nitric Oxide* **50**, 38–45.

Xiao, M., Yang, H., Xu, W., Ma, S., Lin, H., Zhu, H., Liu, L., Liu, Y., Yang, C., Xu, Y., et al. (2012). Inhibition of α -KG-dependent histone and DNA demethylases by fumarate and succinate that are accumulated in mutations of FH and SDH tumor suppressors. *Genes Dev.* **26**, 1326–1338.

Xie, Z., Dai, J., Dai, L., Tan, M., Cheng, Z., Wu, Y., Boeke, J.D., and Zhao, Y. (2012). Lysine succinylation and lysine malonylation in histones. *Mol. Cell. Proteomics* **11**, 100–107.

Xu, X., Gammon, M.D., Zeisel, S.H., Bradshaw, P.T., Wetmur, J.G., Teitelbaum, S.L., Neugut, A.I., Santella, R.M., and Chen, J. (2009a). High intakes of choline and betaine reduce breast cancer mortality in a population-based study. *FASEB J.* **23**, 4022–4028.

Xu, X., Zur Hausen, A., Coy, J.F., and Löchelt, M. (2009b). Transketolase-like protein 1 (TKTL1) is required for rapid cell growth and full viability of human tumor cells. *Int. J. Cancer* **124**, 1330–1337.

Xu, W., Yang, H., Liu, Y., Yang, Y., Wang, P., Kim, S.H., Ito, S., Yang, C., Wang, P., Xiao, M.T., et al. (2011). Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of α -ketoglutarate-dependent dioxygenases. *Cancer Cell* **19**, 17–30.

Yan, S.R., and Berton, G. (1996). Regulation of Src family tyrosine kinase activities in adherent human neutrophils. Evidence that reactive oxygen intermediates produced by adherent neutrophils increase the activity of the p58c-fgr and p53/56lyn tyrosine kinases. *J. Biol. Chem.* **271**, 23464–23471.

Yanagida, O., Kanai, Y., Chairoungdua, A., Kim, D.K., Segawa, H., Nii, T., Cha, S.H., Matsuo, H., Fukushima, J., Fukasawa, Y., et al. (2001). Human L-type amino acid transporter 1 (LAT1): characterization of function and expression in tumor cell lines. *Biochim. Biophys. Acta* **1514**, 291–302.

Ye, J., Mancuso, A., Tong, X., Ward, P.S., Fan, J., Rabinowitz, J.D., and Thompson, C.B. (2012). Pyruvate kinase M2 promotes de novo serine synthesis to sustain mTORC1 activity and cell proliferation. *Proc. Natl. Acad. Sci. USA* **109**, 6904–6909.

Ye, J., Fan, J., Venneti, S., Wan, Y.W., Pawel, B.R., Zhang, J., Finley, L.W., Lu, C., Lindsten, T., Cross, J.R., et al. (2014). Serine catabolism regulates mitochondrial redox control during hypoxia. *Cancer Discov.* **4**, 1406–1417.

Ying, H., Kimmelman, A.C., Lyssiotis, C.A., Hua, S., Chu, G.C., Fletcher-Sanankone, E., Locasale, J.W., Son, J., Zhang, H., Coloff, J.L., et al. (2012). Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* **149**, 656–670.

Ying, J., Rahbar, M.H., Hallman, D.M., Hernandez, L.M., Spitz, M.R., Forman, M.R., and Gorlova, O.Y. (2013). Associations between dietary intake of choline and betaine and lung cancer risk. *PLoS ONE* **8**, e54561.

Yoon, C.Y., Shim, Y.J., Kim, E.H., Lee, J.H., Won, N.H., Kim, J.H., Park, I.S., Yoon, D.K., and Min, B.H. (2007). Renal cell carcinoma does not express argininosuccinate synthetase and is highly sensitive to arginine deprivation via arginine deiminase. *Int. J. Cancer* **120**, 897–905.

Young, R.M., Ackerman, D., Quinn, Z.L., Mancuso, A., Gruber, M., Liu, L., Giannoukos, D.N., Bobrovnikova-Marjon, E., Diehl, J.A., Keith, B., and Simon, M.C. (2013). Dysregulated mTORC1 renders cells critically dependent on de-saturated lipids for survival under tumor-like stress. *Genes Dev.* **27**, 1115–1131.

Yuneva, M., Zamboni, N., Oefner, P., Sachidanandam, R., and Lazebnik, Y. (2007). Deficiency in glutamine but not glucose induces MYC-dependent apoptosis in human cells. *J. Cell Biol.* **178**, 93–105.

Yuneva, M.O., Fan, T.W., Allen, T.D., Higashi, R.M., Ferraris, D.V., Tsukamoto, T., Matés, J.M., Alonso, F.J., Wang, C., Seo, Y., et al. (2012). The metabolic profile of tumors depends on both the responsible genetic lesion and tissue type. *Cell Metab.* **15**, 157–170.

Zhang, C.X., Pan, M.X., Li, B., Wang, L., Mo, X.F., Chen, Y.M., Lin, F.Y., and Ho, S.C. (2013). Choline and betaine intake is inversely associated with breast cancer risk: a two-stage case-control study in China. *Cancer Sci.* **104**, 250–258.

Zhang, J., Fan, J., Venneti, S., Cross, J.R., Takagi, T., Bhinder, B., Djaballah, H., Kanai, M., Cheng, E.H., Judkins, A.R., et al. (2014). Asparagine plays a critical role in regulating cellular adaptation to glutamine depletion. *Mol. Cell* **56**, 205–218.