

IMMUNOLOGIC DEFICIENCY SYNDROMES

V. Žampachová
I. PAŮLF MU

IMMUNODEFICIENCY

Presentation as infections

- Serious
- Persistent
- Unusual
- Recurrent

PRIMARY IMMUNODEFICIENCY

- genetically determined
- humoral and/or cellular arms of adaptive immunity (mediated by B and T lymphocytes) – defects in maturation and/or activation
- defense mechanisms of innate immunity (NK cells, phagocytes, complement)

PRIMARY IMMUNODEFICIENCY

- manifestation mostly in infancy, (6-24 months)
- susceptibility to:
 - recurrent infections by opportunistic pathogens
 - systemic inf. by microorg. normally superficial,
 - unusually extensive inf. by common pathogens
- autoimmune diseases (disorder of regulation)

T-cell defect

- Bacterial sepsis
- Cytomegalovirus, Epstein-Barr virus, severe varicella, chronic infections with respiratory and intestinal viruses
- *Candida, Aspergillus, Pneumocystis jirovecii*
- Aggressive disease with opportunistic pathogens, failure to clear infections

B-cell defect

- Streptococci, staphylococci, *Haemophilus* –skin, respiratory tract
- Enteroviral enteritis, encephalitis
- Severe intestinal giardiasis (protozoan), other GIT infections
- Recurrent sinopulmonary infections, sepsis, chronic meningitis
- panhypogammaglobulinemia x selective deficiency
- ! immunization w. live vaccines (polio, etc.)

Granulocyte defect

- Staphylococci, *Pseudomonas*
- *Candida*, *Nocardia*, *Aspergillus*
- Leukocyte adhesion deficiency
- Defects in phagolysosome function
- Defects in microbicidal activity

Complement defect

- Neisserial infections, other pyogenic infections
- Defects in complement components
- Defects in complement system regulators

PRIMARY IMMUNODEFICIENCY

- **mainly B-cell defect:** X-linked agammaglobulinemia of Bruton, transient hypogammaglobulinemia of infancy, selective IgA deficiency, common variable immunodeficiency (CVID)
- **mainly T-cell defect:** DiGeorge syndrom (thymic hypoplasia), hyper-IgM syndrome
- **B- and T-cell defect:** severe combined immunodeficiency (SCID), Wiskott-Aldrich syndrome (immunodeficiency with thrombocytopenia and eczema – systemic disorder)
- **defect in phagocyte function:** chronic granulomatous disease, leukocyte adhesion deficiency, myeloperoxidase deficiency
- **primary complement deficiencies**

Severe combined immunodeficiency

SCID

- defects in both humoral and cell-mediated immunity
- recurrent, severe infections, wide range of pathogens, incl. *Candida albicans*, *P. jirovecii*, *Pseudomonas*, cytomegalovirus, varicella, many bacteria.
- morbilliform rash shortly after birth – GVH disease due to maternal T-cells
- X-linked (~ 50%), autosomal recessive
- „bubble children“, bone marrow transplantation, gene therapy (! acute T-cell leukemia)

DiGeorge syndrome (thymic hypoplasia)

- T-cell defect; chromosomal deletion; commonly only partial hypoplasia
- failure of embryonal development of the 3rd and 4th pharyngeal pouches (thymus, parathyroids, part of thyroid clear cells → hypocalcemic tetany; heart + great vessels defects)
- T- cell zones depleted (LN – paracortical, periarteriolar sheaths of the spleen)
- ↑ fungal and viral infections

Common variable immunodeficiency

CVID

- relatively common, heterogeneous group of disorders (dg. by exclusion), both sexes, children - adolescents
- hypogammaglobulinemia
- sporadic and inherited forms
- B cells in normal numbers, not able to differentiate into plasma cells
- intrinsic B-cell defects, abnormalities in T helper cell-mediated activation of B cells
- hyperplastic B-cell zones in lymphoid tissue

Common variable immunodeficiency

- recurrent sinopulmonary pyogenic infections
- recurrent herpesvirus infections
- persistent diarrhea due to *G. lamblia*
- enterovirus meningoencephalitis
- ↑ frequency of autoimmune diseases (RA)
- risk of lymphoid malignancy

Isolated IgA deficiency

- common immunodeficiency in Caucasians (1:600), severe reaction after blood transfusion possible
- familial or acquired after some infections (toxoplasmosis, measles, some viral inf.)
- low levels of both serum and secretory IgA
- mostly asymptomatic; possible respiratory, GIT, urogenital recurrent infections
- ↑respiratory tract allergy, autoimmune diseases

Chronic granulomatous disease

- ↓ oxygen radicals production needed for bacteria killing in effective phagocytosis
- X-linked; other types
- ↑ pyogenic bacteria (Staph., G- rods), fungi (Aspergillus)
- respiratory, GIT, skin, ... infections – abscess, giant-cell granuloma
- liver vascular lesions → portal hypertension

SECONDARY IMMUNODEFICIENCY

Due to impaired synthesis and function:

- protein, vitamin and energy deficiency in malnutrition, cachexia in disseminated cancer, anorexia, alcoholism
- prevalent monoclonal Ig in some lymphoproliferative diseases
- bone marrow infiltration or fibrosis (leukemia, myelofibrosis)
- suppression of cell mediated immunity due to acute viral infection (CMV, EBV, measles, etc.), bacterial and protozoal infection – macrophagic dysfunction (leprosy, leishmaniasis)

SECONDARY IMMUNODEFICIENCY

- **iatrogenic** (immunosuppressive and cytostatic drugs, radiotherapy, splenectomy – pneumococcus sepsis)
- diabetes mellitus and other metabolic diseases
- chronic stress
- sarcoidosis (↓ Tcell function)
- certain age groups (old, newborn, immature infants)

Increased catabolism or loss: nephrotic syndrome and renal failure, inflammatory intestinal diseases (IBD, lymphangiectasia)

SECONDARY IMMUNODEFICIENCY

Humoral immunodeficiency

- intestinal lymphangiectasia, IBD → ↓ all Ig classes, commonly + lymphopenia
- nephrotic sy, chronic diarrhea → ↓ IgG
- iatrogenic immunosuppression/cytostatic therapy
- B-cell malignancies
- Splenectomy – spleen B-cell – Ab x polysaccharide antigens – encapsulated microorg. – vaccination x pneumococci

SECONDARY IMMUNODEFICIENCY

Cellular immunodeficiency

- temporary after acute viral infection (CMV, EBV, measles, COVID, etc.)
- iatrogenic immunosuppression/cytostatic
- AIDS

SECONDARY IMMUNODEFICIENCY

Combined immunodeficiency

- Severe general metabolic problems (DM, renal insufficiency), malnutrition, anorexia, chronic alcoholics – inadequate hormones, glucose, vitamins level

SECONDARY IMMUNODEFICIENCY

Defect of phagocytosis


- neutropenia in bone marrow insufficiency
(irradiation, immunosuppressant/cytostatic th.,
some chemicals)
- autoantibodies
- ↑ loss in hypersplenism
- metabolic diaseases
- myeloid leukemia

SECONDARY IMMUNODEFICIENCY


Complement deficiency

- immunocomplex diseases
- sepsis
- severe liver disease


Brain mycotic abscess




Lung mycotic abscess




Fungal structures in necrotic brain




Cryptococcus in skin




Toxoplasmosis in myocardium



Brain toxoplasmosis inflammation



Brain toxoplasmosis cysts



HIV - AIDS

- The goal – ending the *AIDS* epidemic by 2030
- Stopping the new infections
- Everybody infected should have access to treatment
- Pre-exposure prophylaxis

AIDS epidemics

- Estimated 39 million people living with HIV in 2022 (1,5 mil children); 9,2 mil. without therapy
- 1,3 mil newly infected in 2022, 46 % in females
- 630 000 died of AIDS-related illnesses in 2022
- 85,6 million infected since the start of the epidemics
- 40,4 million have died

AIDS epidemics

- AIDS related illnesses important cause of death among women of reproductive age (15–49 years) globally

AIDS epidemics

- Decrease in new infections and deaths, increase in number of people living with HIV (Highly Active Antiretroviral Therapy - HAART)

AIDS epidemics

- HIV infection outside of sub-Saharan Africa: National epidemics concentrated among key populations at higher risk (men who have sex with men – MSM, injecting drug users; prisoners, sex workers, sexual partners of key population) – 94 % of new infections

HIV ISSUES

- Blood safety
- HIV treatment: antiretroviral therapy should begin immediately after diagnosis
- Prevention of mother-to-child transmission
- Co-management of tuberculosis and HIV treatment
- HIV testing in the general and most-at-risk population
- Preexposure prophylaxis

HIV - AIDS

More than 90% of children living with HIV acquired the virus during pregnancy, birth or breastfeeding - forms of HIV transmission that can be prevented.

AIDS epidemics

- Europe, Australia and Canada: mortality rates among people living with HIV in the first five years after infection now ~ in the HIV-uninfected population
- Mortality among HIV-infected people increases with the duration of infection
- Increasing complications of chronic HAART – highly active antiretroviral therapy

Noninfectious HIV-related comorbidities:

- The premature aging process in HIV-infected people
- 2x ↑ risk of myocardial infarction
- ↑ risk of osteoporosis incl. fractures (even in adolescents!)
- ↑ risk of chronic renal failure
- Non-AIDS tumors

Noninfectious HIV-related comorbidities:

- hyper-activated immunological profile, accelerated T-cell senescence
- accelerated process of immune senescence and inflammatory aging during HIV infection → increased risk of age-related diseases
- long-term tolerability of HAART regimens

HAART complications

- Diarrhea, nausea , and vomiting.
- Lipodystrophy: fat in adipous tissue redistributed to other regions, i.e.face and limbs → thin, breasts, stomach and/or neck enlarge.
- Glucose intolerance, diabetes. Lactic acidosis.
- Liver toxicity – acute hepatitis incl. liver failure. Pancreatitis.
- Nephrotoxicity
- Neuropathy
- Osteonecrosis, osteoporosis, osteopenia

HAART complications

- Cardiovascular complications: toxicity, endothelial dysfunction, atherosclerosis, dyslipidemia. Myocardial infarction
- Reconstitution of the immune system (major goal of HAART treatment): risk of debilitating Immune reconstitution inflammatory syndrome (IRIS) - ↑ CD4 count + function. Immune response against antigens associated with infection diseases (TB, MAC, Pneumocystis pneumonia, CMV, HZV). 10-25% of patients
- Drug interactions

HAART resistance

- Drug switching necessary
- New regimens/drugs
- Timely start of therapy

HIV-2

- Endemic in West Africa.
- Limited spread outside this area, suspicion – in persons of West African origin/risk contact
- Prevalence of HIV-2 disproportionately high in countries with strong socioeconomic ties to West Africa (e.g., France; Spain; Portugal; and former Portuguese colonies such as Brazil, Angola, Mozambique, and parts of India near Goa).

HIV-2

- Clinical course - longer asymptomatic stage, lower plasma HIV-2 viral loads, and lower mortality rates compared with HIV-1 infection
- Resistance-associated mutations develop commonly in HIV-2 patients on therapy

HIV infection of cells

- T-lymphocytes (CD4+)
- macrophages/monocytes (viral reservoir, replication and transport)
- mucosal and follicular dendritic cells
- cells in CNS (microglia)

Immune dysfunctions in AIDS

- **Lymphopenia** (selective loss of CD4+ T-cells – direct cytophatic effect, apoptosis of noninfected)
- **Decreased T-cell function in vivo** (loss of memory T-cells, susceptibility to opportunistic infections and neoplasms, decreased delayed-type hypersensitivity)
- **Polyclonal B-cell activation** (hypergammaglobulinemia, CIC, inability of new antibody response)
- **Altered monocyte or macrophage functions** (decreased chemotaxis, phagocytosis, antigen presentation; increased spontaneous secretion of TNF, IL-1 etc.)

Phases of HIV infection

- **Acute retroviral syndrome** (3-6 wks after infection, in 40-90%, self-limited in 2-4 wks)
- **Chronic phase** (clinical latency, persistent generalized lymphadenopathy – PGL)
- **Progression to AIDS** (AIDS-related complex – ARC, AIDS indicator conditions: constitutional, neurologic, opportunistic infection, neoplasm)


Acute HIV infection

- **Suspect:** Signs or symptoms of acute HIV infection with recent (within 2–6 weeks) high risk of exposure
- **Possible signs:** fever, lymphadenopathy, skin rash, myalgia/arthralgia, headache, diarrhea, oral ulcers, leucopenia, thrombocytopenia, transaminase elevation.

Acute HIV infection

- **High-risk exposures** include sexual contact with a person infected with HIV or at risk of HIV, sharing of injection drug use paraphernalia, or contact of potentially infectious blood with mucous membranes or breaks in skin.
- **Differential diagnosis:** Epstein-Barr virus (EBV)- and non-EBV (e.g., cytomegalovirus [CMV])-related infectious mononucleosis syndromes, influenza, viral hepatitis, streptococcal infection, syphilis

Persistent generalized lymphadenopathy



Opportunistic infections and neoplasms

- **Protozoal and helminthic** (cryptosporidiosis, toxoplasmosis, giardiasis, etc.)
- **Fungal** (Pneumocystis, candidiasis, cryptococcosis, coccidiomycosis, histoplasmosis)
- **Bacterial** (mycobacteriosis – atypical, TB; salmonellosis, nocardiosis)
- **Viral** (CMV, Herpes simplex, Varicella-zoster, progressive multifocal leukoencephalopathy – JC polyoma virus)
- **Neoplasms** (Kaposi sarcoma – HHV 8, B-cell non-Hodgkin lymphomas, primary brain lymphomas – EBV, aggressive cervical and anal carcinomas – HPV)

HIV neurologic disease

- *Acute* aseptic meningitis
- *subacute and chronic*: HIV-associated neurocognitive disorders
- HIV meningoencephalitis – AIDS-dementia complex, vacuolar myelopathy, myopathy and peripheral neuropathy
- before HAART, clinical signs of neurologic lesion in 40-60% of patients (HIV, opportunistic infection, tumor)
- now ↓ – chronic encephalitis – microglial nodules + multinucleated giant cell, microfoci of necrosis

10981-07

11/12/06

20:21

0270 10


← H O U D




GRY: 414
CLR: FED
ZOOM: 1
HDS: 100
LEV: 15




HIV encephalopathy



HIV encephalitis



p24 immunohistochemistry



Copyright © 2002, Elsevier Science (USA). All rights reserved.

CNS infections

- Toxoplasmosis
- Cryptococcosis
- Progressive multifocal leukoencephalopathy (JC virus)
- Cytomegalovirus
- HSV, VZV in disseminated infections

Toxoplasmosis


- protozoa with complicated life-cycle
- brain abscess, mostly in cortex and gray nuclei
- acute lesion: central necrosis, mixed inflammatory reaction, macrophages („soap bubble“), toxoplasma pseudocyst
- chronic lesion: cystic space with macrophages, hemosiderin


Toxoplasmosis




Copy

Toxoplasma encephalitis





Toxoplasma encephalitis





Toxoplasma encephalitis

Cryptococcus

- fungus, PAS+ capsule
- ~ 10% of AIDS patients, now ↑ diagnosis in Africa – prophylaxis started
- meningitis mostly


Cryptococcal meningitis





Cryptococcal meningitis

Cryptococcal meningitis



PML: progressive multifocal leukoencephalopathy – demyelination






Copy

LUNG INFECTIONS

- Pneumocystis
- Candidiasis, histoplasmosis, coccidiomycosis
- CMV (+ in combination)
- TBC
- Toxoplasmosis
- Nocardiosis

LUNG INFECTIONS


- **Common, diffuse infiltrates:** CMV, Pneumocystis, drug reaction
- **Common, focal infiltrates:** Mycobacterium tbc, mycobacterium avium-intracellulare (MAC), G-rods, Staph. aureus, Aspergillus, Candida, malignant tumor
- **Uncommon, diffuse infiltrates:** bacteria, Aspergillus, Cryptococcus, malignant tumor
- **Common, diffuse infiltrates:** Cryptococcus, Mucor, Pneumocystis, Legionella




Copy

Pneumocystis pneumonia

Pneumocystis pneumonia




Pneumocystis pneumonia



TBC

- early in the course of HIV infection
- reactivation/reinfection
- pulmonary and/or disseminated
- multiple and/or highly resistant mycobacteria
- problems in combination therapy (HIV + TBC)

TBC



Invasive fungal infections

- *Aspergillus* spp (esp. *A. fumigatus*)

- Epidemiology


- Widespread, grows on rotting vegetation. Spores commonly present in air. Immunosuppression - important predisposing factor.

- Clinical presentations in HIV/AIDS


- Aspergilloma (fungal ball) - develops in cavities (lungs, sinuses, less common).

- Invasive disease - tissue destruction, pneumonia.

Fungal colony



Aspergillus pneumonia - angioinvasion



Histoplasmosis


- macrophagic intracellular parasite - fungus
- clinical presentation + morphology ~tbc
- variable course: localized/self limited – coin lesion in the lung
- chronic progressive similar to tbc
- localized extrapulmonary (mediastinum, liver, adrenals, meninges)
- disseminated in immunocompromised

Histoplasmosis

Morphology


- epithelioid cell granuloma + caseous necrosis, cavities – fibrosis – calcification
- in immunodeficient – no granuloma; accumulation of macrophages with fungal yeasts
- dg. – identification of fungal bodies (x tbc, coccidiomycosis), culture, Ab

Histoplasmosis




Copy


Coccidioidomycosis



Coccidiomyces



Interstitial pneumonia - viral




GIT INFECTIONS


Very common, persistent diarrhea


- Cryptosporidiosis, isosporidiosis (protozoa; watery diarrhea, major fluid loss; dg.- oocysts in the stool)
- Atypical mycobacteriosis (M. avium-intracellulare complex)
- Salmonella, Shigella
- CMV

Erosive gastritis



Haemorrhagic colitis






CRYPTOSPORIDIUM

copy


CMV colitis




CMV colitis



Protozoan colitis (amoebiasis)



Protozoan colitis (amoebiasis)




Mycobacterium avium complex (MAC) enteritis




copy

MAC enteritis




copy

MAC enteritis



Bacterial thrombus



HIV + hepatitis co-infection

- Common coinfection of HIV + HBV and/or HCV
- ↑ acute HCV in HIV infected
- accelerated progression of chronic hepatitis to cirrhosis + liver failure
- problems in HAART / HCV drug interaction and toxicity
- value of the transplantation?

SKIN + ORAL INFECTIONS

- Chronic, relapsing, non-healing
- Commonly ulcers
- EBV + HIV – oral hairy leukoplakia
- Candida
- HSV, VZV

Oral lesions

- Oral lesions due and according to the rate of loss of T-helper cells.
- Oral lesions - prominent features of AIDS and HIV infection.
- Early studies: approximately 90% of HIV+ patients will present with at least one oral lesion in the course of their illness.
- Current studies report the prevalence of oral lesions has significantly declined (HAART)

Oral hairy leukoplakia

- Associated with chronic shedding of EBV in the oral cavity.
- Presentation: Poorly demarcated, corrugated, white plaques on lateral aspect of tongue.
- Unlike thrush, cannot be removed by scraping.
- Occurs with immunosuppression (esp. AIDS) and warrants HIV workup.
- Diagnosis by microscopy and in situ hybridization
- Management includes establishing diagnosis and treating immunosuppression.

HIV/AIDS oral-pharyngeal syndromes

- Interferes with oral hygiene
 - More oral pharyngeal pathology
- Interferes with nutritional intake
 - Wasting syndrome
- **HIV treatment compliance** may be impacted by oral pain, xerostomia, dysphagia
- Psychosocial dimensions
 - Avoidance of social contact due to facial appearance
 - Depressive effects of persistent oral pain

Oral lesions strongly assoc. with HIV

- Candidiasis – erythematous, hyperplastic, pseudomembranous
- Hairy leukoplakia (EBV)
- HIV-associated periodontal disease – necrotizing ulcerative gingivitis, HIV periodontitis, necrotizing stomatitis
- Kaposi's sarcoma (HHV-8)
- Non-Hodgkin's malignant lymphoma (EBV)


HIV/AIDS oropharyngeal syndromes – most common

- Candidiasis 28%-75%
- Necrotizing gingivitis
- HSV, CMV, HIV, EBV ulcers
- Recurrent aphthous ulcers
- Zalcitabine ulcers
- Kaposi's sarcoma
- Dental abscesses

Oral hairy leukoplakia




Oral hairy leukoplakia



HSV in immunocompromised




Hard palate HSV in AIDS



Copyright © 2003, Elsevier Science (USA). All rights reserved.


HSV



CMV Ulcerations



Oral-pharyngeal candidiasis




HIV-associated neoplasia

- HHV-8: Kaposi sarcoma
- EBV: non-Hodgkin's malignant lymphoma, primary brain ML
- HPV: aggressive anal, cervical squamous cell carcinoma
- with HAART: general increased risk of malignancy

Human herpes virus 8

- HHV-8 is found to be associated with Kaposi sarcoma in virtually all cases.
 - Includes AIDS, post-transplant, African and Mediterranean cases.
 - HHV-8 is found in KS lesions, saliva, blood and semen of infected individuals.
- Associated with body cavity based B-cell lymphoma.
- Lesions on mucosal membranes possible, usually starts on skin.


Kaposi sarcoma




Kaposi sarcoma




Kaposi sarcoma



Kaposi sarcoma



Kaposi sarcoma + CMV colitis



copy

HIV lymphoma


- Solitary lump or nodule, swelling, nonhealing ulcer
- The swelling may be ulcerated or may be covered with intact, normal-appearing mucosa.
- Usually painful, rapid growth.
- Common association with EBV
- Several histopathologic types, atypical localization

HIV lymphoma.




Copyright © 2003, Elsevier Science (USA). All rights reserved.

Primary brain malignant lymphoma




Primary brain malignant lymphoma




Human papilloma viruses


- Human papilloma virus lesions appear most commonly in immunocompromised individuals.
- Diagnosis based on history, clinical appearance, and biopsy.
- Common in early HIV infection.
- Spiky warts, raised, cauliflower-like appearance.



HPV – koilocytosis - LSIL



HPV - immunohistochemistry



Invasive squamous cell carcinoma

Invasive cervical carcinoma



Cervical squamous cell carcinoma

