MUNI MED

# RAT Module (EN)

Instructions for students in SIMUportfolio platform

#### **Module RAT – before running the test**

The test is available in the RAT Execute section.

MUNI Med	≡	🖙 🗘 🖞 冊 RAT Student 🕛
	RAT Execute	
슈 Portfolio	☐ 2022-06-16 12:00	the test can only be started after it
RAT      Execute	TZKM: RAT test 1 ● 108: seminary room	has been started by the teacher (facilitator)
	L iRAT	COMPLETE THE TEST PREVIEW
	<b>:≗:</b> tRAT	COMPLETE AS A GROUP PREVIEW

## **Running the iRAT test**

To fill out, choose "complete the test".

=	🖙 🗘 🖞 🌐 RAT Student 🕛
RAT Execute	
<ul> <li>☐ 2022-06-16 12:00</li> <li>TZKM: RAT test 1</li> <li>108: seminary room</li> </ul>	
	COMPLETE THE TEST PREVIEW
≠≛t tRAT	COMPLETE AS A GROUP PREVIEW
	E   2022-06-16 12:00   TZKM: RAT test 1   108: seminary room

#### Start the iRAT test

After starting the test, first select the group (table number), then confirm the group.

MUNI Med	≡	🖙 ¢ 1	error RAT Student
	IRAT TZKM: RAT test 1		BACK FINISH
<ul> <li>Portfolio</li> <li>RAT ✓</li> </ul>	Select your group         Choose a group         Excorpská UNIE         Evropská strukturální a investiční fondy         Operativní program Výzkum, vývoj a vzdělávaň		V

## **Completing the iRAT test**

#### Complete the test within the set limit.

MUNT	🗮 💭 🗘 🖞 🜐 RAT Student 🕛
MED	Lirat TZKM: RAT test BACK FINISH
슈 Portfolio	Test: RAT Student (rat.student)
🖡 RAT 🗸 🗸	1. Which of the following statement is NOT true about Haemophilia A?
	Selected answer B
	A If the woman is a carrier, her son has a 50 % of having haemophilia and her daughter has a 50 % chance of being a carrier.
	• B If the man has Haemophilia, all his daughters will be carriers and all his sons will healthy (meaning not having haemophilia).
	C The amount of factor VIII deficiency correlates with the severity of haemorrhage.
	D Synthetic vasopressin is a therapeutic option used to elevate factor VIII levels. It is usually administered to patients with a mild form of haemophilia.
	E With Haemophilia A prothrombin time (PT) is expected to be prolonged.
	2. Which of the following factors will NOT be deficient in severe liver impairment?


## **Ending the iRAT test**

To send the answers, select the "finish" button and confirm sending.

MUNI	🚍 🖓 🗘 🖞 🜐 RAT Student 🕛
MED	IRAT TZKM: RAT test BACK FINISH Horig arter the situ dose subcutaneous administration shall we take the sample:
<b>命</b> Portfolio	Vybraná odpověď D
RAT V	• A aPTT, 25-30s, 4 hours after administration
	B antiXa 0,6-1,0, 3 h Submit answers?
	C anti Xa 10-20, no Yes No
	D aPTT, 60 seconds and more, immediately after the 3 rd dose
	Can not be routinely monitor in subcutaneous administration
	10. What is the initial treatment of haemodynamically stable patient with a highly suspected or a confirmed PE?
	Vybraná odpověď A
	• A Embolectomy

## **Running the tRAT test**

The tRAT test is completed for the entire group by the 1st student who chose the option "complete the test for the group".



#### **Notes**

- you complete the tRAT test similarly to the iRAT (after being started by the facilitator)
- both individual and team answers are available during the test