




14

Limbic system

Limbic system

Limbus = border






Voluntary

Somatic nervous system Inputs — mainly from outer environment Control – skeletal muscle

Automatic


Autonomic nervus system Inputs — mainly inner environment Control – smooth/cardiac m., glands






Voluntary Fornix Pineal gland Corpus callosum COMPONENTS IN THE CEREBRUM Potencial conflict COMPONENTS IN - Cingulate gyrus THE DIENCEPHALON Parahippocampal Anterior group of thalamic nuclei gyrus - Hippocampus Hypothalamus -Mamillary body-Automatic Amygdaloid body













Limbic system

Limbus = border




Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA






Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA





Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA





Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA

Arousal type 1 (somatic)

Arousal type 2 (limbic)

ARAS (ascendent retikulation activation system)

- Effect of stimulation
 - Habituation
 - Minimal activation of "reward/punishing" system

- Effect of stimulation
 - Minimal habituation
 - Strong activation of "reward/ punishing" system
 - Central gray area –CGA negative
 - Ventral tegmental area VTA positive



Arousal type 1 (somatic)

Arousal type 2 (limbic)

ARAS (ascendent retikulation activation system)

- Effect of stimulation
 - Habituation
 - Minimal activation of "reward/punishing" system

- Ascendent connections
 - Somatosensetivity, visual s., auditory s., vestibular s., cerebellum
- Descendent connections
 - Neocortex, corpus striatum, thalamus

- Effect of stimulation
 - Minimal habituation
 - Strong activation of "reward/ punishing" system
 - Central gray area –CGA negative
 - Ventral tegmental area VTA positive
- Ascendent connections
 - Mainly viscerosenzitivity, pain

- Descendent connections
 - Hypothalamus and other limbic areas, amygdala



Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA

Arousal type 1 (somatic) Effect of stimulation ARAS (ascendent retikulation activation) >Increased EEG activity > Activation of sympathetic nervous system Cooperation of both systems is a key to maintaining consciousness (through neuromodulation)

mections

neocortex, corpus striatum, thalamus


- Descendent connections
 - Hypothalamus and other limbic areas, amygdala




Acetylcholine

- Nucleus basalis (Meynerti) abd other nuclei
- Nicotin receptors
- Muscarin receptors

- Sleep/wake regulation
- Cognitive functions
- Behavior
- Emotions








Norepinefrine


- Locus coeruleus
- Nuclei raphe caudalis
- Vigilance
- Responsiveness to unexpected stimuli
- Memory
- Learning





Dopamine

- Nigrostriatal system
 - Movement
 - Sensory stimuli
- Ventrotegmentno-mesolimbicfrontal system
 - Reward
 - Cognitive function
 - Emotional behavior
- Tubero-infundibular system
 - Hypotalamic-pituatory regulation
- D1 receptors excitatory
- D2 receptors inhibitory




http://www.slideshare.net/drpsdeb/presentations



Serotonin


- Nuclei raphe rostralis
- Nuclei raphe caudalis
- Anxiety/relaxation
- Impulsive behavior
- Sleep




http://www.slideshare.net/drpsdeb/presentations




Sleep and wakefulness









Sleep






Sleep









LIGHT NON-REM SLEEP

- > Stages 1-2
- May drift in and out of sleep several times at the start
- Easy to wake up, disturbs easily

DEEP NON-REM SLEEP

- Stages 3-4
- Difficult to wake up
- Sleep inertia when woken

REM

- Eyes move rapidly under closed eyelids
- Most dreaming occurs here
- Brain is active, muscles are relaxed
- Can't move voluntarily signals from the brain to the postural muscles are blocked



http://www.dailymail.co.uk/sciencetech/article-3042230/Sleeping-habits-world-

revealed-wakes-grumpy-China-best-quality shut-eye-South-Africa-wakes-earliest.html


Sleep and wakefulness

Brainstem nuclei responsible	Neurotransmitter	Activity state
WAKEFULNESS		
Cholinergic nuclei of pons-midbrain junction	Acetylcholine	Active
Locus coeruleus Raphe nuclei	Norepinephrine	Active
	Serotonin	Active
NON-REM SLEEP		
Cholinergic nuclei of pons-midbrain junction	Acetylcholine	Decreased
Locus coeruleus Raphe nuclei	Norepinephrine	Decreased
	Serotonin	Decreased
REM SLEEP ON		
Cholinergic nuclei of pons-midbrain junction	Acetylcholine	Active
Raphe nuclei	Serotonin	Inactive
REM SLEEP OFF		
Locus coeruleus	Norepinephrine	Active




Hypothalamus

- Key center of autonomic regulations and coordination
- Integration of the information from inner and outer environment
- Behavioral modulation
- Regulation of autonomic nervous system
- Maintenance of homeostasis



http://biology.about.com/od/anatomy/p/Hypo thalamus.htm






Hypothalamus

Key center of autonomic regulations and coordination


enviro Biological clock - circadian /seasonal activity ✓ Autonomic nervous system regulation ✓ Endocrine system regulation √Food and water intake regulation ✓ Regulation of body temperature Behavid √"Immediate" behavior regulation (e.g. when hunger) Regulati

√ "Long-term" behavior regulation (e.g. maternal beh.) nervous ✓ Instinctive behavior regulation (e.g. sexuality) Maintena pituitary



Influence of hypothalamus on neocortex

- Via neuromodulating systems
 - Consciuosness (see above)
 - Mood
- Via thalamus
 - Via nucleus mediodorsalis to orbitofrontal cortex (influence on decision making)
 - Influence gating function of other thalamic nuclei
- Papez circuit




cortex

Orbitofrontal







Copyright @ 2007 Wolters Kluwer Health | Lippincott Williams & Wilkins


http://www.slideshare.net/drsunilsuthar/neurobiology-of-emotion



Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Acces sed). License: Creative Commons BY-NC-SA



Prof. Gerald Schneider



Copyright @ 2007 Wolters Kluwer Health | Lippincott Williams & Wilkins

 $\mathbf{f}\mathbf{x} = \text{fornix bundle}$ Association areas (neocortex) Cingulate cortex Paralimbic areas, Retina -> Pretectal Laterodorsal nuclei nucleus entorhinal area of thalamus Subiculum Mammillary (Hypothalamus ... <u>fx</u> Hippocampusı → Septal area (Ach) bodies Dentate gyrus Hippocampal formation

mt = mammillothalamic tract

Courtesy of MIT Press, Used with permission.

Schneider, G. E. Brain Structure and its Origins: In the Development and in Evolution of Behavior and the Mind. MIT Press, 2014, ISBN: 9780262026734.




Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Acces sed). License: Creative Commons BY-NC-SA



Prof. Gerald Schneider

particular place



Copyright © 2007 Wolters Kluwer Health | Lippincott Williams & Wilkins

 $\mathbf{f}\mathbf{x} = \text{fornix bundle}$ Association areas (neocortex) Cingulate cortex Paralimbic areas. Retina -> Pretectal Laterodorsal nuclei nucleus entorhinal area of thalamus Spatial orientation and emotions associated with Subiculum Mammillary (Hypothalamus ... <u>fx</u> Hippocampusı →Septal nuclei area (Ach) bodies Dentate gyrus Hippocampal formation Courtesy of MIT Press, Used with permission. Schneider, G. E. Brain Structure and its Origins: In the Development and in Evolution of Behavior and the Mind. MIT Press, 2014, ISBN: 9780262026734.

 $\mathbf{mt} = \text{mammillothalamic tract}$

Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Acces sed). License: Creative Commons BY-NC-SA



Prof. Gerald Schneider

- Suggestion: the ascending axons of this circuit are continuously activating memories of places that lie ahead, in the direction indicated by the current
 Thus, decisions about direction of locomotion are direction of the head. influenced by memories of those places, including their good or bad values. • Axons in the Papez circuit are of more than one type.
- Only the ones signaling head direction have been characterized.
- What is the hippocampus sending to other parts of the hypothalamus? It may alter motivational levels according to remembered information about locations in the current frame of reference.

mt = mammillothalamic tract $\mathbf{f}\mathbf{x} = \text{fornix bundle}$ Association areas (neocortex) Cingulate cortex Retina -> Pretectal Paralimbic areas. Laterodorsal nuclei entorhinal area of thalamu Subiculum <u>fx</u> Hippocampus Mammillary Hypothalamus Septal bodies Dentate gyrus Hippocampal formation Courtesy of MIT Press, Used with permission.





Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Acces sed). License:Creative Commons BY-NC-SA



Prof. Gerald Schneider

- Origins of endbrain: Structures underlying olfaction
- Two major links between olfactory system and the motor systems of the midbrain
 - Through the ventral endbrain, which became corpus striatum and basal forebrain (including much of the septal area)
 - Outputs to hypothalamus, (epithalamus, subthalamus), midbrain
 - These outputs affected locomotion and orienting movements
 - The links were plastic, so <u>habits</u> were formed according to rewarding effects mediated, e.g., by taste effects.
 - 2) Through the medial part of the dorsal endbrain, which became medial pallium—the hippocampal formation
 - Outputs to ventral striatum, hypothalamus, epithalamus
 - The links were plastic, but the "habits" formed were different: The association of place with good or bad consequences of approach.

mt = mammillothalamic tract $\mathbf{f}\mathbf{x} = \text{fornix bundle}$ Association areas (neocortex) Cingulate cortex Paralimbic areas. Retina -> Pretectal Laterodorsal nuclei nucleus entorhinal area Anterior hucle of thalamus Subiculum <u>fx</u> Hippocampus Mammillary ← Hypothalamus ← Septal nuclei (Ach) bodies area Dentate gyrus Hippocampal formation Courtesy of MIT Press, Used with permission. Schneider, G. E. Brain Structure and its Origins: In the Development and in

Evolution of Behavior and the Mind. MIT Press, 2014, ISBN: 9780262026734.



Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts oriented... Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Ac sed). License: Creative Commons BY-NC-SA



Prof. Gerald Schneider

Object (

Location

oriented...

- Origins of endbrain: Structures underlying olfaction
- Two major links between olfactory system and the motor systems of the midbrain
 - Through the ventral endbrain, which became corpus striatum and basal forebrain (including much of the septal area)
 - Outputs to hypothalamus, (epithalamus, subthalamus), midbrain These outputs affected locomotion and orienting movements The links were plastic, so habits were formed according to rewarding effects mediated, e.g., by taste effects.
 - Through the medial part of the dorsal endbrain, which became medial pallium—the hippocampal formation
 - Outputs to ventral striatum, hypothalamus, epithalamus The links were plastic, but the "habits" formed were different: The association of place with good or bad consequences of approach.

 $\mathbf{f}\mathbf{x} = \text{fornix bundle}$ Association areas (neocortex) Cingulate cortex Paralimbic areas. Retina -> Pretectal Laterodorsal nuclei nucleus entorhinal area Anterior hucl of thalamus Subiculum <u>fx</u> Hippocampus Mammillary Hypothalamus →Septal nuclei (Ach) bodies area Dentate gyrus Hippocampal formation Courtesy of MIT Press, Used with permission. Schneider, G. E. Brain Structure and its Origins: In the Development and in

Evolution of Behavior and the Mind. MIT Press, 2014, ISBN: 9780262026734.

mt = mammillothalamic tract

Spatial orientation and emotions associated with particular place

Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA




Prof. Gerald Schneider



Evolution of corpus striatum:

basic outline of a story

1. Beginnings: a link between olfactory inputs and motor control: The link becomes "Ventral striatum". It was a modifiable link (capable of experience-induced change).



3. Early expansions of endbrain: striatal and pallial.

4. Pre-mammalian & then mammalian expansions of cortex and striatum: For the striatum, the earlier outputs and inputs remain as connections with neocortex expand.

Figure 1. Postulated beginnings in primitive chordates




Figure 2. Other inputs reached the striatum




Figure 3. Early expansion of striatal and adjacent "limbic" areas




Figure 4. Pre-mammalian, and then mammalian expansions



Courtesy of MIT Press. Used with permission. Schneider, G. E. Brain structure and its Origins: In the Development and in Evolution of Behavior and the Mind. MIT Press, 2014. ISBN: 9780262026734.

Working/Short term memory - "RAM" Long term memory
– "Hard disk"



- Connections of striatum and hippocampus are plastic
- Plasticity is a base of learning
- Learning is a forming of long-term memory

Working/Short term memory - "RAM" Long term memory – "Hard disk"



- Connections of striatum and hippocampus are plastic
- Plasticity is a base of learning
- Learning is a forming of long- term memory
- Procedural memory (implicit)
 - Based on striatum
 - Habitual learning motor skills, but also social habits
 - "Construction of the algorithms"



- Connections of striatum and hippocampus are plastic
- Plasticity is a base of learning
- Learning is a forming of long- term memory
- Procedural memory (implicit)
 - Based on striatum
 - Habitual learning motor skills, but also social habits
 - "Construction of the algorithms"
- Declarative memory (explicit)
 - Based on hippocampus
 - Explicit information is stored and later recollected
 - "Construction of the maps (relationships)" spatial or abstract



- Connections of striatum and hippocampus are plastic
- Plasticity is a base of learning
- Learning is a forming of long- term memory
- Procedural memory (implicit)
 - Based on striatum
 - Habitual learning motor skills, but also social habits
 - "Construction of the algorithms"
- Declarative memory (explicit)
 - Based on hippocampus
 - Explicit information is stored and later recollected
 - "Construction of the maps (relationships)" spatial or abstract

Object oriented:

Can I eat it and how to eat it?

Location oriented:

Where am I and what has happened here?




Cognitive maps

Type of mental representation which serves an individual to acquire, code, store, recall, and decode information about the relative locations and attributes of phenomena in their everyday or metaphorical spatial environment.

Wikipedia

- ✓ Unique
- ✓ Egocentric, but multiple

Later generalized to refer to a kind of semantic network / representing an individual's personal knowledge or schemas.






Amygdala

Corticomedial: Inputs from olfactory bulbs, hypothalamus & lateral amygdala; outputs to hypothalamus, amygdala, ANS

Basolateral: Inputs from thalamus, neocortex, hippocampus; outputs to prefrontal cortex, ventral striatum, other amygdala nuclei

Central: Intra-amygdalar inputs; outputs through stria terminalis (see later slides)

- Connections to all major cortical and subcortica lstructures
- Modiffied corpus striatum
- Plasticity memory formation






Amygdala


Corticomedial: Inputs from olfactory bulbs, hypothalamus & lateral amygdala; outputs to hypothalamus, amygdala, ANS

Basolateral: Inputs from thalamus, neocortex, hippocampus; outputs to prefrontal cortex, ventral striatum, other amygdala nuclei


Central: Intra-amygdalar inputs; outputs through stria terminalis (see later slides)

- Connections to all major cortical and subcortica lstructures
- Modiffied corpus striatum
- Plasticity memory formation
- "Influence of information from outer environment on limbic system"
- "Amygdala hijack"
- "Affective tags"
 - Both possitive and negative
 - Higher responsiveness to negative





Gerald Schneider. 9.14 Brain Structure and Its Origins, Spring 2014. (Massachusetts Institute of Technology: MIT OpenCourseWare), http://ocw.mit.edu (Accessed). License:Creative Commons BY-NC-SA





MUNI MED

83. The importance of limbic system and brief characterization of basic functions – somatic and limbic arousal systems, sleep and wakefulness

- Concept, definition and structures of limbic system
 - Integration of information from inner and outer environemnt
 - Hypothalamus is a central structure...
- Somatic vs. limbic arousal system
- Habituation, association with reward punishing system, connections...

- Sleep/wakefulness cooperation of somatic and limbic activation system via neuromodulation
- Phases of sleep, basic EEG characteristics

$\mathsf{M} \mathsf{U} \mathsf{N} \mathsf{I}$ $M \in D$

84. The importance of limbic system and brief characterization of basic functions – learning and memory, the influence of hypothalamus on neocortex, the role of amygdala

- Concept, definition and structures of limbic system
 - Integration of information from inner and outer environemnt
 - Hypothalamus is a central structure...
 - Brief overview of hypothalamic functions
 - Influence of hypothalamus on neocortex

- Learning and memory
 - Learning is based on plasticity, learning is forming of long-term memory
 - Explicit memory hippocampus
 - Implicit memory striatum
- Amygdala
 - Influence of information from outside (neocortex) on limbic system
 - Amygdal hijack, affective tags

#