Otázky k praktické státní zkoušce konané v předmětu ZL-KZL094c simulovaná výuka

- 1. Caries lesion on mesial surface of maxillary molar **16 nebo 26**, crista obliqua is affected. Preparation for composite, making filling.
- 2. Caries lesion on distal surface of maxillary molar **16 nebo 26**, crista obliqua is affected. Preparation for composite, making filling.
- 3. Caries lesion on mesial and distal surface of maxillary molar **16 nebo 26**, crista obliqua is intact. Preparation for composite, making filling.
- 4. Caries lesion on mesial and distal surface of maxillary molar **16 nebo 26**, crista obliqua is affected. Preparation for composite, making filling.
- 5. Caries lesion on distal surface of mandibulary molar **36 or 46**, dental caries is also on the occlusal surface. Preparation for composite, making filling.
- 6. Caries lesion on mesial surface of mandibulary molar **36 or 46**, dental caries is also on the occlusal surface. Preparation for composite, making filling.
- 7. Caries lesion on mesial and distal surface of mandibulary molar **36 or 46**, fissures are intact (no lesion on occlusal surface). Preparation for composite, making filling.
- 8. Caries lesion on mesial and distal surface of mandibulary premolar **35 or 45**, dental caries is also on the occlusal surface. Preparation for composite, making filling.
- 9. Caries lesion on mesial and distal surface of maxillary premolar 15 or 25, dental caries is also on the occlusal surface. Preparation for composite, making filling.
- 10. Caries lesion on central maxillary incisor **11 or 21**, on mesial and distal surface, the incisal edge on distal side is affected. Preparation and making composite filling.
- 11. Caries lesion on central maxillary incisor **11 or 21**, one mesial and distal side, incisal edge is not affected. There is also cervical lesion in enamel. Suggestion of suitable materials, preparation and filling.
- 12. Caries lesion on central maxillary incisor **11 or 21**, on mesial surface, the incisal edge is affected.
- 13. Caries lesion on distal surface of maxillary molar **16 nebo 26**, crista obliqua is intact. There is also cervical caries out of enamel. Suggestion of suitable materials, preparation and filling.
- 14. Caries lesion on distal surface of maxillary molar **16 nebo 26**, crista obliqua is intact. There is also cervical caries in enamel. Suggestion of suitable materials, preparation and filling.
- 15. Caries lesion on distal surface of maxillary molar **16 nebo 26**, crista obliqua is intact. There is also a small lesion in mesial part of occlusal surface. Preparation for composite, making filling.
- 16. Caries lesion on mesial surface of mandibulary premolar **35 or 45**, there i salso v-shaped defect. Suggestion of suitable materials, preparation and filling.
- 17. Traumatic defect on central maxillary incisor **11** or **21**, incisal edge and 1/3 of the crown are lost. Preparation and making composite reconstruction.
- 18. Caries lesion on the mesial surface of central maxillary incisor **11 or 21.** There is also cervical lesion located partly in enamel, partly in cementum. Suggestion of suitable materials, preparation and filling.
- 19. Caries lesion on mesial and distal surface of maxillary premolar 14 **or 24**,dental caries is also on the occlusal surface. Preparation for composite, making filling.
- 20. Caries lesion on mesial and distal surface of maxillary premolar **15 or 25**, there is no caries on occlusal surface. Preparation for composite, making filling.