


Collection of exercises from ELEMENTARY
GEOMETRY for the study of teaching at
primary school

Leni Lvovská

October 2019



*Geometry has two treasures: a pythagorean theorem and a golden section.
The first one is worth gold, the second one is more precious stone.*

Johannes Kepler (1571 - 1630)[10]


Introduction

This collection of elementary geometry problems was developed as a supporting material to the geometry textbooks for the future elementary school teachers. These texts namely contain only a limited number of exercises and no solved tasks. This booklet offers the students a number of solved tasks as well as another set of exercises. At the same time, it follows the current trend of inter-subject connections and in the provided tasks and examples shows how geometry is related to the other subjects as well as to the world around us.

Many tasks work with the magnetic kit Geomag. If you do not have it, these tasks can be demonstrated using skewers and balls of modeling. For the creation of illustrations, the GeoGebra software was used. It is therefore easy to use the GeoGebra tutorial software directly in the classroom or on a standalone task. There are direct references to selected dynamic applets and stepped constructions for specific constructions.

This text was developed with the support of the project MUNI / FR / 1193/2018, Innovation of four subjects Geometry for Teachers of the 1st level of the elementary school with Geomag kit and Geogebra educational software at the Faculty of Education, Masaryk University in Brno.


Many thanks to Helena Durnová for the preparation of the English version of this text and Pavel Kříž for the support of the typesetting in the \LaTeX system.



A regular hexagon or cube?

Exercises 1.2. How did geometry begin to form in the distant past? (Formulate the answer in several sentences.)

Exercises 1.3. What do you know about the books we call Euclid's *Elements*?



Exercises 1.4. Correctly associate branches of geometry with the names of important mathematicians, who worked in them:

- René Descartes (1596 – 1650),
 - Johann Carl Friedrich Gauss (1777 – 1855),
 - Georg Friedrich Bernhard Riemann (1826 – 1866),
 - David Hilbert (1862 – 1943).
- a) German mathematician and physicist. He was interested in geometry, mathematical analysis, number theory, astronomy, electrostatics, geodesy and optics. He strongly influenced most of these fields of knowledge. He stood at the birth of non-Euclidean geometry.
 - b) His work *La Géométrie* is often considered the beginning of analytic geometry as a science.
 - c) German mathematician, who in his work *Foundations of Geometry* constructed discipline currently called Euclidean geometry, he created the system of axioms of Euclidean geometry.
 - d) German mathematician, who contributed significantly to the development of mathematical analysis and differential geometry. Algebraic geometry and complex surface theory were also developed on the basis of his ideas, which became the core of differential geometry on manifolds and topology.

Solution: René Descartes (b), Johann Carl Friedrich Gauss (a), Georg Friedrich Bernhard Riemann (d), David Hilbert (c).

Exercises 1.5. Explain the difference between an axiom and a mathematical theorem. Give an example of an axiom and a mathematical theorem.

2 Basic geometric formations and their properties

Example 2.1. Which geometric formations can arise as the intersection of two half-lines that are lie on the same line? Show and describe.

Solution: Point, line segment, line, half line.

Example 2.2. Investigate all possible relative positions of three different lines lying in one plane. Show and describe.

Řešení: Let's denote the lines a, b, c . Then the following situations can occur:

- a) all lines are mutually parallel, ie. $a \cap b = \emptyset \wedge b \cap c = \emptyset$,
- b) two lines are parallel and the third line is parallel to them, e.g.

$$a \cap b = \emptyset \wedge a \cap c = X \wedge B \cap C = Y$$

- c) all lines are mutually parallel and pass through a single common point, $a \cap b \cap C = P$,
- d) all lines are mutually parallel and intersect at different points.

Exercises 2.3. Draw a line AB . On the line AB mark:

- a) C , so that A is between C and B ,
- b) D , so that B is between A and D ,
- c) a point P that does not lie on the AB line but lies on the AD line.

Exercises 2.4. Draw line KL . Select D between KL , and mark:

- a) R , so that K is between R and L ,
- b) S , so that L is between K and S ,
- c) T , so that S is between L, T .

Decide which statement is true:

- 1) $S \in \mapsto KL$,
- 2) $\mapsto RS \cap \mapsto KL = KL$,

3) $\mapsto RD \cap ST = \emptyset$,

4) $R \in \leftrightarrow KL$,

Exercises 2.5. There is a line p and a point A that does not lie on it. Draw:

- a) a point M that belongs to the $\mapsto pA$,
- b) a point P that lies in both halves defined by p ,
- c) a point N that lies in the half-plane opposite to $\mapsto pA$.

Exercises 2.6. Three different points A, B, C are given.

- a) How many line segments, half-lines, and lines are determined by these points? How do these numbers depend on the position of the points given?
- b) Which point sets can be the intersection of two of these line segments (half-lines, lines)?

Show and discuss.

Exercises 2.7. Let point R lies between P, Q . From half-lines PR, PQ, RP, RQ, QR, QP choose pairs of half-lines that: a) coincide, b) are opposite, c) one is part of the other, d) their intersection is a line segment.

Exercises 2.8. Determine what shapes may arise as an intersection of:

- a) a line segment and a half-plane,
- b) a half-line and a half-plane,
- c) a line and a half-plane.

For all cases, consider the situation in a single plane. Show and describe.

Exercises 2.9. There are n straight lines in the plane, of which no two intersect and no three meet at the same point. How many significant intersections of these lines are there?

Example 2.10. How many different lines are determined by n points that lie in one plane and no three lying on one straight line?

Solution: For a single point, the task is meaningless. Let's outline the situation for some finite number of points: for two points there will be one straight line, for three points just three lines, four points will determine six lines, five points will be ten lines, etc. Now we can do the following: from each point we lead a line to $(n - 1)$ points, but in this way I count them each line twice. The result is:

$$\frac{n(n - 1)}{2}.$$

Exercises 2.11. In the plane there are n lines, two of which intersect and no three of them meet the same point. How many intersections there are?

Exercises 2.12. Determine what shapes may arise as an intersection of two half-planes. Consider the situation in a single plane.

Exercises 2.13. Select points A, B inside one half-plane, which is determined by the line p . Inside the opposite half-plane, select C, D so that the lines AB and CD are parallel to the line p . On line AB select M , on line CD select N . How must the points M, N be chosen so that the line segment MN contains a point of line p lying between M and N ?


Example 2.14. Construct a cuboid of $ABCDEFGH$ (using GeoMag or using skewers and plasticine).

A) Determine all incident lines with cuboid edges that are with BC :

- parallel,
- intersecting,
- skew.

B) Using the points of the cuboid, you list three planes that form a bundle of planes and write down the intersection of these three planes.

Řešení:



- parallel: $\leftrightarrow AD, \leftrightarrow EF, \leftrightarrow HG$
- intersecting: $\leftrightarrow AB, \leftrightarrow EB, \leftrightarrow DC, \leftrightarrow CF$
- skew: $\leftrightarrow EH, \leftrightarrow FG, \leftrightarrow AH, \leftrightarrow DG$

A bundle of planes consists of planes $\leftrightarrow ABC, \leftrightarrow ABE$ a $\leftrightarrow AF$:

$$\leftrightarrow ABC \cap \leftrightarrow ABE \cap \leftrightarrow ABF = \leftrightarrow AB.$$


Example 2.15. Construct a regular tetrahedral pyramid $ABCDV$ (using GeoMag kit or skewers and plasticine).

A) Determine all straight lines specified by A, B, C, D, V that are:

- parallel to BC ,
- intersecting to BC ,
- skew to BC .

B) Using the pyramid points A, B, C, D, V , give an example of the three planes that make up the bunch of planes and write the intersection of the three planes.

Solution:



- parallel: $\leftrightarrow AD$,
- intersecting: $\leftrightarrow AB$, $\leftrightarrow BV$, $\leftrightarrow CV$, $\leftrightarrow CD$,
- skew $\leftrightarrow AV$, $\leftrightarrow DV$.

A bunch of planes is made up of planes $\leftrightarrow ABC$, $\leftrightarrow ABV$ a $\leftrightarrow BCV$:

$$\leftrightarrow ABC \cap \leftrightarrow ABV \cap \leftrightarrow BCV = \leftrightarrow \{B\}.$$

3 Convex and non-convex set, convex and non-convex angle

Exercises 3.1. How can we find out whether a geometrical figure is convex or non-convex? Sort geometric shapes into convex and non-convex: a line segment, line, circle, triangle, quadrilateral, pentagon, circle with hole.

Exercises 3.2. Look around and try to see the angles determined by the edges of the board or the edges of the bench, parts of the window frame, but also the angles formed, for example, the legs of the chair and the floor. Mark such angles in the illustration.



Exercises 3.3. Draw the lines $\leftrightarrow SC$ and $\leftrightarrow SD$. Mark with a red arc the convex angle $\sphericalangle CSD$ and with a blue one non-convex angle $\sphericalangle CSD$. Mark the point E of angle $\sphericalangle CSD$ and the point F of angle $\sphericalangle CSD$. Can you determine point H , which is the point of the angle $\sphericalangle CSD$ as well as of the angle $\sphericalangle CSD$?

Exercises 3.4. Draw the angle $\sphericalangle ADB$. Mark point H in it. Draw the angle $\sphericalangle ADH$. Write down all convex angles.

Exercises 3.5. Draw three lines with a common S origin. Mark one of the points A, B, C on each of the lines. Mark the curves in these angles and write them down.

Exercises 3.6. Sketch two **convex** planar formations such that their

- a) union is a convex set,
- b) union is a non-convex set,
- c) intersection is a convex set,
- d) intersection is a non-convex set.

Exercises 3.7. Sketch two **non-convex** planar formations such that their

- a) union is a convex set,
- b) union is a non-convex set,
- c) intersection is a convex set,
- d) intersection is a non-convex set.

Exercises 3.8. Sketch the following and determine whether it is a convex or non-convex set:

- a) a triangle ABC without its vertices,
- b) triangle KLM without one inner point of one side,
- c) union of the inside of any triangle and two different points of its perimeter,
- d) difference of a convex angle AVB and its arm VA ,
- e) difference of square $ABCD$ and the union of its two sides,
- f) union of the inside of square $ABCD$ and its two sides,
- g) circle.


Example 3.9. Investigate all geometric shapes that may arise as an intersection of two triangles. Show and describe.

Solution: The intersection of two triangles may be:

- A) a point, e. g. $\triangle ABC \cap \triangle EFD = \{D\}$,
- B) a line segment, e.g. $\triangle ABC \cap \triangle EFD = DC$,
- C) a triangle, e. g. $\triangle ABC \cap \triangle EFD = \triangle DMN$,
- D) a quadrilateral, e.g. $\triangle ABC \cap \triangle EFD = \text{quadrilateral } OPQR$,

E) a pentagon, e.g. $\triangle ABC \cap \triangle EFD = \text{pentagon } FSTUV$,

F) hexagon, eg. $\triangle ABC \cap \triangle EFD = \text{hexagon } KLMNOP$.




Exercises 3.10. Choose pairs of convex angles (neither full nor zero ones). Investigate which geometrical shapes can arise as the intersections of these angles. Draw and describe all cases.

Exercises 3.11. Select different non-parallel lines p, q and mark their intersection V . Select P on the line p , Q on the line, q . Define each pair of vertical and adjacent angles determined by the intersecting lines p and q using the half-planes pQ, qP and the half-planes opposite to them. Use symbolic notation.


Exercises 3.12. Model from the Geomeg kit and then draw: a) isosceles triangle, b) equilateral triangle, (c) a square; (d) a regular pentagon; (d) a regular hexagon.

Example 3.13. Model the Geomag kit with the following specifications, and then practice your imagination to solve them:


A) Move 3 equal lines (yellow bars) to form 2 large and one small triangle. The task has two solutions.




B) Remove 3 equal lines (yellow sticks) to form 3 squares. The task has two solutions.



C) Remove one line (yellow bar) to get 2 squares. The task has two solutions.



D) Move 4 identical lines (yellow bars) to create four squares again, but not all of the same size.



Solution: See at the end of the text.

Exercises 3.14. Draw a regular hexagon $ABCDEF$ with the center S and mark the pairs of angles in the picture:



- a) adjacent (not supplementary),
- b) supplementary,
- c) vertical,
- d) corresponding,
- e) alternate exterior angles,
- f) alternate interior angles.

Example 3.15. Model a regular tetrahedron $ABCD$ and then display it. Determine its intersection with the $EFGH$ halfspace if A is between E and C , B is between F and C , and G between D and C .



4 Circle, round, triangle, quadrilateral, regular polygon

Exercises 4.1. Tangram is the oldest known puzzle in the world, it comes from ancient China. It is a square divided in a thoughtful way into seven parts, from which various geometric figures, objects, animals and human figures can be assembled. Make your tangram from a square of paper according to the attached pictures:



Then build using all seven parts:


- a) triangle,
- b) parallelogram,
- c) trapezoid.

Čínští matematici, kteří se tangramem zabývali, zjistili, že ze sedmi částí tangramu lze sestavit celou sérii konvexních mnohoúhelníků:

- a) 1 triangle,
- b) 6 quadrilaterals,
- c) 2 pentagons,
- d) 4 hexagons.

So if you have mastered the geometric shapes a) - d), you can try this somewhat more difficult task.

Example 4.2. The picture shows a few well-known road signs. Answer the following questions:




- 1) Jaké geometrické útvary se nacházejí na obrázcích? (jednobodovou množinu neuvažujeme)
- 2) Construct with a ruler and compass all geometric shapes from task 1).
- 3) Use the ruler and compass to construct the centers of both circles on the first mark? What is the relationship between the two circles?
- 4) What is the content of the triangle that forms the second tag if its side is 900mm? Try to solve the problem in several ways (repeat Heron's formula).
- 5) The first tag has a diameter of 700mm. The side of the triangle on the second mark is 900mm. Which of these brands do we need more sheet metal for?
- 6) Take a picture of the tags you meet on your way to school and formulate similar questions.

Solution: 1) Line, circle, round, equilateral triangle, square, rectangle, regular octagon. 3) These are concentric circles that have a common center. This center can be found, for example, using two arbitrary different chords, taking advantage of the fact that the axis of each line that is a chord of a circle is a line passing through the center of the circle.

Example 4.3. Construct a square if its side is AB . Select claims that are false:

- All angles are equal in a square.
- Right angle is exactly one square.
- Two sides in a square must be horizontal.
- In, adjacent pages must be perpendicular to each other.
- The angle between the diagonal and the adjacent side of the square is 45^{circ} .
- Diagonals in a square form an angle of 60^{circ} .

Solution:



False statements are statements b), c), f).

Exercises 4.4. If in an isosceles triangle ABC the angle at base AB equals three times the angle at the vertex C and if the angle $\sphericalangle BAC$ at the base is divided into three equal angles (so that M, N are those points of BC for which $\sphericalangle NAB \cong \sphericalangle MAN \cong \sphericalangle CAM$), then $AB \cong AN \cong BM, AM \cong CM$. Prove.


Exercises 4.5. A point A lying outside the circle $k(S, r)$ leads the secant CD so that $AC < AD$ and $|AC| = r$. Prove that

$$\sphericalangle ASC = \frac{1}{3} \sphericalangle BSD,$$

where the point B is the point of intersection of the line AS with the circle k such that S lies between the points A, B .

Exercises 4.6. Inside the ABC triangle, select S . Prove that the sum of the lines SA, SB, SC is greater than half the sides of the triangle, ie

$$SA + SB + SC > \frac{1}{2}(AB + BC + CA). \quad (1)$$



Solution: Point S is the inner point of the triangle ABC , so there are three other triangles for which the triangular inequality holds:

for triangle ABS : $AS + BS > AB$,

for triangle ACS : $AS + CS > AC$,

for triangle BCS : $BS + CS > BC$.

Adding the right and left sides of the inequalities we get:

$$2 \cdot AS + 2 \cdot BS + 2 \cdot CS > AB + BC + AC, \quad (2)$$

thus proving inequality (1).

Example 4.7. Prove that for the sum of the centroids t_a, t_b, t_c of triangle ABC , the relation is:

$$\frac{1}{2}(a + b + c) < t_a + t_b + t_c < a + b + c. \quad (3)$$

Řešení: First we prove inequality

$$\frac{1}{2}(a + b + c) < t_a + t_b + t_c. \quad (4)$$

Let's denote A_1 center of BC , B_1 center of AC and C_1 center of AB triangle ABC . The triangular inequality follows

for triangle ABA_1 : $t_a + \frac{a}{2} > c$,

for triangle ACC_1 : $t_c + \frac{c}{2} > b$,


for triangle BCB_1 : $t_b + \frac{b}{2} > a$.

Adding the right and left sides of the inequalities we get:

$$t_a + t_b + t_c + \frac{1}{2}(a + b + c) > a + b + c, \quad (5)$$

i.e.

$$t_a + t_b + t_c > \frac{1}{2}(a + b + c). \quad (6)$$



Let us prove inequality

$$t_a + t_b + t_c < a + b + c. \quad (7)$$

Let the points A_1 , B_1 , C_1 be again the centers of the sides BC , AC and AB of the triangle. Let's construct A' so that A_1 is the center line of AA' . The quadrilateral $ABA'C$ is a parallelogram, its diagonals are halved. So $AC \cong BA'$ holds. The triangular inequality for triangle ABA' implies:

$$2t_a < b + c. \quad (8)$$

Similarly, constructing B' and C' so that B_1 is the center of the BB' line and C_1 is the center of the CC' line:

$$2t_b < a + c \quad (9)$$

and

$$2t_c < a + b. \quad (10)$$

Adding the right and left sides of the three inequalities we get:

$$2t_a + 2t_b + 2t_c < 2a + 2b + 2c, \quad (11)$$

that is, we proved inequality (7).

Example 4.8. Prove that the sum of the lines that connect the inner point P of the triangle to the endpoints of one side is less than the sum of the remaining two sides of the triangle.

Proof: For example, according to the task assignment,

$$AP + BP < AC + BC \quad (12)$$

We can now prove the argument (12). Since the P point belongs to the inside of the ABC triangle, there must be a X point on the side of BC and on the AP half-line after P . For triangles ACX and BPX we express a triangular inequality

$$\text{for triangle } ACX: AX < AC + CX,$$

$$\text{for triangle } BPX: BP < XB + PX.$$

After adding both inequalities we get:


$$AX + BP < AC + CX + BX + PX. \quad (13)$$

Expressing the line AX as the sum of the lines $AP + PX$ and considering that $CX + BX = BC$:

$$(AP + BP) + PX < AC + (CX + XB) + PX,$$

$$(AP + BP) + PX < (AC + BC) + PX$$

and thus the inequality (12) is proven.



Exercises 4.9. The o line is the axis of the AB line. The X point is any inner half-point oA . Prove that: $AX < BX$.

Exercises 4.10. Point U is the inner point of the triangle ABC . Prove that the following applies: $\sphericalangle AUB > \sphericalangle ACB$, $\sphericalangle BUC > \sphericalangle BAC$ and $\sphericalangle AUC > \sphericalangle ABC$.

Exercises 4.11. If the point X lies on the axis of the given convex angle AVB , then it has the same distances from its arms. Prove.

Exercises 4.12. If the center of gravity of the triangle coincides with its height, the triangle is isosceles. Prove.

Exercises 4.13. In the ABC triangle is $\sphericalangle BAC = \alpha = 50^\circ$, $\sphericalangle ABC = \beta = 60^\circ$, $\sphericalangle ABC$ crosses AC v point D . Sort lines AB, BC, CD, AD, AC, BD by size.

Exercises 4.14. Determine the magnitude of the inner angles of the triangle $A_1B_1C_1$, whose vertices are the intersections of the axes of the outer angles of the triangle ABC .


Exercises 4.15. It is given by an isosceles triangle ABC and a point D which is the center of its base AB . Point D leads perpendicular to the arms AC, BC triangle ABC . Their heels are labeled M, N . Prove that $\triangle DMC \cong \triangle DNC$.

Exercises 4.16. Construct a triangle ABC if three independent data are given:

- | | | |
|---------------------|-------------------------|-------------------------|
| a) c, b, t_c | b) α, c, t_c | c) a, v_a, b |
| d) a, α, v_b | e) b, c, v_a | f) α, v_b, r_v |
| g) b, γ, v_c | h) γ, v_a, v_b | i) c, v_a, v_b |
| j) a, v_a, v_b | k) γ, v_a, v_c | l) r_o, v_c, t_c |
| m) a, b, t_c | n) α, β, r_v | o) α, β, r_o |
| p) b, β, v_b | q) a, β, r_v | r) c, t_a, t_b |
| s) b, β, t_a | t) a, t_a, t_b | u) a, v_a, t_b |
| v) t_a, t_b, t_c | w) t_a, t_b, γ | z) t_a, v_a, v_b |

where r_o is the circle radius described and r_v is the circle radius inscribed with the triangle ABC . Some of these tasks can be found in the following examples.

Example 4.17. Construct a triangle ABC if given: a, α, v_b .



The construction step by step


The record of the construction:

- 1) $BC; |BC| = a$
- 2) $S_a; S_a \in BC \wedge |BS_a| = |S_aC|$
- 3) $k_t; k_t(S_a, S_aC)$... the Thales' circle
- 4) $k_1; k_1(B, v_b)$
- 5) $B_1; B_1 \in k_1 \cap k_t$
- 6) $\rightarrow CB_1$
- 7) $Y; Y \in \rightarrow CB_1$... a point of your own choice
- 8) $\sphericalangle CYX; \sphericalangle CYX = \alpha$
- 9) $\leftrightarrow p; \leftrightarrow p \parallel XY \wedge B \in \leftrightarrow p$
- 10) $A; A \in \leftrightarrow p \cap \rightarrow CB_1$
- 11) $\triangle ABC$

The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/raxdkacg>

Example 4.18. Construct a triangle ABC if given: b, c, v_a .



The construction step by step


The record of the construction:

- 1) $AA_1; |AA_1| = v_a$
- 2) $\leftrightarrow p; \leftrightarrow p \perp AA_1 \wedge A_1 \in \leftrightarrow p$
- 3) $k_1; k_1(A, b)$
- 4) $C; C \in k_1 \cap \leftrightarrow p$
- 5) $k_2; k_2(A, c)$
- 6) $B; B \in k_2 \cap \leftrightarrow p$
- 7) $\triangle ABC$

The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/ny5an7tf>


Example 4.19. Construct a triangle ABC if given: α, v_b, r_v .



The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/smnfqkqf>


Example 4.20. Construct a triangle ABC if given: c, v_a, v_b .



The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/yzcd6acd>


Example 4.21. Construct a triangle ABC if given: a, v_a, v_b .



The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/rkdepcxf>

Example 4.22. Construct a triangle ABC if given: b, c, t_c .



The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/gnr4vvn>

Example 4.23. Construct a triangle ABC if given: b, γ, v_c .

The construction step by step

The record of the construction:

- 1) $AC; |AC| = b$
- 2) $\sphericalangle ACX; \sphericalangle ACX = \gamma$
- 3) $S_b; S_b \in AC \wedge |AS_b| = |S_b C|$
- 4) $k_t; k_t(S_b, S_b A) \dots$ the Thales' circle
- 5) $k_1; k_1(C, v_c)$
- 6) $C_1; C_1 \in k_t \cap k_1$
- 7) $\rightarrow AC_1$
- 8) $B; B \in \rightarrow AC_1 \cap CX$
- 9) $\triangle ABC$

10 / 10

2 s

The construction step by step:
<https://www.geogebra.org/m/u7e5f3qn#material/rssprtnv>

Example 4.24. Construct a triangle ABC if given: γ, v_a, v_b .

The construction step by step

The record of the construction:

- 1) $\sphericalangle XCY; / \sphericalangle CXY/ = \gamma$
- 2) $\leftrightarrow p; \leftrightarrow p \parallel CY \wedge \leftrightarrow pCY/ = v_a$
the radius of the circle $k_1 = v_a$
- 3) $A; A \in \leftrightarrow p \cap CX$
- 4) $\leftrightarrow q; \leftrightarrow q \parallel CX \wedge \leftrightarrow qCX/ = v_b$
the radius of the circle $k_2 = v_b$
- 5) $B; B \in \leftrightarrow q \cap CY$
- 6) ΔABC

The construction step by step:
<https://www.geogebra.org/m/u7e5f3qn#material/sn3wvaed>

Example 4.25. Construct a triangle ABC if given: a, v_a, b .

The construction step by step

The record of the construction:

- 1) $BC; /BC/ = a$
- 2) $\leftrightarrow p, \leftrightarrow p'; / \leftrightarrow pBC/ = / \leftrightarrow p'BC/ = v_a$
- 3) $k_1; k_1 (C, b)$
- 4) $A; A \in k_1 \cap \leftrightarrow p$
- 5) ΔABC

9 / 9

The construction step by step:
<https://www.geogebra.org/m/u7e5f3qn#material/ntwfvxns>

Example 4.26. Construct a triangle ABC if given: α , c , t_c .

The construction step by step The record of the construction:

- 1) AB ; $|AB| = c$
- 2) $\sphericalangle BAX$; $|\sphericalangle BAX| = \alpha$
- 3) S_c ; $S_c \in AB \wedge |AS_c| = |S_cB|$
- 4) k_1 ; $k_1(S_c, t_c)$
- 5) C ; $C \in k_1 \cap \rightarrow AX$
- 6) $\triangle ABC$

7 / 7 2 s

The construction step by step:
<https://www.geogebra.org/m/u7e5f3qn#material/trhbazkf>

Example 4.27. Construct a triangle ABC if given: γ, v_a, v_c .

The construction step by step

The record of the construction:


- 1) $\sphericalangle XCY; / \sphericalangle XCY / = \gamma$
- 2) $\leftrightarrow p; \leftrightarrow p \parallel \mapsto CX \wedge v (\leftrightarrow p, \mapsto CX) = v_a$
the radius of the circle $k = v_a$
- 3) $A; A \in \leftrightarrow p \cap \mapsto CY$
- 4) $S_b; S_b \in AC \wedge /CS_b/ = /S_bA/$
- 5) $k_t; k_t(S_b, S_bA)$
- 6) $k_1; k_1(C, v_c)$
- 7) $C_1; C_1 \in k_1 \cap k_t$
- 8) $\mapsto AC_1$
- 9) $B; B \in \mapsto AC_1 \cap \mapsto CX$
- 10) $\triangle ABC$

11 / 11 2 s

The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/njnjbvh9>

Example 4.28. Construct a triangle ABC if given: α, β, r_v , kde r_v je poloměr kružnice trojúhelníku vepsané.



The construction step by step:

<https://www.geogebra.org/m/u7e5f3qn#material/w547a5au>

Exercises 4.29. Construct a triangle ABC if given:

- | | | |
|---------------------------|-----------------------------|-------------------------------|
| a) $a + b, \gamma, v_a$ | b) $a - b, \gamma, c$ | c) $a + b + c, \alpha, \beta$ |
| d) $a, b, \alpha - \beta$ | e) $a + b + c, \alpha, v_c$ | |


Exercises 4.30. The line AB is given.

- a) Construct the set of all vertices of the convex angle $\sphericalangle ACB = \textit{gamma}$, whose arms pass through the endpoints of the line segment AB .
- b) Construct $\triangle ABC$ if $|AB| = 6, \gamma = 60^\circ, v_C = 4$.

Exercises 4.31. Construct a triangle ABC if given: t_a, t_b, t_c .

Example 4.32. Prove the sentence about the centroids of the triangle: The centroids of each triangle intersect at one point, called the centroid of the triangle. the center of gravity divides each center of gravity into two lines, the one containing the vertex of the triangle is twice the other.

Proof: The triangle ABC is given, the points A_1, B_1, C_1 are the center of its sides BC, AC and AB , the lines AA_1, BB_1 and CC_1 is his center of gravity. In this triangle, we consider the AA_1 and BB_1 mines that intersect at T . We prove that CC_1 goes through T .



Let's construct a line CT and a point U on it so that the point T is the center of the line CU , ie $CT \cong TU$. In the AUC triangle, the line B_1T is the middle bar and therefore $B_1T \parallel AU$. Since the points B_1, T, B lie on one straight line, it is $BT \parallel AU$. Analogously in the BUC triangle, the line A_1T is the middle rung and therefore $A_1T \parallel BU$, and hence $AT \parallel BU$. Hence the quadrilateral $ATBU$ has two parallel sides parallel, ie it is a parallelogram and its diagonals AB and TU are bisected. Hence, the center of the AB side, C_1 , lies on the line CT . This proves that CC_1 goes through T . Therefore, the centroids of the triangle ABC intersect at one point. This point always belongs to the inside of the given triangle.

The properties of the middle rungs B_1T and A_1T of the triangles AUC and BUC and the properties of the parallelogram $AUBT$ further imply:

$$\text{for triangle } AUC: B_1T = \frac{1}{2}AU, AU \cong BT, \text{ e.i. } B_1T = \frac{1}{2}BT,$$

for triangle BUC : $A_1T = \frac{1}{2}BU$, $BU \cong AT$, e.i. $A_1T = \frac{1}{2}AT$.

This proves that the center of gravity T divides each of the AA_1 , BB_1 lines into two parts, the one containing the vertex of the triangle is twice the other. By repeating the considerations in choosing another pair of lines, we obtain further relationships, which imply the truthfulness of the statement of the second part of the sentence.

Exercises 4.33. Prove that two triangles are identical when they match in two sides and in the center of gravity to one of them.

Instruction: Prove the identity of triangles by using triangles, which are created by dividing a given triangle by the centroid.

Exercises 4.34. Above the sides of the acute triangle ABC are equilateral triangles ABH and ACK . Prove line segments CH and BK .

Instructions: The assertion follows from the equality of triangles ACH and AKB .

Exercises 4.35. The triangle ABC is given. Its medians are drawn parallel to the opposite sides. Prove that the intersections of these lines determine the triangle, which is a union of four triangles identical to the triangle ABC .

Instructions: Use the theorems of triangles identity and properties of pairs of angles between parallel lines.

Exercises 4.36. The largest side of the convex quadrilateral $ABCD$ is AB , the smallest CD . Prove that $\sphericalangle ABC < \sphericalangle ADC$.


Instructions: The BD diagonal divides the $ABCD$ quadrilateral into two triangles. The assumption results in inequalities, the sum of which gives us the assertion.

Exercises 4.37. On the diagonal AC of the square $ABCD$ is given the point E so that $AE \cong AB$. The perpendicular to the AC line through the E point crosses the BC side at the F point. Prove that $BF \cong EF$.

Instructions: Prove that triangle ECF is isosceles and triangle AFE is identical to triangle AFB .

Example 4.38. The illustration shows seven different quadrilaterals. Assign them to their names and then add their properties (some properties may belong to more than one quadrilateral):
square, rectangle, rhombus, (generic) parallelogram, non-convex quadrilateral, deltoid, trapezoid.

- a) The opposite sides are always the same.
- b) At least two internal angles are always right.
- c) The diagonals are halved.
- d) The diagonals are identical.
- e) You can circle it.
- f) You can write a circle.
- g) Just one pair of sides are parallel lines.




Solution:

- square $EFGH$, a), b), c), d), e), f),
- rectangle $OPQR$, a), b), c), d), e),
- rhombus $A_1B_1C_1D_1$, a), c), f),
- (general) parallelogram $STUV$, a), c),
- non-convex quadrilateral $ABCD$,
- deltoid $XZYW$, b), e), f),
- trapezoid $KLMN$, g).

Example 4.39. Construct a parallelogram $ABCD$, given the a side, the DAC angle, $|\sphericalangle DAC| = \alpha$, and the diagonal size $e = |AC|$.

Solution: Select the line AB , $|AB| = a$. The point C lies at a distance of e from the point A , ie on the circle $k(A, e)$. In addition, the BC ray also forms an α angle with the AB side. For D , $CD \parallel AB$ and $AD \parallel BC$ apply.



The construction step by step:

1. AB , $|AB| = a$
2. k , $k(A, e)$
3. X , $X \in \rightarrow AB$
4. $\sphericalangle XBY$, $|\sphericalangle XBY| = \alpha$

5. $C, C \in k \cap \mapsto BY$
6. $D, CD \parallel AB \wedge AD \parallel BC$
7. parallelogram $ABCD$

Conclusion: The problem has one solution in the given half-plane.

Exercises 4.40. Construct a parallelogram $ABCD$, given the size of its diagonals e, f and the size of the height v_a .

Exercises 4.41. Construct a parallelogram $PQRS$, given its diagonal PR , the angle RPQ and the distance of the parallel sides PQ and RS .

Exercises 4.42. Construct a rhombus $ABCD$ if $e = |AC|$ is given and the angle DAB is α .

Exercises 4.43. Construct a rectangle $KLMN$ if given $|KL| = 6$ and the angle KSL is 120° , where S is the intersection of the diagonals.

Exercises 4.44. Construct a trapezoid $ABCD$ if all sides of a, b, c, d are given.

Exercises 4.45. Construct a trapezoid $ABCD$, $AB \parallel CD$ if the diagonal sizes are e, f , the angle size $DAB = \alpha$, and the angle size $AEB = \omega$, where E is intersection of diagonals.

Exercises 4.46. Construct trapezoid $ABCD$ if given: side size AB , side size BC , size of both diagonals AC, BD and angle size $AEB = \omega$, where E is the intersection of diagonals .

Exercises 4.47. A quadrilateral which can be described and inscribed by a circle, ie a quadrilateral which is both chord and tangent *two-centered*. Can you identify at least one non-square two-centered quadrilateral?

Exercises 4.48. Construct a circle k if its tangent t is given with a touch point T and another tangent q .

Exercises 4.49. Construct the k circle that touches the m circle at that point T a

- a) is centered on the given line p ,
- b) goes through M ,
- c) it touches a given line q .

Exercises 4.50. A circle k is given and two different points K, L outside. Construct the rhombus $KLMN$ so that one of its vertices lies on a circle k .


Exercises 4.51. Construct a circle that passes through the A point and directly touches the t line at T .

Exercises 4.52. Construct a circle that is centered on the m circle and touches the two

- parallel lines a, b ,
- non-parallel lines c, d .

Example 4.53. Construct a circle with a radius of $r = 2$ cm that touches the outside of the circle $m(0.3 \text{ cm})$ and goes through that point $M, |SM| = 6$ cm.

Solution: We construct a circle $m, m(0, 3 \text{ cm})$ and a point $M, |SM| = 6$ cm. The center S of the searched circle k is at a distance of 2 cm from the point M , ie on the circle $n(M, 2 \text{ cm})$. Also, the distance of the center S from the touch points, eg A , of the searched circle k with the given circle m is 2 cm. The set of all such points will be on a circle with a radius 2 cm larger than the radius of the given circle m , eg on the circle $l(0, 5 \text{ cm})$. The searched center S lies at the intersection of the circle n and l .



The construction step by step:

1. $m, m(O, 3 \text{ cm}); M, |SM| = 6 \text{ cm}$
2. $n, n(M, 2 \text{ cm})$
3. $l, l(O, 5 \text{ cm})$
4. $S, S \in n \cap l$
5. $k, k(S, 2 \text{ cm})$

Conclusion: The problem has two solutions in the plane.

Exercises 4.54. Construct a circle that touches the two concentric circles k_1, k_2 and goes through the point P , which is the inner point of the annulus specified by the circles k_1, k_2 .

Exercises 4.55. There are two concentric circles $k_1(S, r_1), k_2(S, r_2)$. Investigate the set of centers of all circles that touch k_1, k_2 .

Exercises 4.56. Investigate the set of centers of all circles that

- a) have a given radius of r and go through two different points A, B ;
- b) have a given radius r and touch a given line p ;
- c) touch two given parallels a, b ;
- d) touch two given divergences a, b ;
- e) touch a given line p at a given point A ;
- f) touch a given circle k at the given point A ;
- g) have a given radius of r and have $k(S, r_1)$ external touch.

Model these tasks in GeoGebra.


Exercises 4.57. A circle $k(S, r)$ is given, followed by a point A . Investigate the set of centers of all chords of the k circle that pass through A . Model the task in GeoGebra.

Exercises 4.58. A circle $k(S, r)$ is given, and on it a point N that belongs to the outer region of that circle. Investigate the set of centers of all the chords of the k circle that lie on the cuts passing through N .

Model the task in GeoGebra.

Exercises 4.59. Construct a regular a) octagon, b) dvanáctiúhelník, c) šestnáctiúhelník.

Example 4.60. Construct a grid of curves that was used to create a Gothic window.








Attachments




Pictures to exercise 1.1



4	9	2
3	5	7
8	1	6









Solving puzzles from the example 3.13


A)




B)



C)



D)



References

- [1] Francová, M., Lvovská, L., *Texty k základům elementární geometrie pro studium učitelství 1. stupně základní školy*, skriptum PedF MU, Brno 2014.
- [2] Francová, M., Matoušková, K., Vaňurová, M. *Texty k základům elementární geometrie pro studium učitelství 1. stupně základní školy*, skriptum UJEP, Brno 1985.
- [3] Francová, M., Matoušková, K., Vaňurová, M. *Sbírka úloh z elementární geometrie*, skriptum MU, Brno 1996.
- [4] Lomtatidze, L. *Historický vývoj pojmu křivka*, Scintilla Svazek 3, Brno 2007
- [5] Vopěnka, P. *Rozpravy s geometrií*, Academia, Praha 1989
- [6] Struik, D. J. *Dějiny matematiky*, Praha 1963. (z angl. originálu *A concise History of Mathematics*, G. Bell and Sons Ltd., London 1956, přeložili Nový, L. - Folta, J.)
- [7] Katz, V. J. *A history of mathematics: an introduction*, Addison-Wesley Educational Publishers, Inc., 2. vydání, 1998.
- [8] Servít, F. *Eukleidovy Základy (Elementa)*. Nákladem Jednoty českých matematiků a fyziků, Praha, 1907.
- [9] Bečvářová M., *Eukleidovy Základy, jejich vydání a překlady* Dějiny matematiky, svazek 20. Prometheus, Praha, 2001.
- [10] Citáty na téma geometrie, <https://citaty.net/temata/geometrie/>

Contents

1	Historical development of geometry	6
2	Basic geometric formations and their properties	9
3	Convex and non-convex set, convex and non-convex angle	14
4	Circle, round, triangle, quadrilateral, regular polygon	20
	Attachments	44
	References	48