Research centre
for toxic compounds
in the environment

Ecotoxicology – Part 3

Current issues in Research vs Regulation

Ludek Blaha + ecotox colleagues

cetocoen

EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

2007-13
OP Research and
Development for Innovation

UNIVERSITAS MASARYKIANA BRUNENSIS

When

Where

the assessment of toxicity is needed ?

What

to assess for toxicity ?

Research centre
for toxic compounds
in the environment

cetocoen

EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

2007-13
OP Research and
Development for Innovation

When & where the toxicity assessment is needed?

View of the researcher

Anytime!

… depending on researcher's budget

View of the regulator

As the law says!

… what are the law(s)? →



Chemical laws („bulk“)

- Industrial chemicals
- Cosmetics
- PPP (pesticides)
- Biocides
- Human pharmaceuticals
- Veterinary pharmaceuticals

MIXTURES!

nano

nano

nano

nano

§§

REACH (ECHA)

PPP (EFSA)

MPs (EMA)

Two §§ approaches:

- Prospective (chemicals…)
- Retrospective (mixtures …)



§§

WFD – surface w.
GWD – ground w.

Air quality

Food and feed

Soil & Sediments

Wastes

Research centre for toxic compounds in the environment

INVESTING IN YOUR FUTURE

OP Research and Development for Innovation

What to assess for toxicity?



	Current research topics	As required by law
Individual chemicals		
Mixtures		
Contaminated samples		

Researc
for toxic
in the er

What to assess for toxicity?



	Current research topics	As required by law
Individual chemicals	Engineered **nanomaterials**/particles **Ecological effects** (e.g. of pharmaceuticals) **Endocrine** disruption & **chronic** diseases	Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics
Mixtures		
Contaminated samples		

Researc
for toxic
in the er

What to assess for toxicity?



	Current research topics	As required by law
Individual chemicals	Engineered **nanomaterials**/particles **Ecological effects** (e.g. of pharmaceuticals) **Endocrine** disruption & **chronic** diseases	Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics
Mixtures	**Multistressors** +T°C, salinity, pathogens, irradiation, food **Exposome**	LOADING
Contaminated samples		

Researc
for toxic
in the er

What to assess for toxicity?



	Current research topics	As required by law
Individual chemicals	Engineered **nanomaterials**/particles **Ecological effects** (e.g. of pharmaceuticals) **Endocrine** disruption & **chronic** diseases	Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics
Mixtures	**Multistressors** +T°C, salinity, pathogens, irradiation, food **Exposome**	LOADING
Contaminated samples	Can analyzed chemicals explain observed effects ?	**Chemical analyses & limits** **Effect testing rare**: Remediation, dredged sediments (CZ), effluents (DE) LOADING

Nanoparticles - examples













EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

OP Research and Development for Innovation

Toxicity of nanoparticles …



(Mostly unknown) Parameters may Affect ecotoxicity

Composition (chemical)
Surface (size, area)
Charge
Reactivity
Interactions with ions, other chemicals…

➔ Effects on environmental Fate and toxicity

EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

OP Research and Development for Innovation

Ecotoxicity of nanoparticles – RECETOX example

Comparison of toxicity - 4 „appeared to be the same“ particles
(one producer – 4 different lots)
(zerovalent iron – ZVI – Fe^0)



?? Why is H16 so toxic ??
… despite of detailed investigation never revealed

PHARMACEUTICALS



R&D and Manufacturing

Storage Transport

Distribution

Storage Transport

Consumption

Storage Transport

Waste management

Manufacturing waste

Possible releases to the environment

Example 1 - DICLOFENAC

Unexpected effects at NON-TARGET species

- **nephrotoxicity** at vultures
- Relevant also in EU (ESP, EL,CY)





Example 2 – AVERMEKTIN-like antiparasitics

Moxidectin – used e.g. in home „spot on" products

Ivermectin – antiparasitics in large herds

- Used **2-times per season** per sheep/cow
- **Kills 100% parasites** in sheep
- Released in dung - **kills 80-90% larvae of dung flies**
- High concentrations in dung (released 2 days post application)
- **Persistent in the soil** (half-life 30 days)
- Can be washed into adjacent streams (highly toxic to water insects)



POLICYFORUM

Science
AAAS

CONSERVATION

Economic Importance of Bats in Agriculture

Insectivorous bat populations, adversely impacted by white-nose syndrome and wind turbines, may be worth billions of dollars to North American agriculture.

Justin G. Boyles,[1]* Paul M. Cryan,[2] Gary F. McCracken,[3] Thomas H. Kunz[4]





Boyles et al. (2011) Science 332 (60251) 41-42

biology letters
Animal behaviour
Biol. Lett.
doi:10.1098/rsbl.2012.0685
Published online

Maternal predator-exposure has lifelong consequences for offspring learning in threespined sticklebacks

Daniel P. Roche, Katie E. McGhee* and Alison M. Bell

School of Integrative Biology, University of Illinois, Urbana, IL 61801, USA
**Author for correspondence (kemcghee@illinois.edu).*

Stress → multigeneration effects



Epigenetics → DNA methylations



Table 1. Behaviours (mean ± s.e.) of the offspring from the maternal treatments.

	offspring of predator-exposed mothers (s)	offspring of unexposed mothers (s)
initial exploratory behaviour (day 1: 09.00):		
latency to first begin moving	49 ± 30	56 ± 20
latency to enter either chamber for the first time	330 ± 70	326 ± 78
learning the colour association:		
day 1 (09.00): latency to find food reward	426 ± 65	427 ± 61
day 3 (09.00): latency to find food reward	533 ± 48	304 ± 74
day 5 (09.00): latency to find food reward	337 ± 61	158 ± 68

2x difference

Neuroscience and Biobehavioral Reviews 32 (2008) 1073–1086

Contents lists available at ScienceDirect

Neuroscience and Biobehavioral Reviews

journal homepage: www.elsevier.com/locate/neubiorev

Review

The long-term behavioural consequences of prenatal stress

Marta Weinstock*

Department of Pharmacology, Hebrew University, Medical Centre, Ein Kerem, Jerusalem 91120, Israel



Fig. 2. Routes by which maternal stress hormones may induce changes in the foetal brain in the programming of offspring behaviour. The developing foetal brain is sensitive to the actions of excess amounts of glucocorticoids and other hormones. These may alter the structure and function of the limbic system and HPA axis resulting permanent changes in behaviour and neuroendocrine regulation in the offspring. ↑ = increase; ↓ = decrease.

International ring test (2012-13)

Testing comparability of existing and innovative bioassays for water quality assessment

Main questions:

Are current limits (for individual compounds) safe?

Relevance of **“Something from Nothing”** phenomenon ?

3 samples

→ 12 European laboratories – different bioassays

→ ČR – RECETOX: 11 bioassays



Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? *Toxicol Sci* **141(1): 218-233**

EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

OP Research and Development for Innovation

International ring test (2012-13)

Testing comparability of existing and innovative bioassays for water quality assessment

EU WFD priority substances

Different concentrations

EQS = limit *(Environmental Quality Standard)*

	RM 1[a]	RM 2[a]	RM 3[a]
Priority substances mg/L	around or >EQS	< EQS	< EQS
Atrazine	6	0.6	0.6
BaP	0.0017	0.00017	0.00017
Cadmium[b]	0.8	0.08	0.08
Chlorfenvinphos	1	0.1	0.1
Chlorpyrifos	0.3	0.03	0.03
DEHP (Bis(2-ethylhexyl) phthalate)	13	1.3	1.3
Diclofenac	1	0.1	0.1
diuron	2	0.2	0.2
17beta-estradiol	0.004	0.0004	0.0004
fluoranthene	0.063	0.0063	0.0063
Isoproturon	3	0.3	0.3
Ni[b]	40	4	4
4-Nonylphenol	3	0.3	0.3
Simazine	10	1	1
Carbamazepine	-	-	0.5
Sulfamethoxazole	-	-	0.6
Triclosan (Irgasan)	-	-	0.02
DEET	-	-	41
Bisphenol A	-	-	1.5

International ring test (2012-13)

Testing comparability of existing and innovative bioassays for water quality assessment

Example: Effects of mixtures on *D. rerio* fish embryos



Control



Effects of RM 3 (i.e. safe) mixtures

Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? *Toxicol Sci* **141(1): 218-233**

EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

OP Research and Development for Innovation

International ring test (2012-13)

Testing comparability of existing and innovative bioassays for water quality assessment

Example: Effects of mixtures on *X. laevis* frog embryos



Controls

Effects of RM 3 (i.e. safe) mixtures



Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? *Toxicol Sci* **141(1): 218-233**

Research centre for toxic compounds in the environment

Biotest	A	B	C
Microtox	26 and 36% stimulation of luminescence in 15 and 30 mins of exposure, respectively	18 and 35% stimulation of luminescence in 15 and 30 mins of exposure, respectively	22 and 39% stimulation of luminescence in 15 and 30 mins of exposure, respectively
Algae growth inhibition test 96-h exposure	31% inhibition of growth compared to solvent control	20% inhibition of growth compared to solvent control	16% inhibition of growth compared to solvent control
Acute immobilization test with *D. magna*	90% immobilization after 48 hours of exposure; 25% immobilization occurred in 50% concentration - not statistically significant	no effect observed	no effect observed
Reproduction test with *D. magna* (21-d exposure)	100% mortality after 3 days of the test, no reproduction could be evaluated	31 +/- 37 % inhibition of reproduction, not statistically significant	23 +/- 24 % inhibition of reproduction, not statistically significant
FETAX (96-h exposure)	62 +/- 10 % of malformed embryos; no effect on embryo length observed	43 +/- 12 % of malformed embryos; no effect on embryo length observed	34 +/- 14 % of malformed embryos; no effect on embryo length observed
FET (120-h exposure)	effects observed in number of defected embryos - absence of gas bladder, (head) deformities and underdeveloped embryos were observed the most often.	no significant effects observed	effects observed in number of defected embryos, number of underdeveloped embryos and length
In vitro - cytotoxicity	no effect observed compared to solvent control	no effect observed compared to solvent control	no effect observed compared to solvent control
In vitro - estrogenicity	effect under LOQ	effect under LOQ	effect under LOQ
In vitro - dioxin-like toxicity	effect under LOQ	effect under LOQ	effect under LOQ
In vitro - androgenicity	effect under LOQ	effect under LOQ	effect under LOQ
In vitro - antiandrogenicity	effect under LOQ	effect under LOQ	effect under LOQ

Contaminated samples? Case study **“air“**

Active sampling
particles *vs* gaseous phase

- **Reference locality –** agriculture (Košetice observatory)
- **Region A** – industrial (historically OCPs production)
- **Region B** – combined: industry, agriculture, traffic



Novák et al. (2009) Environment International

Chemical analyses







Dioxin-like effects

Labs
on Wed + Thu



- Difference B>A
- Difference B vs A – particles vs gas

Antiandrogenic effects



- Quantitative – comparable
- Clear differences in patterns ... no effects on particles in „B“ (?)

Summary on When, Where, What

- **Regulatory world**
  - Assessment of „chemicals“!

- Contaminated samples
  - effects rarely tested
  - **Great value of bioassays** in assessment of contaminated samples
    - Effects observed (!)
    - **How to set the „limits“?**

- **Research issues and questions**
  - Nanomaterials, Pharmaceuticals, EDCs
  - Mixtures!
  - Exposome

ECHA
EUROPEAN CHEMICALS AGENCY
REACH

Contents lists available at ScienceDirect

Environment International

journal homepage: www.elsevier.com/locate/envint

Review

What level of estrogenic activity determined by *in vitro* assays in municipal waste waters can be considered as safe?

Barbora Jarošová [a], Luděk Bláha [a], John P. Giesy [b], Klára Hilscherová [a,*]

[a] Masaryk University, Faculty of Science, RECETOX, Kamenice 5, CZ-62500 Brno, Czech Republic
[b] Department of Biomedical Veterinary Sciences and Toxicology Centre, University of Saskatchewan, Saskatoon, Saskatchewan, Canada