



Central European Institute of Technology
BRNO | CZECH REPUBLIC

*C9940 3-Dimensional Transmission Electron Microscopy
S1007 Doing structural biology with the electron microscope*

Tanvir (Tapu) Shaikh



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE



February 20, 2017



Syllabus


Week	Date	Instructor	Topic
1	20.2	T. Shaikh	Introduction/History/Optics
2	27.2	J. Novacek	Instrumentation/Tour (?)
3	6.3	J. Novacek	Specimen preparation
4	13.3	T. Shaikh	Image analysis I
5	20.3	T. Shaikh	Image analysis II
6	27.3	J. Novacek	Tomography I
7	3.4	J. Novacek	Tomography I
8	10.4	T. Shaikh	Image analysis III
	17.4		(Easter)
9	24.4	T. Shaikh	Image analysis IV
	1.5		(May Day)
	8.5		(Liberation Day)

History of electron microscopy

Munich: Ernst Ruska & Otto Scherzer



Ernst (and Helmut) Ruska: timeline



Milestones:


- ◆ 1931: Invention of the electrostatic lens
 - magnification: 400X
- ◆ 1933: First electron microscope
 - mag: 7000X (vs. LM: 2000X)
- ◆ 1933: Completed Ph.D. (!)
- ◆ 1939: First viable commercial EM
 - mag: 100,000X
- ◆ 1939: His brother Helmut Ruska images first virus (TMV)
- ◆ 1952: Helmut moves from Siemens to Albany
- ◆ 1986: Nobel Prize in Physics

Ernst Ruska




<http://www.biografiasyvidas.com>

Sketch from 1929



<http://ernst.ruska.de>

Replica of first electron microscope



<http://www.bluesci.org>

First (**viable**)
commercial microscope (Siemens)



<http://ernst.ruska.de>

Helmut Ruska (standing)
next to Siemens-20
at Wadsworth Center in Albany



<http://www.wadsworth.org>

Clarification

First Siemens microscope, 1939



<http://ernst.ruska.de>


First **commercial** EM (1937)
was Metropolitan-Vickers EM1
(EM2 shown)



<http://emu.msim.org.uk>


The first commercial electron microscope was actually by the British company Metropolitan-Vickers in 1937. However, the magnification was worse than for the light microscope, so the Siemens is considered “first.”

Metropolitan Vickers
eventually became
AEI, which built the 1.2
million volt EM-7.



<http://www.wadsworth.org>

Otto Scherzer: timeline



Milestones:


- ◆ 1934: First comprehensive book on electron optics
- ◆ 1936: Spherical aberration → resolution 50-100X the wavelength
- ◆ 1947-1951: Devised correction schemes for aberration correction
- ◆ 1949:
 - “Can atoms be visible in the electron microscope?”
 - “Scherzer focus”
- ◆ Scherzer → Harald Rose (Wadsworth, Darmstadt) → Max Haider

Otto Scherzer



<http://www.microscopy.org>

First book on electron optics



<http://www.microscopy.org>

Scherzer (1949) Physikalische Blätter & Scherzer (1949) Journal of Applied Physics

"Can atoms be visible in the electron microscope?"

PHYSIKALISCHE BLÄTTER

1949 Heft 10.11 Seite 460 — 463

Prof. O. Scherzer

Können Atome im Elektronen-Mikroskop sichtbar werden?

... des Auflösungsvermögens für möglich. Es ist also anzunehmen, daß die weitere Entwicklung des Elektronen-Mikroskops eines Tages nicht nur die schweren Jod-Atome des Moleküls, das wir unseren Betrachtungen zu Grunde gelegt haben, sichtbar machen wird, sondern auch die leichten Kohlenstoffatome und damit die Struktur von Molekülen, die weniger übersichtlich gebaut sind.

<http://www.microscopy.org>

"Scherzer focus"





FIG. 4. The function $\sin(3s\Theta^2 - s^2\Theta^4)$, describing the phase shift in case of optimum contrast.


VOLUME 20, JANUARY, 1949

<http://www.microscopy.org>

Toronto group: E.F. Burton, James Hillier, etc.




Toronto group: timeline




Milestones:

- ◆ 1937: Grad students James Hillier, Albert Prebus designed microscope over Christmas break
- ◆ 1938: First images
- ◆ 1939: E.F. Burton et al.
 - introduced airlock system
 - reduced specimen prep to 300nm thickness
 - resolution: 60Å, limited by specimen and not optics
 - maximum mag: 180,000X

Radio Corporation of America (RCA)



James Hillier: timeline



Milestones:

- ◆ 1940 February
 - Hillier started at RCA
 - enlisted by Vladimir Dworykin (cathode ray tube)
- ◆ 1940 Jul 4: Commercial EM, Model B (EMB)
- ◆ 1941: 300kV, for dealing with thick specimens
- ◆ 1947: first stigmator
 - stigmators were iron screws tapped into the pole piece
 - resolution → 1nm
- ◆ 1973 (as VP of RCA): first videodisc


James Hillier

ed, with Albert Prebus standing




<http://www.museevirtuel.ca>

At RCA Model B, 1940




<http://www.rfcafe.com>

Electron microscopy in the Czech Republic



Electron microscopy in the Czech Republic




Milestones:

- ◆ 1950: Ales Blaha: “Tripod” at the Institute of Theoretical and Experimentation Electrotechnology of the Technical University
- ◆ 1951: Tesla BS241
 - first Czech commercial microscope
 - 50kV
 - resolution: 2nm
- ◆ 1968: First ultrahigh vacuum system (Institute of Scientific Instruments)
- ◆ 1990: Delong Instruments founded
- ◆ 1991: TESCAN founded by engineers from Tesla (TEsla SCANing)
- ◆ 1997: FEI builds factory in Brno

Electron microscopy in the Czech Republic

“Tripod,” 1950



<http://www.isibrno.cz>

First high-vacuum system, 1961



<http://www.isibrno.cz>

Electron microscopy in the Czech Republic

TESLA BS 242

Prozařovací elektronový mikroskop
Oceněný zlatou medailí na světové výstavě
EXPO 1958 v Bruselu

Transmission Electron Microscope
Awarded a Gold Medal at the
EXPO 1958 in Brussels




Tesla Factory in Brno




Technical Museum in Brno

Medical Research Council (MRC), Cambridge







Aaron Klug: timeline




Milestones:

- ◆ 1968: DeRosier & Klug – first 3D EM reconstruction (phage T4)
- ◆ 1982: Nobel Prize in Chemistry



Other notable events



Milestones:


- ◆ 1934: Ladislaus Laszlo “Bill” Marton takes the first image of biological specimen: sections of a plant leaf
- ◆ 1937: Manfred von Ardenne (CRT) develops SEM
- ◆ 1948: Dennis Gabor develops electron holography (Nobel Prize in Physics, 1971)
- ◆ 1974: Ken Taylor & Bob Glaeser – electron crystallography of frozen hydrated catalase
- ◆ 1974: Walter Hoppe – 3D reconstruction of fatty acid synthase using tomography
- ◆ 1975: Richard Henderson – subnanometer electron crystallography
- ◆ 1982: Jacques Dubochet – modern cryo techniques
- ◆ 1987: Joachim Frank – “single particle” reconstruction of 50S ribosome
- ◆ 2007: Direct Electron develops first commercial direct electron detector

*“Oh no, not an electron microscope!
We have enough trouble trying to interpret
the images we get with a light microscope!”*
– Jules Bordet, 1934 (Nobel Prize, 1921)

The Beginnings of Electron Microscopy (1985), ed. Peter W. Hawkes

The basics

Size ranges for structural methods



Comparison of practical requirements

	CryoEM	X-ray	NMR
Sample volume	5 μ L ¹	10 μ L	400 μ L
Number of samples	5-10	1 crystal	1 several (<20 kDa) ² (20 - 40 kDa)
Concentration	50 nM	50 μ M ³	0.5-1.0 mM
Total amount of sample	0.25 pmol	500 pmol ⁴	0.2-0.4 μ mol

¹Droplet on grid, prior to blotting

²For stable proteins of this size, a single U-13C, 15N sample is likely sufficient


³Collecting 10 mg/ml of a 100 kDa protein

⁴Final amount required for the crystal. However, a multiple of this amount is required to try out different crystallization conditions


Adapted from Raj Agrawal

Time for data collection 1-7 days 2h (synchrotron) 1-2 weeks (<20 kDa)
48h (in-house) 6-8 weeks (20 - 40 kDa)

Quick review of light optics




Right-hand rule



www.education.com



Right-hand rule

**Path (red) of
electron deflected
by magnetic field
(pointing out of
screen)**



<http://www.polywellnuclearfusion.com>

Electron in a magnetic lens




nau.edu

<http://www.ammrf.org.au>


*One problem:
Magnetic lenses are terrible*

Spherical aberration

Lens with
Spherical Aberration



Aspherical Lens



photographylife.com

In electron optics, there is no lens setting* equivalent to an aspherical lens. Spherical aberration will thus be a problem (but also a benefit).

Syllabus

Week	Date	Instructor	Topic
1	20.2	T. Shaikh	Introduction/History/Optics
2	27.2	J. Novacek & T. Shaikh	Instrumentation/Tour (?)
3	6.3	J. Novacek	Specimen preparation
4	13.3	T. Shaikh	Image analysis I
5	20.3	T. Shaikh	Image analysis II
6	27.3	J. Novacek	Tomography I
7	3.4	J. Novacek	Tomography I
8	10.4	T. Shaikh	Image analysis III
	17.4		(Easter)
9	24.4	T. Shaikh	Image analysis IV
	1.5		(May Day)
	8.5		(Liberation Day)

Thank you for your attention



Central European Institute of Technology
Masaryk University
Kamenice 753/5
625 00 Brno, Czech Republic

www.ceitec.muni.cz | info@ceitec.muni.cz



EUROPEAN UNION
EUROPEAN REGIONAL DEVELOPMENT FUND
INVESTING IN YOUR FUTURE

