

Research centre for toxic compounds in the environment

Ecotoxicology – Part 3

Current issues in Research vs Regulation

Ludek Blaha + ecotox colleagues





EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FUND INVESTING IN YOUR FUTURE






When Where

the assessment of toxicity is needed

What





Research centre for toxic compounds in the environment





EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FUNI INVESTING IN YOUR FUTURE OP Research and evelopment for Innovation

When & where the toxicity assessment is needed?

View of the researcher



Anytime!

... depending on researcher's budget

View of the regulator



As the law says!

... what are the

law(s)?




 \rightarrow










What to assess for toxicity?






| | Current research topics | As required by law |
|--|-------------------------|--------------------|
| Individual chemicals (prospective) | | |
| Mixtures (prospective) | | |
| Contaminated samples (retrospective) | | |
| Researce for toxic in the e | | |


What to assess for toxicity?

Y




| | | and and a second s |
|---|--|--|
| | Current research topics | As required by law |
| Individual chemicals (prospective) | Engineered nanomaterials /particles Ecological effects (e.g. of pharmaceuticals) Endocrine disruption & chronic diseases | Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics |
| Mixtures (prospective) | | |
| Contaminated samples (retrospective | | |
| Rese for t in the | oxic | |

Nanoparticles - examples



Toxicity of nanoparticles ...



(Mostly unknown) Parameters may Affect ecotoxicity

Composition (chemical) Surface (size, area) Charge Reactivity Interactions with ions, other chemicals...


Effects on environmental Fate and toxicity





Ecotoxicity of nanoparticles – RECETOX example

Comparison of toxicity - 4 "appeared to be the same" particles (one producer – 4 different lots) (zerovalent iron – $ZVI – Fe^{0}$)



?? Why is H16 so toxic ?? ... despite of detailed investigation never revealed











PHARMACEUTICALS




Example 1 - DICLOFENAC

Unexpected effects at NON-TARGET species

- nephrotoxicity at vultures
- Relevant also in EU (ESP, EL,CY)









Research centre for toxic compounds in the environment





EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FUND INVESTING IN YOUR FUTURE OP Research and


Example 2 – AVERMEKTIN-like antiparasitics

Moxidectin – used e.g. in home "spot on" products



Ivermectin – antiparasitics in large herds

- Used 2-times per season per sheep/cow
- Kills 100% parasites in sheep
- Released in dung kills 80-90% larvae of dung flies
- High concentrations in dung (released 2 days post application)
- Persistent in the soil (half-life 30 days)
- Can be washed into adjacent streams (highly toxic to water insects)









Research centre for toxic compounds in the environment





EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FUN INVESTING IN YOUR FUTURE




CONSERVATION

Economic Importance of Bats in Agriculture

Justin G. Boyles,^{1*} Paul M. Cryan,² Gary F. McCracken,³ Thomas H. Kunz⁴

POLICYFORUM Science MAAAS

Insectivorous bat populations, adversely impacted by white-nose syndrome and wind turbines, may be worth billions of dollars to North American agriculture.



Bovles et



esearch centre for toxic compound n the environment

biology letters Animal behaviour

Biol. Lett. doi:10.1098/rsbl.2012.0685 Published online

Stress

→ multigeneration effects

Maternal predatorexposure has lifelong consequences for offspring learning in threespined sticklebacks

Daniel P. Roche, Katie E. McGhee* and Alison M. Bell

School of Integrative Biology, University of Illinois, Urbana, IL 61801, USA *Author for correspondence (kemcghee@illinois.edu).




Table 1. Behaviours (mean \pm s.e.) of the offspring from the maternal treatments.

| | offspring of predator-exposed mothers (s) | offspring of unexposed mothers (s) |
|--|--|---------------------------------------|
| initial exploratory behaviour (day 1: 09.00): | | |
| latency to first begin moving | 49 ± 30 | 56 ± 20 |
| latency to enter either chamber for the first time | 330 ± 70 | 326 ± 78 |
| learning the colour association: | | |
| day 1 (09.00): latency to find food reward | 426 + 65 | 427 ± 61 |
| day 3 (09.00): latency to find food reward | 533 ± 48 2x difference | 304 ± 74 |
| day 5 (09.00): latency to find food reward | $_{337\pm 61}$ 2X unterence | 158 ± 68 |



Contents lists available at ScienceDirect

Neuroscience and Biobehavioral Reviews



journal homepage: www.elsevier.com/locate/neubiorev

Review






Fig. 2. Routes by which maternal stress hormones may induce changes in the foetal brain in the programming of offspring behaviour. The developing foetal brain is sensitive to the actions of excess amounts of glucocorticoids and other hormones. These may alter the structure and function of the limbic system and HPA axis resulting permanent changes in behaviour and neuroendocrine regulation in the offspring. \uparrow = increase; \downarrow = decrease.



Research centre for toxic compounds in the environment

International ring test (2012-13)


Testing comparability of existing and innovative bioassays for water quality assessment

Main questions:

Are current limits (for individual compounds) safe? Relevance of **"Something from Nothing"** phenomenon ?

3 samples

- → 12 European laboratories different bioassays
- → ČR RECETOX: 11 bioassays



Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? *Toxicol Sci* 141(1): 218-233

EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FU INVESTING IN YOUR FUTURE




International ring test (2012-13)

Testing comparability of existing and innovative bioassays for water quality assessment


| EU WFD | | RM 1ª | RM 2 ^ª | RM 3 ª |
|---------------------------|----------------------------|-----------------------|-------------------|---------|
| priority substances | <i>Priority substances</i> | around <u>or</u> >EQS | < EQS | < EQS |
| Different | Atrazine | 6 | 0.6 | 0.6 |
| concentrations | BaP | 0.0017 | 0.00017 | 0.00017 |
| concentratione | Cadmium ^b | 0.8 | 0.08 | 0.08 |
| EQS | Chlorfenvinphos | 1 | 0.1 | 0.1 |
| = limit | Chlorpyrifos | 0.3 | 0.03 | 0.03 |
| (Environmental Quality | DEHP (Bis(2-ethylhexyl) | | | |
| Standard) | phthalate) | 13 | 1.3 | 1.3 |
| | Diclofenac | 1 | 0.1 | 0.1 |
| | diuron | 2 | 0.2 | 0.2 |
| | 17beta-estradiol | 0.004 | 0.0004 | 0.0004 |
| | fluoranthene | 0.063 | 0.0063 | 0.0063 |
| | Isoproturon | 3 | 0.3 | 0.3 |
| | Ni ^b | 40 | 4 | 4 |
| | 4-Nonylphenol | 3 | 0.3 | 0.3 |
| | Simazine | 10 | 1 | 1 |
| | Carbamazepine | - | - | 0.5 |
| | Sulfamethoxazole | - | - | 0.6 |
| | Triclosan (Irgasan) | - | - | 0.02 |
| Resea | DEET | - | - | 41 |
| for to in the | Bisphenol A | - | - | 1.5 |

International ring test (2012-13) Testing comparability of existing and innovative bioassays for water quality assessment

Example: Effects of mixtures on D. rerio fish embryos



Control



Effects of RM 3 (i.e. safe) mixtures

Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? *Toxicol Sci* 141 (1), 218-233



EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FUND

OP Research and Development for Innovation

International ring test (2012-13) Testing comparability of existing and innovative bioassays for water quality assessment

Example: Effects of mixtures on X. laevis frog embryos


Controls

Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? *Toxicol Sci* **141(1): 218-233**



Research centre for toxic compounds in the environment


Effects of RM 3 (i.e. safe) mixtures



| Biotest | Α | В | c |
|-----------------------------------|-------------------------------------|-----------------------------------|-----------------------------------|
| Microtox | 26 and 36% stimulation of | 18 and 35% stimulation of | 22 and 39% stimulation of |
| | luminescence in 15 and 30 mins of | luminescence in 15 and 30 mins of | luminescence in 15 and 30 mins of |
| | exposure, respectively | exposure, respectively | exposure, respectively |
| | | | |
| Algae growth inhibition test 96-h | 31% inhibition of growth compared | 20% inhibition of growth compared | 16% inhibition of growth compared |
| exposure | to solvent control | to solvent control | to solvent control |
| Acute immobilization test with | 90% immobilization after 48 hours | no effect observed | no effect observed |
| D. magna | of exposure; 25% immobilization | | |
| | occurred in 50% concentration - not | | |
| | statistically significant | | |
| Reproduction test with D. | 100% mortality after 3 days of the | 31 +/- 37 % inhibition of | 23 +/- 24 % inhibition of |
| magna (21-d exposure) | test, no reproduction could be | reproduction, not statistically | reproduction, not statistically |
| | evaluated | significant | significant |
| | | | |
| FETAX (96-h exposure) | 62 +/- 10 % of malformed embryos; | 43 +/- 12 % of malformed embryos; | 34 +/- 14 % of malformed embryos; |
| π | no effect on embryo length | no effect on embryo length | no effect on embryo length |
| • | observed | observed | observed |
| FET (120-h exposure) | effects observed in number of | no significant effects observed | effects observed in number of |
| | defected embryos - absence of gas | | defected embryos, number of |
| | bladder, (head) deformities and | | underdeveloped embryos and |
| | underdeveloped embryos were | | length |
| | observed the most often. | | * |
| In vitro - cytotoxicity | no effect observed compared to | no effect observed compared to | no effect observed compared to |
| | solvent control | solvent control | solvent control |
| In vitro - estrogenicity | effect under LOQ | effect under LOQ | effect under LOQ |
| In vitro - dioxin-like toxicity | effect under LOQ | effect under LOQ | effect under LOQ |
| In vitro - androgenicity | effect under LOQ | effect under LOQ | effect under LOQ |
| In vitro - antiandrogenicity | effect under LOQ | effect under LOQ | effect under LOQ |

What to assess for toxicity?






| | Current research topics | As required by law | |
|--|---|--|--|
| Individual chemicals (prospective) | Engineered nanomaterials/particles Ecological effects (e.g. of pharmaceuticals) Endocrine disruption & chronic diseases | Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics | |
| Mixtures (prospective) | Multistressors +T°C, salinity, pathogens, irradiation, food Exposome | LOADING | |
| Contaminated samples (retrospective) | | | |
| Researce for toxic in the el | | | |

What to assess for toxicity?



or toxic compounds in the environment



| | | Current research topics | As required by law | |
|--|--|--|--|--|
| | Individual chemicals (prospective) | Engineered nanomaterials /particles Ecological effects (e.g. of pharmaceuticals) Endocrine disruption & chronic diseases | Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics | |
| | Mixtures (prospective) | Multistressors +T°C, salinity, pathogens, irradiation, food Exposome | LOADING | |
| | Contaminated samples (retrospective) | Can analyzed chemicals explain observed effects ? | Chemical analyses & limits (see lectures: RISK ASSESSMENT part) | |
| | | | Effect testing rare: Remediation, dredged sediments (CZ), effluents | |
| | | | | |
| | | | Commission | |
| | for toxi | c compounds | TECHNICAL REPORT ON AQUATIC EFFECT-BASED MONITORING TOOLS | |

Contaminated samples? Case study "air"

Active sampling particles vs gaseous phase

- Reference locality agriculture (Košetice observatory)
- Region A industrial (historically OCPs production)
- **Region B** combined: industry, agriculture, traffic


Novák et al. (2009) Environment International






EUROPEAN REGIONAL DEVELOPMENT FUN INVESTING IN YOUR FUTURE

OP Research and evelopment for Innovation



Dioxin-like effects

dioxin-like toxicity



Difference B>A
 Difference B vs A – particles vs gas



Research centre for toxic compounds in the environment






EUROPEAN UNION EUROPEAN REGIONAL DEVELOPMENT FUNI INVESTING IN YOUR FUTURE



Antiandrogenic effects

antiandrogenicity



Clear differences in patterns ... no effects on particles in "B

OP Research and

Development for Innovation

VESTING IN YOUR FUTUR

CEIOCOEN

Quantitative – comparable



Summary on When, Where, What

Regulatory world

– Assessment of "chemicals"!

Contaminated samples

- effects rarely tested
 - Great value of bioassays in assessment of contaminated samples
 - Effects observed (!)
 - How to set the "limits"?

Research issues and questions

- Nanomaterials, Pharmaceuticals, EDCs
- Mixtures!
- Exposome



Research centre for toxic compounds in the environment





Contents lists available at ScienceDirect

Environment International

journal homepage: www.elsevier.com/locate/envint

Review

What level of estrogenic activity determined by *in vitro* assays in municipal waste waters can be considered as safe?

Barbora Jarošová ^a, Luděk Bláha ^a, John P. Giesy ^b, Klára Hilscherová ^{a,*}

^a Masaryk University, Faculty of Science, RECETOX, Kamenice 5, CZ-62500 Brno, Czech Republic
^b Department of Biomedical Veterinary Sciences and Toxicology Centre, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

