





Bonding

Potential Energy of a Bond



Stretching vibration

Bond Length and Temperature



Experimental Mo≡Mo bond distance:
at room temp. 2.23 Å, at LN2 t. 2.18 Å

Harmonic oscillator in quantum mechanics
Quantized vibrational energies
 $E^n = (v + \frac{1}{2})\hbar\nu$

Morse - Potential energy curve for a diatomic molecule

D_0 = bond dissociation energy


The ground-state energy is not 0

Bond	wavenumber cm ⁻¹
C–H	3000
C–D	2200

The C–H bond is a stronger spring
The C–D bond has lower zero-point energy
Force constant does not change with isotopic substitution
The C–H bond is longer/shorter than the C–D bond?

$$\nu = \frac{1}{2\pi} \sqrt{\frac{k}{\mu}}$$

Bond Length and Temperature



$$E^n = (v + \frac{1}{2})\hbar\nu$$


Potential energy curve is not parabolic (harmonic) but unsymmetrical

99.9 % of C-H bonds are in the ground vibrational state ($v = 0$) at room temperature

At high temperature, more high vibronic states are populated

Middle points move towards longer r

Bond Length and Isotopes



A Morse potential curve

The zero point vibrational energy (ZPE) is the lowest possible energy of a system, the ground state energy.

E_D^0 and E_H^0 (for $v = 0$)


the ZPE of R-H and R-D (R is much heavier than H or D) depend upon the reduced mass of the molecule.

The heavier the molecule or atom, the lower the frequency of vibration and the smaller ZPE.

Lighter molecules or atoms have a greater frequency of vibration and a higher ZPE.

Deuterium is heavier than hydrogen and therefore has the lower zero point energy.

Bond Length and Isotopes




Due to the anharmonicity of the C-H/D vibrational potential energy function and the lower ZPE of a C-D bond, the average C-D bond length is typically $\sim 0.005 \text{ \AA}$ shorter than an equivalent C-H bond.

Secondary Isotope Effect - Steric effect of a CD_3 group smaller than CH_3 .

Kinetic Isotope Effects

The kinetic isotope effect (KIE) = isotopically substituted molecules exhibit different reaction rates, the change in rate of a reaction due to isotopic substitution.
An isotopic substitution does not influence the electronic potential energy surface.

hydrogen (^1H) - deuterium (^2H) - tritium (^3H)
carbon (^{12}C , ^{13}C), nitrogen (^{14}N , ^{15}N), oxygen (^{16}O , ^{18}O), chlorine (^{35}Cl , ^{37}Cl)



Different bond dissociation energies for R-D and R-H: E_{D} is greater than E_{H}
This difference in energy due to isotopic replacement results in differing rates of reaction k – the rate determining step

The effect is measured in kinetic isotope effects:
The reaction rate for the conversion of R-D is slower than the reaction rate for the conversion of R-H.

Kinetic Isotope Effects

Isotope replacement does not change the electronic structure of the molecule or the potential energy surfaces of the reactions

The greater the mass the more energy is needed to break bonds. A heavier isotope forms a stronger bond - less of a tendency to dissociate. The increase in energy needed to break the bond results in a slower reaction rate and the observed isotope effect:

the ratio $k_H/k_D = 1$ to 7

Heavy atom isotope effects = the substitution of carbon, oxygen, nitrogen, sulfur, and bromine, with effects that are much smaller and are usually between 1.02 and 1.10. The maximum isotopic rate ratio is proportional to the square root of the inverse ratio of isotopic masses.

Secondary kinetic isotope effects = rate changes due to isotopic substitutions at a site other than the bond breaking site in the rate determining step of the reaction, α , β , and γ effects

Kinetic Isotope Effects

Harmonic oscillator frequency
Bond stretching frequency

$$\nu = \frac{1}{2\pi} \sqrt{\frac{k}{\mu}}$$

Reduced mass

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$

bond	μ
C-H	0.92
C-D	1.71
¹² C- ¹² C	6.00
¹² C- ¹³ C	6.24

Zero point energy

$$E^0 = \frac{1}{2} h \nu$$

Reaction rate

$$k = A e^{-\frac{E^0}{kT}}$$

Isotope effect

$$\frac{k_H}{k_D} = e^{\frac{h(\nu_D - \nu_H)}{2kT}}$$

Pauling's Rules


Five principles which could be used to determine the structures of complex ionic/covalent crystals

Pauling's Rule no. 1 Coordination Polyhedra


A coordinated polyhedron of anions is formed about each cation.

Cation-Anion distance is determined by sums of ionic radii.


Cation coordination environment is determined by radius ratio.



1 Metallic radius




2 Covalent radius




3 Ionic radius

$$r(O^{2-}) = 140 \text{ pm} \text{ (Linus Pauling)}$$

Periodic Trends in Atomic Radii



Coordination Polyhedra



Bond Length

Pauling $R(A - B) = r_A + r_B$

Schomaker-Stevenson Rule

polar bonds are shorter than the sum of covalent radii

Original $c = 9 \text{ pm}$, $n = 1$

Modified $c = 8.5 \text{ pm}$, $n = 1.4$

$$R(A - B) = r_A + r_B - c |\chi_A - \chi_B|^n$$

Allred-Rochow electronegativities give the best fit

Ionic Radii

R.D. Shannon and C.T. Prewitt, Acta Cryst. B25, 925-945 (1969)

R.D. Shannon, Acta Cryst. A32, 751-767 (1976)

As the coordination number (CN) increases, the Ionic Radius increases

Sr^{2+}

CN Radius, Å

6 1.32

8 1.40 As the oxidation state increases, cations get smaller
9 1.45 (6-fold coordination, in Å)

10 1.50

12 1.58 Mn^{2+} 0.810
 Mn^{3+} 0.785
 Mn^{4+} 0.670

Ti^{2+} 1.000
 Ti^{3+} 0.810
 Ti^{4+} 0.745

Ionic Radii

The radius increases down a group in the periodic table.

The exception - 4d/5d series in the transition metals - the lanthanide contraction

(6-fold coordination, in Å)

Al^{3+} 0.675

Ga^{3+} 0.760

In^{3+} 0.940

Tl^{3+} 1.025

Right to left across the periodic table the radius decreases.

Ti^{4+} 0.745

(6 coordinate radii, in Å)

Zr^{4+} 0.86

Hf^{4+} 0.85


La^{3+} 1.172

Nd^{3+} 1.123

Gd^{3+} 1.078

Lu^{3+} 1.001

Cation/Anion Radius Ratio



CN	r/R
12 – hcp/ccp	1.00 (substitution)
8 – cubic	0.732 – 1.00
6 – octahedral	0.414 – 0.732
4 – tetrahedral	0.225 – 0.414

optimal radius ratio
for
given CN
ions are in touch

Pauling's Rules

Pauling's Rule no. 2 Bond Strength

The strength of an electrostatic bond: $s_{ij} = \text{valence} / \text{CN}$

The bond valence sum of each ion equals its oxidation state.

The valence of an ion (V_i , equal to the oxidation state of the ion) is equal to a sum of the valences of its bonds (s_{ij}).

In a stable ionic structure the charge on an ion is balanced by the sum of electrostatic bond strengths (s_{ij}) to the ions in its coordination polyhedron.

TiO_2 (Rutile) Ti - oxidation state of +4, coordinated to 6 oxygens.


$$V_{\text{Ti}} = +4 = 6 (s_{ij}) \quad s_{ij} = +2/3$$

The bond valence of oxygen, coordinated by 3 Ti atoms

$$V_{\text{O}} = 3 (s_{ij}) = 3 (-2/3) = -2$$

Each bond has a valence of s_{ij} with respect to the cation

and $-s_{ij}$ with respect to the anion.



Bond Strength

Brown, Shannon, Donnay, Allmann:

Correlation of the valence of a bond s_{ij} with the (experimental) bond distance d_{ij} .

$$s_{ij} = \exp \frac{R_{ij} - d_{ij}}{b}$$

R_{ij} = standard single bond length - determined empirically from (many) structures where bond distances and ideal valences are accurately known.

Tables of R_{ij} values for given bonding pairs (i.e. Nb-O, Cr-N, Mg-F, etc.) have been calculated, just as tables of ionic radii are available.

A constant $b = 0.37$

Bond Strength

Correlation of the valence of a bond s_{ij} with the (experimental) bond distance d_{ij} .

$$s_{ij} = \exp \frac{R_{ij} - d_{ij}}{b}$$

$$v_i = \sum s_{ij} = \sum \frac{z_i}{CN}$$

Use of the bond valence concept


- A) To check experimentally determined structures for correctness, or bonding instabilities
- B) To predict new structures
- C) To locate light atoms such as hydrogen or Li ion, which are hard to find experimentally
- D) To determine ordering of ions which are hard to differentiate experimentally, such as Al^{3+} and Si^{4+} , or O^{2-} and F^-
- E) To check/confirm oxidation states of atoms ($\text{Co}^{2+}/\text{Co}^{3+}$, $\text{Fe}^{2+}/\text{Fe}^{3+}$)

Pauling's Rules

Pauling's Rule no. 3 Polyhedral Linking

The presence of shared edges, and particularly shared faces decreases the stability of a structure. This is particularly true for cations with large valences and small coordination numbers.


Avoid shared polyhedral edges and/or faces.



Polyhedral Linking

The Coulombic interactions - maximize the cation-anion interactions (attractive), and minimize the anion-anion and cation-cation interactions (repulsive).

- increasing the coordination number
- decreasing the cation-anion distance



The cation-cation distance decreases, (the Coulomb repulsion increases) as the

- degree of sharing increases (corner < edge < face)
- CN decreases (cubic < octahedral < tetrahedral)
- cation oxidation state increases (this leads to a stronger Coulomb repulsion)


Pauling's Rules

Pauling's Rule no. 4 Cation Evasion


In a crystal containing different cations those with large valence and small coord. number tend not to share polyhedral elements (anions).

Perovskite, CaTiO_3

Ca^{II} 12-coordinate CaO_{12} cuboctahedra share **FACES**



Ti^{IV} 6-coordinate TiO_6 octahedra share only **VERTICES**



Pauling's Rules

Pauling's Rule no. 5 Environmental Homogeneity the rule of parsimony

The number of chemically different coordination environments for a given ion tends to be small.

Once the optimal chemical environment for an ion is found, if possible all ions of that type should have the same environment.

Covalent Radius of Fluorine

1938 Brockway - electron diffraction on $F_2(g)$
 $R(F-F) = 145 \text{ pm}$ $d(F) = 73 \text{ pm}$


1960 Pauling – backbonding in A–F

1990 Reed and Schleyer – pi bonding in A–F


1992 Gillespie and Robinson - 54 pm
Longer if a lone pair on EAF_n
Shorter in an incomplete octet BF_3

1997 Gillespie - 60 pm, the X–F bond length decreases with a decreasing CN, the smaller the bond angle ($\angle F-A-F$) the longer the bond length

2009 Pyykkö - 64 pm in a single bond, 59 pm in a double bond and 53 pm in triple bond character



Covalent Radius of Fluorine



Period 2

Fluorine radii decrease across the 2nd period with increasing electronegativity of the central atom and therefore with decreasing ligand charge.


Period 3

The six-coordinated fluorides fall on the same smooth curve as for the Period 2 fluorides = close-packed.

The 3- and 4-coordinated fluorides of the Period 3 elements do not fall on this curve = not close packed.

Additive Covalent Radii

Additive covalent radii = approximate a bond length as the sum of two atomic radii
Predominantly covalent



the A–B bond is not too ionic

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----

1 H 32	3 Li 133	4 Be 102	5	6	7	8	9	10	11	12	13	14	15	16	17	2 He 46	
-	-	-														-	
3 Li 124	4 Be 90	5	6	7	8	9	10	11	12	13	14	15	16	17	18		
-	85																
11 Na 155	12 Mg 139	13 V 112	14 Cr 111	15 Mn 105	16 Fe 109	17 Co 103	18 Ni 96	19 Cu 101	20 Zn 120	21 Ga 117	22 Ge 111	23 As 114	24 Se 107	25 Br 106	26 Kr 107		
160	132	108	106	103	103	102	101	120	-	121	114	106	107	110	108		
-	127																
19 K 196	20 Ca 171	21 Sc 148	22 Ti 136	23 V 134	24 Cr 122	25 Mn 119	26 Fe 116	27 Co 111	28 Ni 110	29 Cu 112	30 Zn 118	31 Ga 124	32 Ge 121	33 As 121	34 Se 116	35 Br 114	36 Kr 117
193	147	116	117	112	111	105	109	103	101	115	120	117	111	114	107	109	121
-	133	114	108	106	103	103	102	96	101	120	-	121	114	106	107	110	108
37 Rb 210	38 Sr 185	39 Y 163	40 Zr 154	41 Nb 147	42 Mo 138	43 Tc 128	44 Ru 125	45 Rh 125	46 Pd 120	47 Ag 128	48 Cd 136	49 In 142	50 Sn 140	51 Sb 140	52 Te 136	53 I 133	54 Xe 131
202	157	130	127	125	121	120	114	110	117	139	144	136	130	133	128	129	135
-	139	124	121	116	113	110	103	106	112	137	-	146	132	127	121	125	122
55 Cs 232	56 Ba 196	La–Lu	72 Hf 152	73 Ta 146	74 W 137	75 Re 131	76 Os 129	77 Ir 122	78 Pt 123	79 Au 124	80 Hg 133	81 Tl 144	82 Pb 144	83 Bi 151	84 Po 145	85 At 147	86 Rn 142
209	161		128	126	120	119	116	115	112	121	142	142	135	141	135	138	145
-	149		122	119	115	110	109	107	110	123	-	150	137	135	129	138	133
87 Fr 223	88 Ra 201	Ac–Lr	104 Rf 157	105 Db 149	106 Sg 143	107 Bh 141	108 Hs 134	109 Mt 129	110 Ds 128	111 Rg 121	112 Cn 122	113	114 Fl 143	115	116 Lv 162	117	118
218	173		140	136	128	128	125	125	116	116	137		136	162	175	165	157
-	159		131	126	121	119	118	113	112	118	130						

57 La 180	58 Ce 163	59 Pr 137	60 Nd 137	61 Pm 135	62 Sm 134	63 Eu 134	64 Gd 135	65 Tb 135	66 Dy 132	67 Ho 133	68 Er 133	69 Tm 131	70 Yb 131	71 Lu 131
139	137	138	137											
139	131	128												
89 Ac 186	90 Th 175	91 Pa 169	92 U 170	93 Np 171	94 Pu 172	95 Am 166	96 Cm 166	97 Bk 168	98 Cf 168	99 Es 165	100 Fm 167	101 Md 173	102 No 176	103 Lr 161
153	143	138	134	136	135	135	136	136	139	140	140	139	139	141
140	136	129	118	116										

The Same or Not the Same

Three sigma criterion



$$\Delta = A \pm B$$

$$\sigma_{\Delta} = \sqrt{\sigma_A^2 + \sigma_B^2}$$

$$\Delta < 3\sigma_{\Delta} \quad A, B - \text{are identical}$$


$$\Delta > 3\sigma_{\Delta} \quad A, B - \text{are different}$$

Crystallographically imposed
 C_3 axis??




Al-N(1) 1.983(6)	Si(1)-N(3) 1.697(6)
Al-N(2) 1.805(5)	Si(2)-N(2) 1.684(5)
Al-N(3) 1.812(6)	Si(3)-N(4) 1.690(6)
Al-N(4) 1.810(6)	

Ionic Character vs. Electronegativity




$$i = 1 - \exp [-0.21(\chi_A - \chi_B)^2]$$


van Arkel-Ketelaar Triangle



van Arkel-Ketelaar Triangle



Normal vs. Dative Bond




	13	14	15	16	
5	B boron 10.81 [10.806, 10.821]	C carbon 12.011 [12.009, 12.012]	N nitrogen 14.007 [14.006, 14.008]	O oxygen 15.999 [15.999, 16.000]	
13	Al aluminium 26.982 [28.084, 28.086]	Si silicon 28.085 [28.084, 28.086]	P phosphorus 30.974 [32.059, 32.076]	S sulfur 32.06 [32.059, 32.076]	
2	n [10]	Ga gallium	Ge germanium	As arsenic	Se selenium
0					


Normal vs. Dative Bond

Heterolytic Bond Cleavage

Charged + Diamagnetic



Neutral + Diamagnetic



Neutral + Radicals

Charged + Radical

Homolytic Bond Cleavage

Badger's Rule

Badger's rule relates vibrational frequencies to bond distances for various atomic pairs.

$$r_e = \left(\frac{C_{ij}}{k} \right)^{\frac{1}{3}} + d_{ij}$$

r_e = equilibrium bond distance

k = force constant (can be substituted with ν , the stretching frequency)


C_{ij} and d_{ij} = empirically fitted values for a particular atomic pair i-j

Plots of bond distances vs $\nu^{-2/3}$ or $k^{-1/3}$ for a set of compounds comprising a particular atomic pair are fit to equation to determine C_{ij} and d_{ij} , which then allows a newly measured force constant or vibrational frequency to be used to calculate an unknown bond distance (or vice versa).

$$\nu = \frac{1}{2\pi} \sqrt{\frac{k}{\mu}} \Rightarrow k = \mu(2\pi\nu)^2$$

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$


Badger's Rule



$$r_e = \left(\frac{C_{ij}}{k} \right)^{\frac{1}{3}} + d_{ij}$$

$$\nu = \frac{1}{2\pi} \sqrt{\frac{k}{\mu}}$$


Badger's Rule



$$r_e = \left(\frac{C_{ij}}{\mu(2\pi\nu)^2} \right)^{\frac{1}{3}} + d_{ij}$$


Normalized Badger's Rule

Normalized frequency



$$r_e = \left(\frac{181}{\nu_\mu} \right)^{1/3} + 0.259$$

Normalized Badger's Rule



Bond Order

Not an observable quantity

G. N. Lewis 1916

Electron sharing between two atoms

Bond order = a measure of the number of electrons shared between two atoms.

Resonance structures

G. Herzberg

Bond order is equal to half the difference between the number of bonding electrons and the number of ‘loosening electrons’ in the molecule.

R. S. Mulliken

A non-integral bonding power of either sign is attributable to every outer electron - the Mulliken population analysis.

The effective bond order (EBO)

Each bonding orbital i has a natural orbital (NO) occupation number b_i . The corresponding antibonding orbital has the occupation number ab_i .

$a = 0$ – full bond, $a = 1$ – no bond

$$\text{EBO} = \sum_i \frac{(b_i - ab_i)}{2}$$

Multiple Bonds

Dominant form of oxygen is O=O, a colourless paramagnetic gas.





Ozone (O₃) is the only other allotrope.

Why does not it form O₈?

Lone-pair repulsion across the short O-O distance.

O is small, and its orbitals highly contracted, and so electrons feel a lot of e-e repulsion.

Bond energies



S=S is thermodynamically unstable wrt -S-S- (*catenation*) but stable above 700 °C (Why?)

π -Bonds

Diffuseness of orbitals

2s and 2p orbitals are only shielded from the nucleus by the 1s electrons.

Therefore they are highly contracted, and π -overlap is efficient for the second period elements.

3s and 3p orbitals in the 3rd row and higher elements are more diffuse, and so this side-on p-p overlap is less effective.

Repulsion between core electrons

Second-row elements have only two core electrons. Third-row and heavier elements have a significantly larger number of core electrons that cause repulsion between two heavy elements bonded to each other. This prevents the atoms to approach each other closer and form stronger π bonds.


Atomic size

the large jump (50%) in the covalent radii between the 2nd and 3rd rows of the periodic table (for example, C (0.77 Å), Si (1.17 Å). Higher coordination numbers observed for the 3rd row.

Thermochemistry


π -bond energies in heavy double bonds too small to prevent polymerization.

Atomic Radii



B. Cordero et al. *Dalton Trans.*, 2008, 2832–2838

π -Bonds



Atom n	C 2	Si 3	Ge 4	Sn 5	Pb 6
Orbital energy					
ns	-19.39	-14.84	-15.52	-13.88	-15.41
np	-11.07	-7.57	-7.29	-6.71	-6.48
Ionization energy					
ns	16.60	13.64	14.43	13.49	16.04
np	11.26	8.15	7.90	7.39	7.53
Electron affinity					
Polarizability	1.26	1.39	1.23	1.11	0.36
Electronegativity	1.76	5.38	6.07	7.7	6.8
Mulliken					
Pauling	1.92	1.46	1.40	1.30	1.21
Allen	2.55	1.90	2.01	1.96	2.33
Atomic radius					
ns	2.28	1.76	1.81	1.68	1.91
np	1.58	2.20	2.19	2.48	2.39
	1.74	2.79	2.88	3.22	3.22

Carbon - 2s and 2p electrons approximately **the same spatial extension** (the core electrons occupy only the 1s orbital).

Heavier Group 14 elements, the np valence electrons ($n > 2$) are spatially separated from ns due to Pauli repulsion with the $(n-1)p$ electrons in the inner shell.

Difficult to hybridize ns and np orbitals for heavier atoms, tend to preserve the valence ns electrons as core-like electrons while carbon shows a preference for an effective hybridization of the s and p - the strong overlap binding ability.

π -Bonds

In Group 14 all E=E species are unstable!




	kJ mol^{-1}	E=E	E-E	ΔH
C	602	—2(356)	—110	
Si	310	—2(226)	—142	
Ge	270	—2(188)	—106	
Sn	190	—2(151)	—112	

C=C double bonds are thermodynamically less stable than two corresponding single bonds and are kinetically stabilized. The preparation of polymers involves a catalyst to overcome the kinetic barrier and induce an exothermic reaction.

The Si=Si bond are much weaker, the driving force for polymerization is large, hard to prevent polymerization.

Stabilisation of Si=Si



Bulky groups can inhibit catenation

Si=Si 216.0 pm

a typical Si–Si 232 pm - 8% shortening

Tetraphenylethene - C=C 12% shortening

the Z → E isomerization energy of unsymmetrical disilene, (Mes)(tBu)Si=Si(tBu)(Mes)

the barrier to inversion $E_{act} = 131 \text{ kJ mol}^{-1}$

trans-stilbene only 179 kJ mol⁻¹



The barrier in the disilene is high

West, R.; Fink, M.J.; Michl, J. *Science* **1981**, *214*, 1343

Masamune, S. et al. *J. Am. Chem. Soc.* **1982**, *104*, 1150

π -Bonds


Differences in reactivity?






Triplet - Singlet

R_2C carbene triplet
 R_2Sn singlet



triplet (T)



singlet (S)



Stabilisation of Sn=Sn



Distannene = the first isolable species with multiple bonding between two heavier main group elements.

Sn-Sn distance = 2.768(1) Å

Sn elemental = 2.80 Å



$\text{Ph}_3\text{Sn}-\text{SnPh}_3$ = 2.770(4) Å

Lappert. M.F. et al. *J. Chem. Soc., Chem. Commun.* **1976**, 261





Distannene Sn=Sn

Double Bond in Heavy Elements



Compound	M–M/Å	$\delta^*/^\circ$	$\gamma^*/^\circ$
[Ge(C ₆ H ₃ Et ₂ -2,6)] ₂	2.213(2)	12	10
[Ge(C ₆ HMe ₃ -2,3,4-Bu ^t -6)] ₂	2.2521(8)	0	20.4
[Ge(SiMePr ⁱ) ₂] ₂	2.267(1)	0	6.5
[Ge(SiPr ⁱ) ₂] ₂	2.298(1)	0	16.4
[Ge(Mes)(C ₆ H ₃ Pr ⁱ -2,6)] ₂	2.301(1)	36	7
[Ge{CH(SiMe ₃) ₂ } ₂] ₂	2.347(2)	32	0
[GeCl(C ₆ H ₃ Mes ₂ -2,6)] ₂	2.443(2)	39	0
{GeN(Bu ^t)(CH ₂) ₃ N(Bu ^t)SiN(Bu ^t)(CH ₂) ₂ N(Bu ^t)} ₂	2.451(2)	41.3	42.3
[{Ge(SiBu ^t) ₃ } ₂ Ge(SiBu ^t) ₃] ₂	2.239(4)	—	—
[{Ge(SiBu ^t) ₃ } ₃][BPh ₄] ₂	2.226(4)	—	—
[Ge(C ₆ H ₃ Mes ₂ -2,6)] ₃	2.35(7)	—	—
K[{Ge(C ₆ H ₃ Mes ₂ -2,6)} ₃] ₂	2.422(2)	—	—
[Sn{CH(SiMe ₃) ₂ } ₂] ₂	2.768(1)	41	0
[Sn{Si(SiMe ₃) ₃ } ₂] ₂	2.8247(6)	28.6	63.2
[Sn{C ₆ H ₂ (CF ₃) ₃ -2,4,6} {Si(SiMe ₃) ₃ }] ₂	2.833(1)	41.5	0
[Sn(C ₆ HMe ₃ -2,3,4-Bu ^t -6)] ₂	2.910(1)	21.4, 64.4	—
[Sn{C ₆ H ₂ (CF ₃) ₃ -2,4,6}] ₂	3.639(1)	46	0
[K(THF) ₆][{SnC ₆ H ₃ Trip ₂ -2,6}] ₂	2.8123(9)	95.20	0
[Pb{C ₆ H ₂ (CF ₃) ₃ -2,4,6} {Si(SiMe ₃) ₃ }] ₂	3.537(1)	40.8	0



Double Bond in Heavy Elements



Triple Bond - Disilyne







$\text{Si}\equiv\text{Si}$ 2.0622(9) Å

half the magnitude of the bond shortening of alkynes/alkenes

the substituents at the $\text{Si}\equiv\text{Si}$ group are trans-bent, a bond angle of 137.44(4)°

Heavier Gr14 Alkyne Analogues


Compound	M≡M (\AA)	M≡M-C bending angle (degrees)	Shortening* (%)
	2.0622(9) (ref. 21)	137.44(4)	11.87
	2.285(6) (ref. 22)†	128.67(8)	6.35
	2.6675(4) (ref. 23) †	125.1(2)	5.07
	3.1811 (ref. 24)‡	94.26(4)	-9.69

* Shortening with respect to a single bond.

† Ar' = C₆H₃-2,6(C₆H₃-2,6-*i*-Pr₂)₂.

‡ Ar* = C₆H₃-2,6(C₆H₂-2,4,6-*i*-Pr₃)₂.


Heavier Gr14 Alkyne Analogues






The $6s^2$ valence electrons are stabilized by relativistic effects - they participate less in bonding.


The Pb-Pb single bond formed by head-to-head overlap of 6p orbitals, which have a larger radius than f 6s and hence form longer bonds

Vinylidene / Acetylene Isomers



Triple M≡M Bond





Triple M≡M Bond


	Configuration	rM-M / pm	Magnetism
Cr ₂ Cl ₉ ³⁻	$\sigma^2(\delta/\pi)^4$	310	Paramagnetic
Mo ₂ Cl ₉ ³⁻	$\sigma^2(\delta/\pi)^4$	253-288	Variable
W ₂ Cl ₉ ³⁻	$\sigma^2(\delta/\pi)^4$	242-250	Diamagnetic

Cr₂Cl₉³⁻ two CrCl₃ fragments held together by 3 bridging Cl, no d-orbital overlap, no direct Cr-Cr bond, paramagnetic with 6 unpaired electrons.

Mo₂Cl₉³⁻ the Mo-Mo distance depends on the cation present in the crystal structure, variable bonding and magnetism.

W₂Cl₉³⁻ good overlap and a triple W-W bond with no unpaired electrons.


Triple M≡M Bond



$\text{Mo}_2(\text{NMe}_2)_6$
a $\sigma^2\pi^4$ triple bond
a short Mo–Mo bond length of 2.214(2) Å





M.H. Chisholm, F.A. Cotton, B.A. Frenz, W.W. Reichert, L.W. Shive, B.R. Stults, J. Am. Chem. Soc. 98 (1976) 4469

$\text{W}_2(\text{NMe}_2)_6$
a W–W bond length of 2.294(1) Å




M.H. Chisholm, F.A. Cotton, M. Extine, B.R. Stults, J. Am. Chem. Soc. 98 (1976) 4477

Triple M≡M Bond in Gr13




N. Holzmann, A. Stasch, C. Jones, G. Frenking, *Chem. Eur. J.* **2011**, *17*, 13517.

Triple E \equiv E Bond in Gr13




ground state of B_2



ground state of E_2 ($E = Al-In$)





Triple E \equiv E Bond in Gr13




The ground state, seen in ligand free B_2 , has two doubly occupied bonding orbitals (blue) but one doubly occupied antibonding orbital (red), giving a bond order of 1 and hence a single bond.


In contrast, the third excited state observed in $(NHC) \rightarrow B \equiv B \leftarrow (NHC)$ has three doubly occupied bonding orbitals and hence a triple bond.

Triple E \equiv E Bond in Gr13






Braunschweig et al., *Science* **2012**, *336*, 1420.





Braunschweig et al., *Angew. Chem. Int. Ed.* **2014**, *53*, 9082.



Braunschweig et al., *Angew. Chem. Int. Ed.* **2015**, online.

Quadruple Bond

1964 – F. A. Cotton – the first bond order higher than 3



$\delta-\delta^*$ transition $14700 \text{ cm}^{-1} (1.82 \text{ eV})$

EBO = 3.2

the sum of the partial bond orders 0.92 (σ), 1.74 (π) and 0.54 (δ)

Formal Shortness Ratio (FSR)

Bond	Ratio	Bond	Ratio
$\text{C}\equiv\text{C}$	0.783	$\text{Cr}\equiv\text{Cr}$	0.771
$\text{N}\equiv\text{N}$	0.786	$\text{Mo}\equiv\text{Mo}$	0.807
		$\text{Re}\equiv\text{Re}$	0.848


A short bond?

The FSR is a dimensionless number given by the ratio of the atom–atom distance, d , of a bond and the sum of the radii of the two atoms involved, $r_A + r_B$:

$$\text{FSR} = d/(r_A + r_B)$$



The advantage of this formalism is its interelement applicability. The FSR is a useful tool for comparing formal quintuple bonding and short metal–metal distances

Quadruple Bond



Quadruple Bond

the Mo–Mo distances changes as torsional strain is introduced by bridging diphosphines with concomitant diminution of the d_{xy} – d_{xy} overlap, i.e., the δ bond.




internal twist

rotation about the metal–metal bond
average of 4 torsion angles


$\chi = 0^\circ$ - an eclipsed configuration (b.o. = 4)

$\chi = 45^\circ$ - a staggered configuration (b.o. = 3)

$\text{Mo}_2\text{X}_4(\text{PR}_3)_4$ and $\text{Mo}_2\text{X}_4(\text{diphosphine})_2$
 $\text{X} = \text{Cl}$ and Br



Quadruple Bond in PES



PES spectrum of the $[\text{Re}_2\text{Cl}_8]^{2-}$ anion

$$1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$$

Oxidation $\text{Mo}_2(\text{TiPB})_4$ to $[\text{Mo}_2(\text{TiPB})_4]^+$

$\text{Mo}_2^{4+} \rightarrow \text{Mo}_2^{5+}$
one unpaired electron (EPR)

Mo–Mo distance 0.06 Å longer
electron removed from a bonding δ orbital

$$\text{b.o.} = 3.5$$

Quintuple Cr-Cr Bonding




Strategies for stabilization of multiple bonds

Maximum of valence electrons at metal

Minimum number of attached ligands

Bulky ligands to prevent intermolecular reactions

Monovalent m-terphenyl ligands



Bond lengths:

$\text{Cr-Cr} = 1.8351 \text{ \AA}$


$\text{Cr(1)-C(1A)} = 2.131 \text{ \AA}$

$\text{Cr(1)-C(7A)} = 2.2943 \text{ \AA}$


Bond angle: $\text{Cr(1A)-Cr(1)-C(1)} = 102.78^\circ$

T. Nguyen, A. D. Sutton, M. Brynda, J.C. Fettinger, G. J. Long, P. P. Power, *Synthesis of stable compound with fivefold bonding between two chromium(I) centers*, Science 2005, 310, 844. Doi 10.1126/science.1116789

Quintuple Cr-Cr Bonding





in THF, Ar' = C₆H₃-2,6(C₆H₃-2,6-iPr₂)₂




Quintuple Cr-Cr Bonding

$$\text{Cr-Cr bonding} = \sigma + 2\pi + 2\delta$$



Quintuple bond = five electron pairs play a role in holding the metal atoms together. It does not imply that the bond order is five or that the bonding is very strong since the ground state of the molecule mixes with higher energy configurations with less bonding character. This gives lower, usually non-integer, bond orders.

Quintuple Cr-Cr Bonding



Cr-Cr



R = Mes 1.7404(8) Å

R = Xylyl 1.7397(9) Å

Cottons “formal shortness ratio” (FSR)


FSR = 0.733

the FSR of dinitrogen is 0.786




Quintuple Cr-Cr Bonding

The Hein–Cotton concept
three-atom bridging ligands to establish short metal–metal distances






Importance of the ligand in establishing short metal–metal bonds:
The N–N distances correlate with the Cr–Cr bond lengths.

The Shortest M–M Bond



a

Cr-Cr, 1.7293(12) Å




5


Steric pressure

a bulky substituent at the bridging carbon
a further reduced M-M distance
Guanidinates π system is delocalized and planar

Quintuple Mo-Mo Bond



Bonding in R-MM-R



R. Hoffmann - the potential energy surfaces of RMMR, R = H, F, Cl, Br, CN, and CH₃; M = Cr, Mo, and W.


The potential energy surface is complex with several local minima.

The linear geometry is always a minimum, but almost never the global minimum;

A preference for either a trans-bent conformation or one of the R groups bridging

The potential energy surface is relatively flat, the lowest energy conformation - the best compromise between maximization of the MM bonding and minimization of orbital interactions that are MR antibonding
the MM quintuple bond persists

Bonding in R-MM-R



R. Hoffmann - the potential energy surfaces of RMMR, R = H, F, Cl, Br, CN, and CH_3 ; M = Cr, Mo, and W.

The potential energy surface is complex with several local minima.

The linear geometry is always a minimum, but almost never the global minimum;

A preference for either a trans-bent conformation or one of the R groups bridging

The potential energy surface is relatively flat, the lowest energy conformation - the best compromise between maximization of the MM bonding and minimization of orbital interactions that are MR antibonding

the MM quintuple bond persists

Sextuple Bonding in Gr₆ M₂ Dimers

M₂ (M = Cr, Mo, W)

Matrix isolation

IR and UV spectroscopy

Short M-M distances


Singlet ground state

Laser-evaporated Cr₂ in the gas phase

d(Cr-Cr) = 1.68 Å

Spectroscopic studies of Cr₂ generated from pulsed photolysis of Cr(CO)₆
d(Cr-Cr) = 1.71 Å



Mo-Mo molecule in the gas phase at low temperatures
d(Mo-Mo) = 1.93 Å



Sextuple Bonding in Gr6 M₂ Dimers

p-orbitals too high in energy

M₂ (M = Cr, Mo, W)




overlap **increases down a group** and can reach the bonding manifold

δ-bonds are weaker than π-bonds due to poor overlap between orbitals.

M-M bonding energy increases down a group which is in contrast to the p-block.


Sextuple Bonding in Gr₆ M₂ Dimers



Cr_2 – poor delta bonding, 4 AF coupled electrons, larger 4s orbitals generate a considerably longer bond than the 3d-3d. This unbalance weakens the 3d bonds and makes the 4s bond repulsive at equilibrium geometry, the repulsive interaction between the closed 3p, which have about the same radial extension as 3d.

Effective bond order (EBO)
Dissociation energy (DE)

M	EBO	d(M-M), Å	DE, eV
Cr	3.5	1.66	1.65
Mo	5.2	1.95	4.4
W	5.2	2.01	5.4



Multiple Bonding

The bond energy only 1.66 eV for Cr_2 sextuple bond
but 3.17 eV for quintuple bond in $[\text{PhCrCrPh}]$
= no direct relation between bond order and bond energy, a complex quantity depends on many factors, such as atomic promotion energy, the interplay between attractive nuclear forces and electron repulsion, spin–orbit coupling—decreases the bond energy

Bond orders higher than 6?

Lanthanides – 4f orbitals?

Actinides – 5f orbitals participate in bonding

The EBO is always smaller than the maximum formal bond order (MBO = the number of electrons forming the bond divided by two)

Molecule	MBO	EBO	D_0 [eV]
Cr_2	6	3.5	1.6
$[\text{PhCrCrPh}]$	5	3.5	3.2
Ac_2	3	1.7	1.2
Th_2	4	3.7	3.3
Pa_2	5	4.5	4.0
U_2	6	4.2	1.2

Multiple Bonding Trends In Groups

Main-Group Elements

Tendency to multiple bonding **decreases down the group**

s + 3×p orbitals available

s/p size difference increases and propensity to hybridization decreases down the group

Transition Metals

Multiple bonding **increases down the group**

s + 5×d orbitals available

difference in size between the smaller nd and larger (n+1)s orbitals decreases down the group


relativistic effects contract s and p orbitals while d and f are expanded

the s and d orbitals in second- and especially third-row transition metals are more equal in size. This greater equivalency in size considerably enhances the bond strength

Inverted Bond

isoelectronic molecules C_2 , CN^+ , BN, and CB^- singlet ground state $X^1\Sigma_g^+$

valence bond (VB) theory and full configuration interaction (FCI)




bond order = 2

$2\sigma_g$ and $2\sigma_u$ orbitals are both filled, two π -bonds unsupported by an underlying σ -bond and two σ lone pairs.

bond order = 3


sp-hybridized carbons, one σ - and two π -bonds, two electrons in the outwardly pointing hybrids



triplet state $c^3\Sigma_u^+$, electrons are unpaired, 26.4 kcal mol⁻¹ above the ground state
these electrons maintain a significant bonding interaction in the ground state

Sigma Hole Interactions


Electrostatic potential (ESP) – observable quantity



Halogen/chalcogen/pnictogen/tetrel bonding

= a noncovalent interaction between a covalently-bonded atom of Groups 14–17 and a negative site, *e.g.* a lone pair of a Lewis base or an anion.

σ -hole = a region of positive electrostatic potential on the extension of one of the covalent bonds to the atom caused by the anisotropy of the atom's charge distribution




Heavy atoms without hybridization
4p_z occupied by only one electron

Sigma Hole Interactions

Electrostatic potential (ESP)

$$V(\mathbf{r}) = \sum_{\text{A}} \frac{Z_{\text{A}}}{|\mathbf{R}_{\text{A}} - \mathbf{r}|} - \int \frac{\rho(\mathbf{r}') d\mathbf{r}'}{|\mathbf{r}' - \mathbf{r}|}$$



Bond strength 13 – 100 kJ mol⁻¹
H-bond in (H₂O)₂ 20 kJ mol⁻¹

Directionality increase Cl < Br < I

Bond-Stretch Isomers

Isomerism


- the *molecular conformation* specifies a combination of relative atomic positions conferring on the molecule a certain stability.
- the *potential energy (hyper)surface* (PES) characterizes through its minima the various molecular conformations that could be expected for a given assembly of atoms, and defines by means of saddle points the thermodynamic pathways interconnecting them.

Isomers = molecular conformations corresponding to distinct minima on the PES, separated by an energy barrier high enough to impede immediate interconversion at room temperature, separable, $100 - 120 \text{ kJ mol}^{-1}$ or more.

Conformers = barrier is lower, not separable under ambient conditions, but could be observed and characterized at lower temperature.

Bond-Stretch Isomers

the potential energy (hyper)surface (PES)




Bond-Stretch Isomers

R. Hoffmann 1972

Bond-stretch *isomers* = distinct, separable and stable at room temperature

Bond-stretch isomerism (BSI) = the unusual phenomenon whereby molecules differ only in the length of one or more bonds



NOT Bond-stretch *isomers*:


Complex $\text{Re}(\text{cis-Cl}_2)(\text{NCMe})(\text{NO})\text{-trans-(PMe}_3)_2$

Crystallizes in $P2_1/a$ with 44 molecules in the asymmetric unit

a dense and low-symmetry hydrogen bonding network

the Re-N(NO) and the Re-N(NCMe) distances vary by as much as 0.10 or 0.12 Å, respectively.

[2.2.2]propellane




S orbital = symmetric combination wrt σ_h , transannular C–C bond

A orbital = antibonding, antisymmetric


Interaction (through-bond coupling) between **A** and the high-lying σ^* orbitals of the C–C bonds (2–3, 5–6 and 7–8) results in a stabilization of **A**.

A crossing between **A** and **S**, **A** becomes the HOMO when the transannular distance is larger than 2.25 Å and gives rise to a diradical form.

The orbital crossing makes the interconversion between the diradical and the tricyclic form symmetry forbidden and should generate isomers = equilibrium conformations separated by an energy barrier



Not Bond-Stretch Isomers



**Table I. Selected Bond Lengths (Å) for
Mo(O)Cl₂(PM₂Ph)₃ and Mo(O)Cl₂(PEt₂Ph)₃^a**


	blue Mo(O)Cl ₂ (PM ₂ Ph) ₃	green Mo(O)Cl ₂ (PEt ₂ Ph) ₃
Mo–O	1.676(7)	1.803(11)
Mo–Cl(1) ^b	2.551(3)	2.426(6)
Mo–Cl(2) ^b	2.464(3)	2.479(5)
Mo–P(1)	2.500(3)	2.521(5)
Mo–P(2)	2.541(3)	2.582(6)
Mo–P(3)	2.558(3)	2.556(6)

^a Taken from ref 12a. ^b Cl(1) and Cl(2) are trans and cis to O, respectively.


Not Bond-Stretch Isomers

Distortional M=O isomers?


Blue



Green




- Mo(IV)
- Mo(V)
- Mo(VI)





1.675(3) Å

1.871(3) Å




Distribution of Mo=O distances (Å) for monooxo complexes in the oxidation states +4, +5, and +6.

Not Bond-Stretch Isomers




Yoon, K.; Parkin, G.; Rheingold, A. L.
J. Am. Chem. Soc. 1992, 114, 2210



Not Bond-Stretch Isomers

Crystallographic disorder
difficult to detect, results in the incorrect
determination of bond lengths and the
incorrect formulation of compounds

Cocrystallization of structurally related
molecules resulting in the formation of
single-crystal solid solutions




Yoon, K.; Parkin, G.; Rheingold, A. L. J. Am. Chem. Soc. 1992, 114, 2210

Chromium Dimer?

Valence electron configuration:

$(3d\sigma_g)^2(3d\pi_u)^4(3d\delta_g)^4(4s\sigma_g)^2$ sextuple bond, Cr–Cr distance: 1.6788 Å (2.5 Å in Cr metal)
Singlet, observed dissociation energy = 1.44 ± 0.05 eV

The optimal bonding regions are quite different for the 3d and 4s orbitals




1st minimum: $Re = 1.59$ Å; $De = 1.38$ eV

2nd minimum: $Re = 2.40$ Å; $De = 1.14$ eV

Balance between covalent bonding (d-d) at short distance and (s-s) bonding and antiferromagnetic coupling of the 3d electrons at long distance

Calculated 2nd minimum of the ground-state energy curve is extremely shallow corresponding isomer rather short-lived and difficult to trap and to characterize

Not Bond-Stretch Isomers





1: R=tBu, R'=iPr,

2: R=Dur, R'=iPr



3: R=tBu, R'=Ph,

4: R=Dur, R'=Et,

5: R=Ph, R'=Ph,




compound	1 $(i\text{Pr}_2\text{PB}t\text{Bu})_2$	2 $(i\text{Pr}_2\text{PBDur})_2$	3 $(\text{Ph}_2\text{PB}t\text{Bu})_2$	4 $(\text{Et}_2\text{PBDur})_2$	5 $(\text{Ph}_2\text{PBPh})_2$
$\delta^{31}\text{P}$ (ppm) ^{a)}	+3	-26	-56	-55	-54
$\delta^{11}\text{B}$ (ppm) ^{a)}	+25	-13	-12	-19	-13
B–B (Å)	2.57	2.24	1.99	1.89	1.83
τ (°)	180	130	118	115	114



Bond-Stretch Isomers

Planar P_2B_2 core, $B-B = 2.57 \text{ \AA}$



^{31}P solid-state NMR $\delta = 5.9 \text{ ppm}$ (diradical)



^{31}P solution $\delta = -28 \text{ ppm}$ (bicyclic)

^{11}B solution $\delta = -9 \text{ ppm}$ (bicyclic)

Changing populations
diradical \rightarrow bicyclic
ratio of 1:7 at r.t.

bicyclic – diradical
coalescence

Diradical
 $\delta = 4.0 \text{ ppm}$




$\delta = 32.2 \text{ and } 41.8 \text{ ppm}$
(A:X = 1:1)

M_2X_2 Rings

Isomers with or without ligand–ligand bonds

Electron transfer from ligands X to metal M



FEC =

8

6

4

Metal configuration

d^n

d^{n+1}

d^{n+2}

Bridge

$(X_2)^{x-}$

$(X_2)^{2-x}$

$(X_2)^{4-x}$

M

X

L_2

β

Δ_{XX}

Experimental data

Pd

Te

$(PEt_3)_2$

108

0.07

Pt

S

$(PpyPh_2)_2$

103

0.96

Pt

S

dppe

86

1.10

Pt

Se

$(PPh_3)_2$

100

0.70

Pt

Te

$(PEt_3)_2$

106

0.32

Pt

Te

$(PPh_3)_2$

100

0.32

Pt

Te

dppe

86


0.53

β is the LML bond angle, Δ_{XX} is the difference b/w the X–X distance and the atomic radii sum.⁹⁹

Linear Metal M-M-M Frameworks




Ox. state of M?




Linear Metal M-M-M Frameworks

symmetrical (*s*) Co₃ chain



unsymmetrical (*u*) Co₃ chain



less unsymmetrical at low temperature


both the *s* and the *u* isomers are in an $S = 1/2$ ground state at low temperature

Linear Metal M-M-M Frameworks

Linear triatomic system: M-M-M

the equivalent atomic orbitals of every type give rise to a set of three molecular orbital (MO) combinations:

- (i) a bonding MO, lowest in energy
- (ii) a nonbonding, antisymmetric, localized on the terminal atoms
- (iii) an antibonding MO



Linear Metal M-M-M Frameworks

Five orbital sets:

One σ set - d_{z^2} orbital combinations (z is collinear with the framework axis)

Two degenerate π sets - combinations of d_{xz} and d_{yz} orbitals


Two δ sets - combinations of d_{xy} and $d_{x^2-y^2}$ orbitals.

One δ set accepts the lone pairs of the equatorial dpa ligands

The 3d atomic orbitals are compact in space and the 3d–3d overlap between π and δ orbitals is weak except at supershort distances


The nine metal MOs belonging to the two π sets and to the remaining δ set are not split in energy

Linear Metal M-M-M Frameworks



Three-electron, three-center system

18 electrons in the π and δ orbital sets do not take part in any metal–metal interaction, *localized* on individual metal atoms
The bonding in the M-M-M fragment is exclusively due to the 3 σ electrons, which are *delocalized* over the metal framework



Three-electron, three-center system

12 electrons in the π/δ block singly occupied, the 10 unpaired electrons—including the one in the σ nonbonding orbital and shared between the terminal Cr atoms—are *spin-coupled-antiferromagnetic* interaction

Linear Metal M-M-M Frameworks

Structure $\Delta d_{\text{Cr-Cr}} = 0$	S value	Spin Coupling Atomic Spin Populations	Relative Energies
Symmetric $\Delta d_{\text{Cr-Cr}} = 0$	2	$\uparrow\uparrow\uparrow \text{---} \downarrow\downarrow\downarrow \text{---} \uparrow\uparrow\uparrow$ 3.52 -3.00 3.52	0.0
Non-symmetric $\Delta d_{\text{Cr-Cr}} = 0.106$	2	$\uparrow\uparrow\uparrow \text{---} \downarrow\downarrow\downarrow \text{---} \uparrow\uparrow\uparrow$ 3.60 -3.00 3.44	+0.97
Non-symmetric $\Delta d_{\text{Cr-Cr}} = 0.679$	2	$\uparrow\uparrow\uparrow \text{-----} \downarrow\downarrow \overline{\overline{\overline{\uparrow\uparrow\uparrow}}}$ 3.79 -2.27 2.50	+4.25 AF to bonding
Non-symmetric $\Delta d_{\text{Cr-Cr}} = 0.679$	2	$\uparrow\uparrow\uparrow \text{-----} \uparrow\uparrow\uparrow \overline{\overline{\overline{\downarrow\downarrow\downarrow}}}$ 3.84 2.17 -2.09	+10.12 Short = bonding Long = ferromg
Symmetric $\Delta d_{\text{Cr-Cr}} = 0$	5	$\uparrow\uparrow\uparrow \text{---} \uparrow\uparrow\uparrow \text{---} \uparrow\uparrow\uparrow$ 3.50 2.93 3.50	+30.8
state ($S = 5$) is destabilized by 30.8 kcal mol ⁻¹			


Bond-Stretch Isomers?

Co-Co-Co

A ground-state potential energy curve (PES) has only one shallow minimum corresponding to the *s* conformation

Both the shape of this PES and the symmetric position of its minimum are consequences of the three-electron, three-center bond

The crystal forces could influence the molecular geometry or population of a low-energy, high-spin excited state induce a temperature-dependent distortion of the framework




Cr-Cr-Cr

A shallow, symmetric PES governed by σ metal electrons
 π and δ electrons, though localized on the metal atoms take part in the bonding through their magnetic coupling

Spin State Isomers

$3d^4 - 3d^7$ cations in O_h complexes

LS, $S = 0$




$10Dq > \Pi$

$10Dq$ – ligand-field splitting


Π – electron pairing energy

$10Dq(LS) \neq 10Dq(HS)$

HS, $S = 2$




$10Dq < \Pi$




Spin Crossover (SCO)

a reversible change in the spin state of from low spin (LS) to high spin (HS) affected by the application of external stimuli:

- Temperature
- Pressure
- Photoexcitation
- Magnetic field
- Electric fields




Fe(II) complexes (O_h) - most abundant cases



$$r(\text{Fe}-\text{N}): 1.95-2.00 \text{ \AA} \quad 2.15-2.20 \text{ \AA}$$


the Fe-N bond lengths and orbital overlap change upon SCO, and therefore $10Dq$ is different for the LS and HS states of the same complex

Spin Crossover (SCO)



Spin State Isomers


Magnetometry



HS molar fraction γ

$$\gamma = \frac{\chi T - (\chi T)_{\text{LS}}}{(\chi T)_{\text{HS}} - (\chi T)_{\text{LS}}}$$

^{57}Fe Mössbauer Spectroscopy




Relativistic Effects

Einstein's special theory of relativity (1905)

$$m = m_0 / \sqrt{1 - (v/c)^2}$$

space-time; a single continuum (one entity)


Bohr atomic model (1913)



$$a_0 = \frac{\epsilon_0 h^2}{m Z e^2 \pi}$$

$$v_{electron} = \left(\frac{2\pi e^2}{nh} \right) Z$$

	1s electron velocity m/s	relativistic mass	relativistic radius contraction
H (Z = 1)	$v = 2.18 \cdot 10^6$ $v = 0.00727 \cdot c$	$m = 1.0000265 m_0$	~ 0 %
Au (Z = 79)	$v = 1.73 \cdot 10^8$ $v = 0.577 \cdot c$	$m = 1.23 m_0$	~ 20 %
Fm (Z = 100)	$v = 2.18 \cdot 10^8$ $v = 0.727 \cdot c$	$m = 1.46 m_0$	~ 30 %



a_0 – Bohr radius

ϵ_0 – permittivity of free space

h – Planck constant

m – mass of an electron


Z – atomic number


e – elementary charge

$c = 3 \cdot 10^8$ m/s


Relativistic Effects

- **direct effect:** stabilization (decrease) of *s*- and *p*-orbitals
- **indirect effect:** destabilization (expanding) of *d*- in *f*-orbitals due to the increased shielding by *s*- and *p*-orbitals









- color Cu, Au, Ag instead Cu, Ag, Au
- very high first ionization energy (Au 9.23 eV, Ag 7.58 eV, Cu 7.73 eV)




- the highest electron affinity of all metals: simple reduction to auride anion Au^- (Cs^+Au^- is known since 1931, it has CsCl structure) – **pseudohalogen**
- $\text{Au}_2(\text{g})$ dissociation energy: 221 kJ/mol



- gold is the most electronegative of all metals




Relativistic Atomic Radius and Bond-Length Contractions



Expected order: **Cu < Ag < Au**

Actual order: **Cu < Au < Ag**


$$\begin{array}{ll} d(\text{Ag-P}) & d(\text{Au-P}) \\ 2.44 \text{ \AA} & 2.35 \text{ \AA} \end{array}$$



$$\begin{array}{ll} d(\text{Ag-S}) & d(\text{Au-S}) \\ 2.35 \text{ \AA} & 2.30 \text{ \AA} \end{array}$$




Distance, Å	Ag	Au
In solid state $d(\text{M-M})$	2.889	2.885
Ionic radius for two-coordinate M^{1+}	1.33	1.25
Ionic radius for four-coordinate M^{1+}	1.46	1.37

$$\begin{array}{ll} d(\text{Ag-H}) & d(\text{Au-H}) \\ 1.62 \text{ \AA} & 1.52 \text{ \AA} \end{array}$$





$$\begin{array}{ll} d(\text{Ag-As}) & d(\text{Au-As}) \\ 2.66 \text{ \AA} & 2.59 \text{ \AA} \end{array}$$

Auropphilicity



L-Au⁺ isolobal to H⁺, R⁺



Attractive interactions Au¹⁺...Au¹⁺:

d(Au–Au)
3.27 Å

aurophilicity or auophilic interaction

two-coordinate Au 5d¹⁰ closed-shell interactions

energy comparable to hydrogen-bonding (5–10 kcal/mol)

distances shorter 2.50–3.50 Å than the sum of van der Waals radii




(3.80 Å) or d(Au-Au) in ccp (2.89 Å)

correlation effects + relativistic effects ~20 %

metallophilicity or metallophilic interactions



d¹⁰-d¹⁰ Au⁺, Ag⁺, Cu⁺, Hg²⁺, Pt⁰, Pd⁰, Tl³⁺, also Ir

Auophilic Interactions



Auration Reactions

L-Au⁺ isolobal to H⁺, R⁺




(a) Homoleptic substitution

C(AuL) ₄	—	—	—
[C(AuL) ₅] ⁺	[N(AuL) ₄] ⁺	[O(AuL) ₃] ⁺	—
[C(AuL) ₆] ²⁺	[N(AuL) ₅] ²⁺	[O(AuL) ₄] ²⁺	—
—	—	—	—
—	—	S(AuL) ₂	—
—	[P(AuL) ₄] ⁺	[S(AuL) ₃] ⁺	[Cl(AuL) ₂] ⁺
—	[P(AuL) ₅] ²⁺	[S(AuL) ₄] ²⁺	—
—	[P(AuL) ₆] ³⁺	[S(AuL) ₅] ³⁺	—
—	—	[S(AuL) ₆] ⁴⁺	—
—	—	—	—
—	[As(AuL) ₄] ⁺	[Se(AuL) ₃] ⁺	[Br(AuL) ₂] ⁺
—	—	[Te(AuL) ₃] ⁺	—



(b) Heteroleptic substitution

RC(AuL) ₃	—	—	—
—	[R ₂ N(AuL) ₂] ⁺	—	—
—	[RN(AuL) ₃] ⁺	—	—
[R ₂ C(AuL) ₃] ⁺	[RN(AuL) ₄] ²⁺	—	—
—	[RP(AuL) ₃] ⁺	[RS(AuL) ₂] ⁺	—
—	[RP(AuL) ₄] ²⁺	[RS(AuL) ₃] ²⁺	—
—	[R ₂ P(AuL) ₃] ²⁺	—	—
[(L)B(AuL) ₄] ⁺	—	[RSe(AuL) ₂] ⁺	—



Auration Reactions




- 1 $L = PR_3$, $X = Cl$
- 2 $L = \text{carbene}$, $X = Cl$
- 3 $L = CN$, $X = Cl$
- 4 $L = PPh_2S$, $X = PPh_3$





Auration Reactions



Auration Reactions



Auration Reactions



Bond Energies of Auophilic Interactions

Compound/model	Au–Au contact mode	Method	Bond energy/kcal mol ⁻¹
(CO)AuCl	Unsupported intermolecular	MP2	10.0
(Et ₃ P)AuCl	Unsupported intermolecular	DFT	9.5
(NHC)AuCl	Unsupported intermolecular	DFT	8.6
[Au(CN) ₂] ⁻	Unsupported inter-anionic	EH/MP2	7.2
[Cl(AuPH ₃) ₂] ⁺	Unsupported inter-cationic	MP2/DFT	19.8
S(AuPH ₃) ₂	Unsupported intermolecular	MP2/DFT	29.8
[HS(AuPH ₃) ₂] ⁺	Unsupported inter-cationic	MP2/DFT	^a
[(Et ₃ PAu) ₂ C≡CB ₁₁ H ₁₁]	Unsupported inter-cationic	NMR	8.8
(dppe)Au ₂ [S ₂ C ₂ (CN) ₂]	Fully supported intramolecular	UV-vis	15.0
[(Xantphos) ₂ Au ₂] ²⁺	Fully supported intramolecular	NMR	11.6

^a Non-bonding due to Coulomb repulsion; attraction verified upon addition of anionic point charges.

H^+	R^+	LAu^+
H_2O	R_2O	$(\text{LAu})_2\text{O}$
H_3O^+	R_3O^+	$(\text{LAu})_3\text{O}^+$
H_4O^{2+}	R_4O^{2+}	$(\text{LAu})_4\text{O}^{2+}$
H_3N	R_3N	$(\text{LAu})_3\text{N}$
H_4N^+	R_4N^+	$(\text{LAu})_4\text{N}^+$
H_5N^{2+}	R_5N^{2+}	$(\text{LAu})_5\text{N}^{2+}$
H_4C	R_4C	$(\text{LAu})_4\text{C}$
H_5C^+	R_5C^+	$(\text{LAu})_5\text{C}^+$
H_6C^{2+}	R_6C^{2+}	$(\text{LAu})_6\text{C}^{2+}$

Hydrogen Bond




D–H bond elongates = weakened
D–H bond dipole increases
Dipole-dipole electrostatic energy increases
Attractive interaction increases

Charge transfer from A (lone pair, pi density)
to sigma* (antibonding) MO of D–H

D–H bond weakened = elongated

D–H vibration decreases = RED shift ($10 - 100 \text{ cm}^{-1}$)

D–H dipole increases = intensity in IR increases



Improper (Blue-Shifting) H-bond

P. Hobza et al.: JPC A 102, 2501 (1998)

benzene...H-X (X = CH₃, CCl₃, C₆H₅)



P. Hobza, Z. Havlas: Chem. Rev. 100, 4253 (2000)

Charge transfer from electron donor to distant parts
of electron acceptor (restructuralization of
acceptor electrons)

C–H strengthening, shortening = **BLUE SHIFT**
Rehybridization of C


Indirect mechanism

a two-step mechanism - a charge transfer from
the proton acceptor to a remote part of the proton
donor, followed by a structural reorganization of the
proton donor itself



Improper (Blue-Shifting) H-bond

$[\text{Cp}_2\text{Co}^+]$
salts in (solid state)




The blue spectra = an improper H-bond

- a: $[\text{Co}(\text{CN})_6]^{3-}$
- b: $[\text{PF}_6]^-$
- c: $[\text{Co}(\text{CO})_4]^-$

The red spectra = the proper H-bond

- d: $[\text{Br}_3]^-$
- e: $[\text{I}]^-$

a_{1g} = the in-phase displacement of all the C-H vibrators



Dihydrogen Bonds


The dihydrogen bond = an interaction between a transition metal or main-group **hydride** ($M-H$) and a **protic** hydrogen moiety ($H-X$)

intra- and intermolecular proton–hydride hydrogen bonds

Play a role in:

- crystal packing
- potential hydrogen-storage materials
- organometallic reaction mechanisms

X-ray crystal structure
of trans-[PtH(PhHN₃C₆H₆)(PPh₃)₂]BF₄



NMR Spectral Criteria of Dihydrogen Bonding

- shifts of the ^1H resonance of HX to lower field by 2–4 ppm
- high-field shift of the hydride (M–H) signal by 0.1–0.8 ppm and a 1.5–3-fold decrease of its longitudinal relaxation time ($T_{1\min}$)

rapid exchange on the NMR time scale of free and hydrogen-bonded molecules
weighted average between free and dihydrogen bonded hydrides


- an enhancement of H–H exchange coupling, $J_{\text{H–H}}$ on metal polyhydrides
- 1D nuclear Overhauser effect (NOE) spectroscopy or 2D ^1H nuclear Overhauser enhancement spectroscopy (NOESY)

estimation of H···H distances from spin-lattice T_1 relaxation measurements
short MH···HX contacts cause strong homonuclear dipolar coupling that provides
an additional contribution to nuclear dipole–dipole relaxation

$$r_{\text{H–H}} = 5.815 \left(\frac{T_{1\min}}{\nu} \right)^{\frac{1}{6}}$$

NMR Spectral Criteria of Dihydrogen Bonding

Hydride regions
500 MHz ^1H NMR



AB_2X spin system

$$J_{ab} = 80 \text{ Hz}$$

$$J_{ab} = 220 \text{ Hz}$$

one AB_2X multiplet is observed = an average over the fast exchanging 1a/1b

IR Spectroscopy Criteria of Dihydrogen Bonding

IR short time scale - detection of separate absorptions for free and hydrogen-bonded species.

The formation of a dihydrogen bond $\text{MH}\cdots\text{HX}$ = appearance of a new wider and more intense band, $\nu_{\text{XH}}^{\text{bonded}}$, of the proton donor

a band shift: $\Delta\nu_{\text{XH}} = \nu_{\text{XH}}^{\text{bonded}} - \nu_{\text{XH}}^{\text{free}}$ (red shift up to -450 cm^{-1})

elongation of the proton-donating HX bond

Enthalpies of intermolecular hydrogen bonds ΔH_{HB}
changes in the IR band positions ($\Delta\nu_{\text{XH}}$) and intensities (ΔA_{XH})
(in kcal mol^{-1})

$$\Delta H_{\text{HB}} = -18\Delta\nu_{\text{HX}} / (\Delta\nu_{\text{HX}} + 720)$$

$$\Delta H_{\text{HB}} = -0.30\Delta\nu_{\text{HX}}^{1/2}$$

$$\Delta H_{\text{HB}} = -2.9\Delta A_{\text{HX}}^{1/2} = -2.9(A_{\text{bonded}}^{1/2} - A_{\text{free}}^{1/2})$$


Enthalpies of Dihydrogen Bonds

Complexes of Boron Tetrahydride with Different Proton Donors

ROH	ν_σ in CH_2Cl_2 (cm^{-1})	$\Delta\nu_{\text{OH}}$ (cm^{-1})	$-\Delta H_{\text{exp}}$ in CH_2Cl_2 (kcal/mol)	$r_{\text{H}\cdots\text{H}}$ (\AA)
$\text{CFH}_2\text{CH}_2\text{OH}$	318	247	4.6	1.63
$\text{CF}_3\text{CH}_2\text{OH}$	324	290	5.2	1.55
$(\text{CF}_3)_2\text{CHOH}$	362	402	6.5	1.46



ν_σ = intermolecular $\text{H}\cdots\text{H}$ stretching mode

Enthalpies of Dihydrogen Bonds



Correlation between the enthalpy of DHB formation ($\Delta H^\circ_{\text{DHB}}$; derived from IR data) and the H···H distance (determined from NMR data on $T_{1\min}$) for complexes of fluorinated alcohols
[TFE = $\text{CF}_3\text{CH}_2\text{OH}$, HFIP = $(\text{CF}_3)_2\text{CHOH}$, PFTB = $(\text{CF}_3)_3\text{COH}$] in dichloromethane

Crystallographic Structural Data



Neutron diffraction crystal structure of $\text{ReH}_5(\text{PPh}_3)_3 \cdot \text{indole}$