

Ecotoxicology Current issues in Research vs Regulation

Ludek Blaha + ecotox colleagues









When Where

the assessment of toxicity is needed

What

to assess for toxicity









When & where the toxicity assessment is needed?

View of the researcher



Anytime!

... depending on researcher's budget

View of the regulator



As the law says!


... what are the law(s)?












Two approaches:

- Prospective (chemicals...)
- Retrospective (mixtures ...)






What to assess for toxicity?








	Current research topics	As required by law
Individual chemicals (prospective)		
Mixtures (prospective)		
Contaminated samples (retrospective)		
Resear for tox		

What to assess for toxicity?









		White consequent (MIII)
	Current research topics	As required by law
Individual chemicals (prospective)	Engineered nanomaterials/particles Ecological effects (e.g. of pharmaceuticals) Endocrine disruption & chronic diseases	Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics
Mixtures (prospective)		
Contaminated samples (retrospective)		
Research for toxic in the co		

Nanoparticles - examples



Toxicity of nanoparticles ...



(Mostly unknown)
Parameters may
Affect ecotoxicity

Composition (chemical)
Surface (size, area)
Charge
Reactivity
Interactions with ions,
other chemicals...


→ Effects on environmental Fate and toxicity





Ecotoxicity of nanoparticles – RECETOX example

Comparison of toxicity - 4 "appeared to be the same" particles (one producer – 4 different lots) (zerovalent iron – ZVI – Fe⁰)



?? Why is H16 so toxic ??

... despite of detailed investigation never revealed









PHARMACEUTICALS





Storage



Transport



Distribution

Storage



Transport



Consumption

Storage



Transport



Waste management


Manufacturing wast

Possible releases to the environment

Example 1 - DICLOFENAC

Unexpected effects at NON-TARGET species


- nephrotoxicity at vultures
- Relevant also in EU (ESP, EL,CY)



















Example 2 – AVERMEKTIN-like antiparasitics

Moxidectin – used e.g. in home "spot on" products














Ivermectin – antiparasitics in large herds

- Used **2-times per season** per sheep/cow
- Kills 100% parasites in sheep
- Released in dung kills 80-90% larvae of dung flies
- High concentrations in dung (released 2 days post application)
- Persistent in the soil (half-life 30 days)
- Can be washed into adjacent streams (highly toxic to water insects)



















POLICYFORUM


CONSERVATION


Economic Importance of Bats in Agriculture

Justin G. Boyles, 1* Paul M. Cryan, 2 Gary F. McCracken, 3 Thomas H. Kunz⁴



Insectivorous bat populations, adversely impacted by white-nose syndrome and wind turbines, may be worth billions of dollars to North American agriculture.









Biol. Lett. Published online

doi:10.1098/rsbl.2012.0685

Maternal predatorexposure has lifelong consequences for offspring learning in threespined sticklebacks



Daniel P. Roche, Katie E. McGhee* and Alison M. Bell

School of Integrative Biology, University of Illinois, Urbana, IL 61801, USA

*Author for correspondence (kemcghee@illinois.edu).

Stress

→ multigeneration effects





Epigenetics

→ DNA methylations




Table 1. Behaviours (mean \pm s.e.) of the offspring from the maternal treatments.

	offspring of predator-exposed mothers (s)	offspring of unexposed mothers (s)
initial exploratory behaviour (day 1: 09.00):		
latency to first begin moving	49 ± 30	56 ± 20
latency to enter either chamber for the first time	330 ± 70	326 ± 78
learning the colour association:		
day 1 (09.00): latency to find food reward	426 ± 65	427 ± 61
day 3 (09.00): latency to find food reward day 5 (09.00): latency to find food reward	$^{533 \pm 48}_{337 \pm 61}$ 2x difference	304 ± 74 158 ± 68

Testing comparability of existing and innovative bioassays for water quality assessment

Main questions:

Are current limits (for individual compounds) safe? Relevance of "Something from Nothing" phenomenon?

3 samples

- → 12 European laboratories different bioassays
- → ČR RECETOX: 11 bioassays



Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they?

Toxicol Sci 141(1): 218-233



Testing comparability of existing and innovative bioassays for water quality assessment

EU WFD priority substances

Different concentrations


EQS
= limit
(Environmental
Quality
Standard)

			l I
	RM 1ª	RM 2 ª	RM 3 a
Priority substances	around or >EQS	< EQS	< EQS
Atrazine	6	0.6	0.6
BaP	0.0017	0.00017	0.00017
Cadmium ^b	0.8	0.08	0.08
Chlorfenvinphos	1	0.1	0.1
Chlorpyrifos	0.3	0.03	0.03
DEHP (Bis(2-ethylhexyl) phthalate)	13	1.3	1.3
Diclofenac	1	0.1	0.1
diuron	2	0.2	0.2
17beta-estradiol	0.004	0.0004	0.0004
fluoranthene	0.063	0.0063	0.0063
Isoproturon	3	0.3	0.3
Ni ^b	40	4	4
4-Nonylphenol	3	0.3	0.3
Simazine	10	1	1
Carbamazepine	-	-	0.5
Sulfamethoxazole	-	-	0.6
Triclosan (Irgasan)	-	-	0.02
DEET	-	-	41
Bisphenol A	-	-	1.5



Testing comparability of existing and innovative bioassays for water quality assessment

Example: Effects of mixtures on *D. rerio* fish embryos



Control




Effects of RM 3 (i.e. safe) mixtures



Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they? Toxicol Sci 141(1):

Testing comparability of existing and innovative bioassays for water quality assessment

Example: Effects of mixtures on *X. laevis* frog embryos




Effects of RM 3 (i.e. safe) mixtures

Controls

Carvalho, R. et al. (2014) Mixtures of chemical pollutants at European legislation safety concentrations: how safe are they?

Toxicol Sci 141(1): 218-233






Biotest	Α	В	С
Microtox	26 and 36% stimulation of	18 and 35% stimulation of	22 and 39% stimulation of
	luminescence in 15 and 30 mins of	luminescence in 15 and 30 mins of	luminescence in 15 and 30 mins of
	exposure, respectively	exposure, respectively	exposure, respectively
Algae growth inhibition test 96-h	31% inhibition of growth compared	20% inhibition of growth compared	16% inhibition of growth compared
exposure	to solvent control	to solvent control	to solvent control
Acute immobilization test with	90% immobilization after 48 hours	no effect observed	no effect observed
D. magna	of exposure; 25% immobilization		
	occurred in 50% concentration - not		
	statistically significant		
Reproduction test with D.	100% mortality after 3 days of the	31 +/- 37 % inhibition of	23 +/- 24 % inhibition of
magna (21-d exposure)	test, no reproduction could be	reproduction, not statistically	reproduction, not statistically
	evaluated	significant	significant
FETAX (96-h exposure)	62 +/- 10 % of malformed embryos;	43 +/- 12 % of malformed embryos;	34 +/- 14 % of malformed embryos;
	no effect on embryo length	no effect on embryo length	no effect on embryo length
·	observed	observed	observed
FET (120-h exposure)	effects observed in number of	no significant effects observed	effects observed in number of
	defected embryos - absence of gas		defected embryos, number of
	bladder, (head) deformities and		underdeveloped embryos and
	underdeveloped embryos were		length
	observed the most often.		*
	*		*
In vitro - cytotoxicity	no effect observed compared to	no effect observed compared to	no effect observed compared to
	solvent control	solvent control	solvent control
In vitro - estrogenicity	effect under LOQ	effect under LOQ	effect under LOQ
In vitro - dioxin-like toxicity	effect under LOQ	effect under LOQ	effect under LOQ
In vitro - androgenicity	effect under LOQ	effect under LOQ	effect under LOQ
In vitro - antiandrogenicity	effect under LOQ	effect under LOQ	effect under LOQ

What to assess for toxicity?







		Visite Visite
	Current research topics	As required by law
Individual chemicals (prospective)	Engineered nanomaterials/particles Ecological effects (e.g. of pharmaceuticals) Endocrine disruption & chronic diseases	Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics
Mixtures (prospective)	Multistressors +T°C, salinity, pathogens, irradiation, food Exposome	LOADING
Contaminated samples (retrospective)		




What to assess for toxicity?



in the environment





	Current research topics	As required by law
Individual chemicals (prospective)	Engineered nanomaterials/particles Ecological effects (e.g. of pharmaceuticals) Endocrine disruption & chronic diseases	Industry & biocides (REACH) PPPs = pesticides Pharmaceuticals Cosmetics
Mixtures (prospective)	Multistressors +T°C, salinity, pathogens, irradiation, food Exposome	LOADING
Contaminated samples (retrospective)	Can analyzed chemicals	Chemical analyses & limits (see lectures: RISK ASSESSMENT part)
(rotroopootivo)	explain observed effects?	Effect testing rare: Remediation, dredged sediments (CZ), effluents
		LOADING
for toxi	ch centre c compounds	TECHNICAL REPORT ON AQUATIC EFFECT-BASED MONITORING TOOLS

Contaminated samples? Case study "air"

Active sampling particles vs gaseous phase

 Reference locality – agriculture (Košetice observatory)


 Region A – industrial (historically OCPs production)


 Region B – combined: industry, agriculture, traffic

Novák et al. (2009) Environment International










Dioxin-like effects



- Difference B>A
- Difference B vs A particles vs gas











Antiandrogenic effects



Quantitative – comparable

Clear differences in patterns ... no effects on particles in "B







Summary on When, Where, What

- Regulatory world
 - Assessment of "chemicals"!
- Contaminated samples
 - effects rarely tested
 - Great value of bioassays

 in assessment of contaminated samples
 - Effects observed (!)
 - How to set the "limits"?
- Research issues and questions
 - Nanomaterials, Pharmaceuticals, EDCs
 - Mixtures!
 - Exposome





Contents lists available at ScienceDirect

Environment International

journal homepage: www.elsevier.com/locate/envint

Review

What level of estrogenic activity determined by *in vitro* assays in municipal waste waters can be considered as safe?

Barbora Jarošová ^a, Luděk Bláha ^a, John P. Giesy ^b, Klára Hilscherová ^{a,*}

- ^a Masaryk University, Faculty of Science, RECETOX, Kamenice 5, CZ-62500 Brno, Czech Republic
- b Department of Biomedical Veterinary Sciences and Toxicology Centre, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

