




# Hertzprung-Russell Diagram



# Hertzprung-Russell Diagram





# Color-Magnitude Diagram



Gaia Collaboration,  
2018, A&A, 616,  
A10


**Fig. 5.** *Gaia* HRD of sources with **low extinction** ( $E(B - V) < 0.015$  mag) satisfying the filters described in Sect. 2.1 (4,276,690 stars).

# Color-Magnitude Diagram




# Color-Magnitude Diagram

## Pleiades




# Color-Magnitude Diagram

## Praesepe




# Colour and $T_{\text{eff}}$


- Measuring accurate  $T_{\text{eff}}$  for stars is an intensive task – spectra needed and model atmospheres
- Spectral Energy Distribution (SED) fitting, only useful if measurements in the UV are available
- Magnitudes of stars are measured at different wavelengths
- Colours  $\Rightarrow$  Calibrations  $\Rightarrow T_{\text{eff}}$
- The Asiago Database on Photometric Systems (ADPS) lists about 200 different systems



a sequence of stellar flux profiles



# Colour and $T_{\text{eff}}$



Various calibrations can be used to provide the colour relation:

$$(B - V) = f(T_{\text{eff}})$$


Remember that observed  $(B - V)$  must be corrected for interstellar extinction to

$$(B - V)_0$$

Most of the calibrations are for cool type stars

# Absorption = Extinction = Reddening

- $A_V = k_1 E(B-V) = k_2 E(V-R) = \dots$
- *General extinction* because of the ISM characteristics between the observer and the object
- *Differential extinction* within one star cluster because of local environment
- Both types are, in general *wavelength dependent*



Important parameter:

$$R_V = A_V / E(B - V)$$

Normalization factor

Standard value used is 3.1

Be careful, different values used!

Depending on the line of sight

TABLE 2  
OPTICAL/IR EXTINCTION RATIOS FOR  $R = 3.1$

Extinction Ratio (1)	Observed Value (2)	References (3)	Model Curve Value (4)
$A(M)/E(B-V)$ .....	0.08–0.12	1, 2	0.12
$A(L)/E(B-V)$ .....	0.09–0.20	1,2,3,4	0.19
$A(K)/E(B-V)$ .....	0.33–0.38	2, 3, 4	0.36
$A(H)/E(B-V)$ .....	0.52–0.55	1, 2	0.53
$A(J)/E(B-V)$ .....	0.85–0.91	1, 2, 3	0.86
$A(I)/E(B-V)$ .....	1.50	3	1.57
$A(R)/E(B-V)$ .....	2.32	3	2.32
$A(V)/E(B-V)$ .....	3.10		3.10
$E(U-B)/E(B-V)$ .....	$0.70 + 0.05 \times E(B - V)$	5	$0.69 + 0.04 \times E(B - V)$
$E(b-y)/E(B-V)$ .....	0.74	6	0.74
$E(m1)/E(b-y)$ .....	-0.32	6	-0.32
$E(c1)/E(b-y)$ .....	0.20	6	0.17
$E(u-b)/E(b-y)$ .....	1.5	6	1.54

REFERENCES.—(1) Rieke & Lebofsky 1985; (2) Whittet 1988; (3) Schultz & Wiemer 1975; (4) Savage & Mathis 1979; (5) FitzGerald 1970; (6) Crawford 1975.

**Table 3.** Multiband Relative Extinction Values

Band ( $\lambda$ )	$\lambda_{\text{eff},0}$ ( $\mu\text{m}$ )	$A_\lambda/A_{G_{\text{RP}}}$	$A_\lambda/A_{G_{\text{RP}}}$ (from Chen18)	$A_\lambda/A_V$	$A_\lambda/E(G_{\text{BP}} - G_{\text{RP}})$
<i>GAIA</i> $G_{\text{BP}}$	0.5387	$1.700 \pm 0.007$		$1.002 \pm 0.007$	$2.429 \pm 0.015$
<i>GAIA</i> $G_{\text{RP}}$	0.7667	1		$0.589 \pm 0.004$	$1.429 \pm 0.015$
Johnson $B$	0.4525	$2.206 \pm 0.023$		$1.317 \pm 0.016$	$3.151 \pm 0.027$
Johnson $V$	0.5525	$1.675 \pm 0.010$		1	$2.394 \pm 0.018$
SDSS $u$	0.3602	$2.653 \pm 0.024$		$1.584 \pm 0.017$	$3.791 \pm 0.028$
SDSS $g$	0.4784	$2.018 \pm 0.012$		$1.205 \pm 0.010$	$2.883 \pm 0.019$
SDSS $r$	0.6166	$1.421 \pm 0.006$		$0.848 \pm 0.006$	$2.030 \pm 0.016$
SDSS $i$	0.7483	$1.056 \pm 0.002$		$0.630 \pm 0.004$	$1.509 \pm 0.015$
SDSS $z$	0.8915	$0.767 \pm 0.004$		$0.458 \pm 0.003$	$1.096 \pm 0.012$
Pan-STARRS $g$	0.4957	$1.934 \pm 0.010$		$1.155 \pm 0.009$	$2.764 \pm 0.018$
Pan-STARRS $r$	0.6211	$1.413 \pm 0.005$		$0.843 \pm 0.006$	$2.019 \pm 0.015$
Pan-STARRS $i$	0.7522	$1.052 \pm 0.001$		$0.628 \pm 0.004$	$1.503 \pm 0.015$
Pan-STARRS $z$	0.8671	$0.815 \pm 0.002$		$0.487 \pm 0.003$	$1.165 \pm 0.012$
Pan-STARRS $y$	0.9707	$0.662 \pm 0.004$		$0.395 \pm 0.003$	$0.947 \pm 0.011$
2MASS $J$	1.2345	$0.407 \pm 0.007$		$0.243 \pm 0.004$	$0.582 \pm 0.011$
2MASS $H$	1.6393	$0.219 \pm 0.010$	$0.222 \pm 0.012$	$0.131 \pm 0.006$	$0.313 \pm 0.014$
2MASS $K_S$	2.1757	$0.125 \pm 0.010$	$0.130 \pm 0.006$	$0.078 \pm 0.004$	$0.186 \pm 0.009$
<i>WISE</i> $W1$	3.3172	$0.055 \pm 0.011$	$0.066 \pm 0.006$	$0.039 \pm 0.004$	$0.094 \pm 0.009$
<i>WISE</i> $W2$	4.5501	$0.029 \pm 0.011$	$0.044 \pm 0.006$	$0.026 \pm 0.004$	$0.063 \pm 0.009$
<i>WISE</i> $W3$	11.7281	$0.066 \pm 0.016$		$0.040 \pm 0.009$	$0.095 \pm 0.021$
<i>GAIA</i> $G$	0.6419	$1.323 \pm 0.003$		$0.789 \pm 0.005$	$1.890 \pm 0.015$
<i>Spitzer</i> [3.6]			$0.062 \pm 0.005$	$0.037 \pm 0.003$	$0.089 \pm 0.007$
<i>Spitzer</i> [4.5]			$0.044 \pm 0.005$	$0.026 \pm 0.003$	$0.063 \pm 0.007$
<i>Spitzer</i> [5.8]			$0.031 \pm 0.005$	$0.019 \pm 0.003$	$0.044 \pm 0.007$
<i>Spitzer</i> [8.0]			$0.042 \pm 0.005$	$0.025 \pm 0.003$	$0.060 \pm 0.007$

Wang & Chen, 2019,  
ApJ, 877, 116

At *Spitzer* bands, the determination of the relative extinction  $A_\lambda/A_V$  and the extinction coefficient  $A_\lambda/E(G_{\text{BP}} - G_{\text{RP}})$  are based on the relative extinction values from Chen18.

# Absolute magnitude and bolometric magnitude

- **Absolute Magnitude**  $M$  defined as apparent magnitude of a star if it were placed at a distance of 10 pc

$$(V - M_V) - A_V = 5 \log(d) - 5$$

where  $d$  is in pc.  $(V - M_V)$  is also called **distance modulus**.


- Magnitudes are measured in some wavelength. To compare with theory, it is more useful to determine **bolometric magnitude**  $M_{\text{bol}}$  – defined as absolute magnitude that would be measured by a bolometer sensitive to all wavelengths. We define the bolometric correction to be

$$\text{BC} = M_{\text{bol}} - M_V$$

Bolometric luminosity is then

$$M_{\text{bol}} - M_{\text{bol},\odot} = -2.5 \log L/L_{\odot}; M_{\text{bol},\odot} = 4.75 \text{ mag}$$

# Bolometric Correction



BC from Flower, 1996, ApJ, 469, 355