

## Hyades

$\log t = 8.90$


$d = 45$  pc

$[Fe/H] = +0.17$  dex

Width of Main Sequence  
about 1.8 mag in  $M_V$

NO  
Observational error

What are the reasons?



Vertical distance from the main sequence

$$x = a(C_{AB} - C_A) + V_A - V_{AB}$$

Absolute magnitude:

$$M_V = -2.5 \log (L_1 + L_2)$$

Maximum at  $L_1 = L_2 \Rightarrow$

$$M_V = -0.753 \text{ mag}$$


The maximal width of the main sequence due to binary systems is 0.753 mag


# Praesepe: Fossati et al., 2008, A&A, 483, 891


At.N.	Element	"Normal" A-type stars						Solar Abundances	
		HD 72846	HD 73345	HD 73450	HD 73574	HD 74028	HD 74050		
3	Li	< -8.08(−; 1)	< -8.33(−; 1)	< -8.70(−; 1)	< -8.38(−; 1)		< -8.41(−; 1)	< -8.26(−; 1)	-10.99
6	C	-3.58(−; 1)	-3.44(12; 3)	-3.27(−; 1)	-3.36(18; 2)	-3.39(08; 2)	-3.52(−; 1)	-3.49(01; 2)	-3.51(04; 2)
8	O	-3.18(−; 1)	-3.22(01; 2)				-3.70(−; 1)	-3.30(−; 1)	-3.38
11	Na	-5.44(01; 2)	-5.37(01; 2)	-6.28(−; 1)	-5.57(02; 2)	-5.98(−; 1)	-5.64(13; 2)	-5.61(02; 2)	-5.70(14; 2)
12	Mg	-4.18(08; 3)	-4.18(02; 3)	-5.02(18; 2)	-4.37(04; 3)	-4.86(08; 3)	-4.22(05; 4)	-4.56(08; 3)	-4.52(01; 2)
14	Si	-4.62(16; 2)	-4.67(−; 1)	-4.13(−; 1)	-4.19(−; 1)	-4.17(−; 1)	-4.37(−; 1)	-4.16(−; 1)	-4.25(−; 1)
16	S	-4.71(04; 2)	-4.44(03; 4)	-4.35(−; 1)	-4.61(02; 2)	-4.26(01; 2)		-4.50(04; 2)	-4.28(11; 2)
20	Ca	-5.17(−; 1)	-5.39(09; 6)	-5.95(06; 4)	-5.86(16; 5)	-5.37(16; 2)	-6.13(06; 2)	-5.49(15; 6)	-5.68(02; 3)
21	Sc	-8.88(−; 1)	-8.63(07; 3)	-8.57(14; 3)	-8.89(02; 3)	-8.35(−; 1)	-8.96(27; 3)	-8.56(−; 1)	-8.69(14; 2)
22	Ti	-6.88(03; 5)	-6.95(06; 6)	-7.30(11; 5)	-6.98(09; 5)	-6.78(−; 1)	-7.08(15; 5)	-6.83(16; 3)	-6.93(10; 5)
24	Cr	-6.23(06; 3)	-6.22(08; 2)	-6.56(08; 3)	-6.19(16; 3)	-6.23(12; 4)	-6.48(10; 3)	-6.05(13; 4)	-6.44(20; 5)
25	Mn	-6.37(−; 1)	-6.88(−; 1)	-6.52(02; 2)	-6.77(−; 1)	-6.61(−; 1)	-6.62(04; 2)	-6.71(−; 1)	-6.65
26	Fe	-4.55(18; 42)	-4.33(11; 61)	-4.62(09; 15)	-4.49(10; 30)	-4.50(09; 18)	-4.44(13; 16)	-4.28(10; 33)	-4.61(11; 26)
28	Ni	-5.70(18; 2)	-5.58(11; 4)	-5.82(16; 2)	-5.62(08; 4)	-5.93(14; 3)	-5.60(15; 3)	-5.84(−; 1)	-5.68(02; 3)
39	Y	-9.75(−; 1)	-9.46(−; 1)	-9.83(−; 1)	-9.20(−; 1)	-9.56(−; 1)	-9.26(−; 1)	-9.13(−; 1)	-9.10(−; 1)
56	Ba	-9.48(−; 1)	-9.30(06; 2)	-9.50(02; 2)	-8.98(04; 2)	-9.65(−; 1)	-9.52(01; 2)	-8.96(25; 2)	-9.15(−; 1)
		8045	7993	7270	7662	7750	7872	7500	7600
		log <i>g</i>	3.50	3.96	4.20	4.00	4.50	3.66	4.20
		<i>v</i> <sub>mic</sub>	2.5	2.6	2.7	2.6	2.6	2.7	2.7
		<i>v sin i</i>	119	85	138	102	150	188	90
									155


Fe: -4.28 to -4.62dex; 0.34 dex

# Metallicity => different opacity









Von Zeipel theorem (1924,  
MNRAS, 84, 665)

Energy generation rate

$$\epsilon = (\text{const}) \left( 1 - \frac{\omega^2}{2\pi G \rho} \right)$$



From the rotational velocity  $\Rightarrow \epsilon \Rightarrow T_{\text{eff}}$  and  $L$  ( $\log g$ )

# Vega



**Table 7**

A Summary of Physical Parameters for the Pole-on Model

Parameter	Value
Equatorial radius ( $R_{\odot}$ )	$2.75 \pm 0.01$
Polar radius ( $R_{\odot}$ )	$2.40 \pm 0.02$
Polar effective temperature (K)	$10000 \pm 30$
Pole-to-equator $T_{\text{eff}}$ difference (K)	1410
Mean effective temperature (K)	$9560 \pm 30$
Luminosity ( $L_{\odot}$ )	$44 \pm 2$
Mass ( $M_{\odot}$ )	$2.4 \pm 0.1$
Polar surface gravity (cgs, dex)	$4.04 \pm 0.01$
Pole-to-equator log $g$ difference (cgs, dex)	0.26
Mean surface gravity (cgs, dex)	$3.95 \pm 0.01$
Projected rotational velocity (km s $^{-1}$ )	$20.8 \pm 0.2$
Inclination of rotation axis (degrees)	$5.7 \pm 0.1$
Equatorial rotational velocity (km s $^{-1}$ )	$211 \pm 4$
Fraction of breakup velocity (km s $^{-1}$ )	$0.81 \pm .02$



$p$  ... Degree of differential rotation




# Conclusions – Width of the Main Sequence

- Differential reddening:  $k \cdot \Delta E(B-V)$
- Spectroscopic Binaries: 0.753 mag
- Metallicity: up to 1.2 mag for  $M_V$ , but only 0.2 mag for  $(U - B)$  versus  $(B - V)$
- Rotation: 1 mag for  $M_V$ , 0.2 (?) mag for  $(U - B)$  versus  $(B - V)$

# Binary fraction

- Important for the formation and evolution of star clusters
- Critical parameter for the IMF
- Needed for N-body numerical simulations
- Observations are biased in many respects
- Many different types of binary systems

## Metallicity




# Rotational Energy


Lower metallicities seem to favour binary formation

# How to observe the binary fraction?

- Photometric observations of star clusters
  1. “Cluster main sequence”
  2. Eclipsing binaries
  3. Positions (astrometric binaries)
- Spectroscopic observations
  1. Radial velocity variability
  2. Direct detection in spectrum (SB2)




Simulation with randomly distributed mass ratios



Observations of Praesepe with known binary systems

# Observed binaries with the Gaia DR2



**Fig. A.11.** Left: distribution of the probable members of NGC 752. Right: colour-magnitude diagram of the probable members.

# Observed binaries with the Gaia DR2





Fig. A.8: Member stars (blue) together with field stars (grey) for UBC8 in  $(l, b)$  (left) and in proper motion space (middle). The Color-Magnitude Diagram shows the sequence of the identified members (outlining an empirical isochrone) (right).


Ninth Catalogue of Spectroscopic Binary Orbits  
<http://sb9.astro.ulb.ac.be/>


Bin size of one day



Ninth Catalogue of Spectroscopic Binary Orbits  
<http://sb9.astro.ulb.ac.be/>

Bin size of one hundred days





# Results for open clusters

- Sollima et al., 2010, MNRAS, 401, 577
  - NGC 188 (9.63): 21 – 58%
  - NGC 2204 (9.20): 12 – 36%
  - NGC 2243 (9.58): 34 – 70%
  - NGC 2420 (9.08): 17 – 51%
  - NGC 2516 (8.52): 25 – 66%
- Sana et al., 2009, MNRAS, 400, 1479
  - NGC 6611 (6.50): 44 – 67%
- Sana et al., 2008, MNRAS, 386, 447
  - NGC 6231 (6.50): 63% - ?
- Bica & Bonatto, 2005, A&A, 431, 943
  - IC 4651 (9.26): 50 +- 11%
  - NGC 2287 (8.20): 48 +- 45%
  - NGC 2447 (8.60): 21 +- 9%
  - NGC 2548 (8.56): 48 +- 23%
  - NGC 2682 (9.51): 39 +- 16%
  - NGC 3680 (9.20): 25 +- 5%
  - NGC 5822 (9.00): 16 +- 8%
  - NGC 6208 (9.11): 54 +- 30%
  - NGC 6694 (7.85): 18 +- 12%
- Sandhu et al., 2003, A&A, 408, 515
  - NGC 2099 (8.60): ~30%
  - King 5 (9.00): ~30%
  - King 7 (8.80): ~20%

	<i>f</i>		Globular clusters
NGC 288	$0.15 \pm 0.05$ $>0.06$	M3	
NGC 362	$0.21 \pm 0.06$		$0.14 \pm 0.08$
NGC 2808		M4	$0.23^{+0.34}_{-0.23}$
NGC 3201		M15	$\sim 0.07$
NGC 4590	$>0.09$	M22	
NGC 5053	$>0.08$	M30	
NGC 5466	$>0.08$	M55	$>0.06$
NGC 5897	$>0.07$	M71	$0.22^{+0.26}_{-0.12}$
NGC 6101	$>0.09$	M92	
NGC 6362	$>0.06$	Arp 2	$>0.08$
NGC 6397	$<0.07$	Terzan 7	$>0.21$
NGC 6723	$>0.06$	Palmoar 12	$>0.18$
NGC 6752	$0.27 \pm 0.12$	Palmoar 13	$0.30 \pm 0.04$
NGC 6792		47 Tucane	$0.14 \pm 0.04$
NGC 6981	$>0.10$		