



Regulation of cell function by FGFR3 receptor tyrosine kinase

Pavel Krejci


Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, CA
Institute of Experimental Biology, Masaryk University, Brno, Czech Republic



FGFR3-related skeletal dysplasias

Thanatophoric Dysplasia

Achondroplasia






- short long bones
- brachydactyly
- macrocephaly
- low nasal bridge
- spinal stenosis
- temporal lobe malformations

FGFR3-related skeletal dysplasia

STATURE

Hypochondroplasia
Achondroplasia
SADDAN
Thanatophoric Dysplasia





Frequent activating mutations of FGFR3 in human bladder and cervix carcinomas

epithelia (data not shown), we examined the expression and mutational status of *FGFR3* in a series of bladder and cervix carcinomas to determine whether *FGFR3* is involved in epithelial tumorigenesis.

We assessed transcript levels of the two *FGFR3* variants¹⁴, *FGFR3b* and *FGFR3c*,

Frequent translocation t(4;14)(p16.3;q32.3) in multiple myeloma is associated with increased expression and activating mutations of fibroblast growth factor receptor 3

Marta Chesi¹, Elena Nardini², Leslie A. Brents¹, Evelin Schröck³, Thomas Ried³, W. Michael Kuehl¹ & P. Leif Bergsagel²

Activating mutations in *FGFR3* and *HRAS* reveal a shared genetic origin for congenital disorders and testicular tumors

Anne Goriely¹, Ruth M S Hansen¹, Indira B Taylor¹, Inge A Olesen², Grete Krag Jacobsen³, Simon J McGowan⁴, Susanne P Pfeifer⁵, Gilean A T McVean⁵, Ewa Rajpert-De Meyts² & Andrew O M Wilkie¹

Skeleton: hypochondroplasia, achondroplasia, thanatophoric dysplasia, SADDAN, Muenke syndrome

Skin: epidermal nevi, seborrheic keratosis, acanthosis nigricans

Cancer: multiple myeloma, bladder cancer, seminoma

Investors



[Home](#) [Investors](#) [RSS Content](#)

[Overview](#)

[Financial Information](#)

[SEC Filings](#)

[Corporate Governance](#)

[Investors' Kit](#)

[Calendar of Events](#)

RSS Content

BioMarin Announces Program for BMN-111 for the Treatment of Achondroplasia

NOVATO, Calif., Oct 19, 2010 /PRNewswire via COMTEX/ --

BioMarin Pharmaceutical Inc. (Nasdaq: BMRN) today announced its program for BMN-111, a peptide therapeutic for the treatment of achondroplasia. BioMarin plans to file an IND in the fourth quarter of 2011 and to initiate a Phase 1 clinical trial by the first quarter of 2012.

BMN-111, for the treatment of achondroplasia, is an analog of C-type Natriuretic Peptide (CNP), a small cyclic peptide that is a positive regulator of bone growth. It is produced and has a receptor in the growth plate, and along with the fibroblast growth factor receptor 3 (FGFR3), regulates normal bone growth. In addition to short stature, there are complications in achondroplasia that are related to bone compression (e.g. foramen magnum narrowing, spinal stenosis, upper respiratory narrowing) of nervous tissues or other tissues.

Exp Cell Res. 2004;297:152-64.

J Cell Sci. 2005; 118: 5089-100.

J Biol Chem. 2007 ;282:2929-36.

Pediatr Res. 2007; 61(3):267-72.

Invest New Drugs 2007; 25:391-95.

PLoS One 2008; 3:e3961.

J Cell Sci 2008; 121:272-81.

Cell Signal 2009; 21:151-60.

Hum Mol Genet. 2009; 18:227-40.

J Biol Chem 2010; 285:20644-53.








Bone 2010; 47:102-10.

Leukemia 2011; 25:538-50.



Human Mutation 2012; 33:29-41

healthy

TD




FGFR3 inhibits chondrocyte proliferation by arresting their cell cycle in G1 phase





FGFR3 alters the cartilage-like phenotype of chondrocytes

control



FGF2



FGF2 (h)

M C 0.5 1 4 8 12 24 48 72

aggre can


collagen II

collagen I

GAPDH

control


FGF2




proMMP9
matureMMP9



proMMP2
intermediate
matureMMP2




pro MMP3/10
mature MMP3/10




proMMP13
matureMMP13


Vinculin




Phalloidin







DAPI





merge






FGF2






FGFR3 causes premature senescence in chondrocytes






FGF2 (hours) — 8 12 24 48 72 —



FGFR3 arrests chondrocyte growth via RAS-ERK MAP kinase pathway










Activation of Stat1 by mutant fibroblast growth-factor receptor in thanatophoric dysplasia type II dwarfism


**Wu-Chou S. Su*, Motoo Kitagawa*, Ninrong Xue,
Bing Xie, Silvio Garofalo†, Jay Cho†, Chuxia Deng‡,
William A. Horton† & Xin-Yuan Fu**




NATURE | VOL 386 | 20 MARCH 1997



FGFR3 associates with STAT1 and acts as STAT1-kinase in chondrocytes










FGFR3 mutation	cases, n = 591 (% total)
N540K	70 (16.9%)
G380R	308 (52.1%)
R248C	102 (17.3%)
Y373C	40 (6.8%)
K650M	5 (0.8%)
K650E	24 (4%)
other	42 (7.1%)
	520 (88%)
	29 (4.9%)


Chronic FGF stimulus inhibits cytokine/STAT signaling in chondrocytes






2001







2012














C-natriuretic peptide (CNP)



wild-type


CNP↑



Kazuwa Nakao







Nature Medicine 10, 80–86 (2004).







control


FGF2

control



CNP





Simple, mammalian cell-based assay for identification of inhibitors of the Erk MAP kinase pathway

Pavel Krejci · Katerina Pejchalova · William R. Wilcox

Received: 26 December 2006 / Accepted: 4 April 2007 / Published online: 26 April 2007
© Springer Science + Business Media, LLC 2007

Summary The Erk MAP kinase pathway contributes to tumor development and thus represents an important therapeutic target. Several inhibitors of the Erk pathway are presently being evaluated in clinical trials for cancer, but show limited efficiency thus warranting discovery of more

toxic compounds as false-positive hits, given the nature of the RCS response to inhibition of the Erk pathway, i.e. growth.





Keywords Erk · Inhibitor · FGFR3 · Growth arrest ·




Tocriscreen™ *Mini Library*

A collection of 1120 biologically active compounds supplied as DMSO solutions.

NF449



Kinase: FGFR3



Cedars-Sinai Medical Center
Los Angeles, California

Betty Mekikian

Anie Aklian

UCI, Irvine, California

Leslie Thompson

Tamara Kashiwada

Lisa Salazar

UCLA, Los Angeles, California

Matthew Schibler

Masaryk University, Brno, Czech Republic

Jirina Prochazkova

Vita Bryja

Lukas Balek

Tereza Spoustova

Tereza Pospisilova

INSERM U589, Toulouse, France

Bernard Masri

Vincent Fontaine