

Lecture 10:

Wnt morphogenetic system


Wnts (Wingless/Int)

- family of ligands
- 19 members in human and mouse
- glycosylated and palmitoylated extracellular proteins
- short range of action, bind to extracellular matrix
- only in multicellular animals



canonical

(eg. Wnt-1 or Wnt-3a)



non-canonical


(eg. Wnt-5a)

Wnt/ β -kateninová dráha (= kanonická dráha)



- induce axis duplication in *Xenopus*
- induce transformation of mammary cell line C57mg
- signal via nuclear translocation of β -catenin

Wnt/ β -catenin pathway



The Canonical Wnt signalling cascade. Canonical Wnt signalling mediates its effect by binding to their receptors frizzled (Fzd) and co-receptors, LRP 5/6. This causes activation of intracellular Dishevelled (Dvl) which, in turn, inhibits glycogen synthase kinase-3 (GSK3 β). This results in the stabilisation and nuclear translocation of β -catenin, inducing gene transcription via the LEF/TCF family of transcription factors. In the absence of Wnt signalling, a complex containing GSK3 β phosphorylates β -catenin, leading to degradation by ubiquitination. Copyright BTR ©

Wnt secretion





Figure 1. Wnt Secretion



To be secreted, Wnt proteins in the endoplasmic reticulum (ER) need to be palmitoylated by the action of Porcupine. Wnt proteins also require Wntless (Wis/Evi) in order to be routed to the outside of the cell. Loading onto lipoprotein particles may occur in a dedicated endo/exocytic compartment. The retromer complex may shuttle Wis between the Golgi and the endo/exocytic compartment.

Wnt and E-cadherin pathways



Tyr142 of β -catenin:
 phosphorylated – binds Bcl9
 dephosphorylated – binds α -catenin

Epithelio-mesenchymal transition (EMT)



Příklady vývojových procesů regulovaných kanonickou Wnt dráhou

Maternální Wnt/ β -cateninová dráha
determinuje dorsální (horní) pól
vyvíjející se zygoty a embrya




Fig. 2.6 Gastrulation in amphibians. The blastula (first panel) contains several thousand cells and there is a fluid-filled cavity, the blastocoel, beneath the cells at the animal pole. Gastrulation begins (second panel) at the blastopore, which forms on the dorsal side of the embryo. Future mesoderm and endoderm of the marginal zone move inside at this site through the dorsal lip of the b

between the endoderm and ectoderm in the animal region (third panel). The tissue movements create a new internal cavity—the archenteron—which will become the gut. Endoderm in the ventral region also moves inside through the ventral lip of the blastopore (fourth panel) and will eventually completely line the archenteron. At the end of gastrulation the blastocoel has

A

Ear
e
Lat




a

ray
ge 9)
ray
(8)

Wnt/ β -cateninová dráha určuje antero-posteriorní (AP, předozadní) osu těla během gastrulace – podporuje vznik zadních a blokuje vznik předních částí těla

A

Early *Xenopus*
embryo
Lateral view



Xenopus blastula




myší embryo po gastrulaci (E8.5):

Cílové geny Wnt/ β -cateninové dráhy jsou exprimovány v zadní části těla.


Uncx4.1/Mesogenin

Wnt5a^{+/+};LRP6^{+/+}



Wnt5a^{-/-};LRP6^{+/-}


Deplece Wnt/ β -kateninové dráhy při gastrulaci = ztráta zadních částí těla





wild type

Wnt-3a knockout

Deplece inhibitorů Wnt/ β - kateninové dráhy při gastrulaci = ztráta předních částí těla



wild type vs. Dkk1 knockout




β-Catenin Defines Head Versus Tail Identity During Planarian Regeneration and Homeostasis

Kyle A. Gurley, Jochen C. Rink, Alejandro Sánchez Alvarado*

SCIENCE VOL 319 18 JANUARY 2008

Wnt/ β -cateninová dráha je
klíčovým regulátorem aktivace
kmenových buněk jak v
embryogenezi, tak v dospělých
tkáních



Wnt/ β -catenin dráha je velmi často deregulovaná u nádorů!



according to Beachy et al., Nature 2004

Wnt pathway


Colon	Adenocarcinoma	Tumorigenesis by inactivation of APC, Axin; tumorigenesis by stabilization of β -catenin; epigenetic inactivation of SFRPs
Liver	Hepatoblastoma	Tumorigenesis (in mouse) by inactivation of APC and by stabilization of β -catenin
Blood	Multiple myeloma	Cell-growth inhibition by dominant negative TCF4; growth stimulation by Wnt ligand
Hair follicle	Pilomatricoma	Tumorigenesis (in mouse) by overexpression of β -catenin
Bone	Osteosarcoma	Dkk3 and LRP5 expression inhibits tumour cell growth <i>in vitro</i>
Lung	Non-small-cell carcinoma	Apoptosis and cell-growth inhibition by short interfering RNA and a blocking antibody against Wnt2
Pleura	Mesothelioma	Apoptosis and cell-growth inhibition by transfection of SFRP




Reya & Clevers 2005, Nature

The effects of Wnts on stem cells in their niche



Under Normal Physiological Conditions



In Cancers or Tumors



Wnt pathway induces de novo formation of hair follicles



Wnt signaling pathway related polymorphism?



Wnt-dependent *de novo* hair follicle regeneration in adult mouse skin after wounding

Mayumi Ito¹, Zaixin Yang¹, Thomas Andl¹, Chunhua Cui¹, Noori Kim¹, Sarah E. Millar¹ & George Cotsarelis¹




nature

Vol 447 | 17 May 2007 | doi:10.1038/nature05766

LETTERS




Aktivace β -catenininu ve vyvíjející se mozkové trubici:

midbrain (Brn4-promotor)




Zechner et al., 2003: Dev. Biol.;258:406-418.

cortex (nestin enhancer)



Chenn & Walsh, 2002: Science;297:365-369.


Aktivace beta-cateninu v kmenových buňkách zubu:



Wnt/ β -cateninová dráha reguluje
vývoj neurální lišty (neural crest)

Nervová lišta (neural crest)

- populace buněk, vzniká z dorsální (= horní) strany nervové trubice procesem delaminace a migruje několika hlavními cestami do jiných částí embrya




Co všechno z NC vzniká?

- podíl na vzniku cca 40 různých tkání a orgánů


1. Oblast trupu:

- Neurony a gliální buňky sensorického, sympatického a parasympatického systému
- Buňky dřeně nadledvin
- Pigmentové buňky epidermis
- Svalové buňky některých cév





I. Vývoj neurální lišty:



Wnt-3a




- neurální lišta je zdrojem periferního nervového systému, melanocytů, obličejových kostí a svalů, částí srdce a dalších



Wnt signaling při vývoji neurální lišty



Wnt1/3a DKO



Wnt/ β -cateninová dráha v regulaci stárnutí

Augmented Wnt Signaling in a Mammalian Model of Accelerated Aging

Hongjun Liu,¹ Maria M Fergusson,^{1*} Rogerio M. Castilho,^{2*} Jie Liu,¹ Liu Cao,¹ Jichun Chen,³ Daniela Malide,⁴ Ilsa I. Rovira,¹ Daniel Schimel,⁵ Calvin J. Kuo,⁶ J. Silvio Gutkind,² Paul M. Hwang,¹ Toren Finkel^{1†}

SCIENCE VOL 317 10 AUGUST 2007

803

Increased Wnt Signaling During Aging Alters Muscle Stem Cell Fate and Increases Fibrosis

Andrew S. Brack,¹ Michael J. Conboy,¹ Sudeep Roy,¹ Mark Lee,² Calvin J. Kuo,² Charles Keller,³ Thomas A. Rando^{1,4*}



SCIENCE VOL 317 10 AUGUST 2007

807

Klotho myš

-mutantní kmen myši s fenotypem akcelerovaného stárnutí: např. kratší život, arterioskleróza, snížená plodnost nebo kožní atrofie


- protein Klotho je transmembránový protein s velkou extracelulární doménou, ta může být odštěpena a volně cirkulovat v krvi



WT

Klo

aktivita Wnt/ β -cateninové dráhy
ve střevním epitelu









Model 1 – svalová fibróza

- s prodlužujícím se věkem stále častěji při regeneraci svalu vznikají místo svalových buněk buňky fibrózní tkáně – tak přispívají k nižší výkonnosti svalu, která souvisí se stárnutím

Model 2 – parabiotické párování

Fyzické propojení dvou krevních systémů (a tím i dvou vnitřních prostředí) u myši





Cell Stem Cell




Article

mTOR Mediates Wnt-Induced Epidermal Stem Cell Exhaustion and Aging

Rogério M. Castilho,¹ Cristiane H. Squarize,¹ Lewis A. Chodosh,² Bart O. Williams,³ and J. Silvio Gutkind^{1,*}


Complement C1q Activates Canonical Wnt Signaling and Promotes Aging-Related Phenotypes

Atsuhiko T. Naito,^{1,3} Tomokazu Sumida,⁴ Seitaro Nomura,⁴ Mei-Lan Liu,⁴ Tomoaki Higo,¹ Akito Nakagawa,¹ Katsuki Okada,¹ Taku Sakai,¹ Akihito Hashimoto,¹ Yurina Hara,¹ Ipppei Shimizu,⁴ Weidong Zhu,⁴ Haruhiro Toko,⁴ Akemi Katada,⁴ Hiroshi Akazawa,^{1,3} Toru Oka,^{1,3} Jong-Kook Lee,^{1,3} Tohru Minamino,⁴ Toshio Nagai,⁴ Kenneth Walsh,⁵ Akira Kikuchi,² Misako Matsumoto,⁶ Marina Botto,⁷ Ichiro Shiojima,^{1,3} and Issei Komuro^{1,3,4,*}



Nekanonická Wnt dráha

- e.g. Wnt5a





- do not induce axis duplication in Xenopus
- do not induce transformation of mammary cell line C57mg
- do not signal via nuclear translocation of β -catenin





SnapShot: Noncanonical Wnt Signaling Pathways

Mikhail V. Semenov,¹ Raymond Habas,² Bryan T. MacDonald,¹ and Xi He¹

¹Children's Hospital Boston, Harvard Medical School, Boston, MA 02115, USA; ²University of Medicine and Dentistry of New Jersey, Piscataway, NJ 08854, USA




Drosophila – PCP (planar cell polarity)



Molekulární mechanismus ustavení PCP

Box 1 | Molecular interactions between the Fz/PCP core factors

The molecular logic of the formation and separation of the Frizzled–Dishevelled–Diego (FZ–DSH–DGO) and Prickle–Strabismus (PK–STBM) complexes has started to be unravelled. In FIG. 2 are reported examples of the localization of each complex in various tissues. The figure is an apical view of two cells that have attained asymmetric localization of the two complexes. Several lines of



Seifert and Mlodzik, Nature Reviews in Genetics, 2007







Figure 2 | Subcellular distribution of core Fz/PCP factors in *Drosophila melanogaster* and vertebrates. a–c | Examples of cells with epithelial character (marked by grey shading). *Drosophila melanogaster* wing cells and eye R3 and R4 cells and mouse sensory hair cells in the cochlea (inner ear) are shown in a, b and c, respectively. d,e | Examples of dividing cells. The spindle orientation in the *D. melanogaster* sensory organ precursor (SOP) cells depends on the asymmetric distribution of the Frizzled (Fz)/planar cell polarity (PCP) factors (as shown in d), as does the orientation of neuroectodermal cells in zebrafish (as shown in e; note that during mitosis the asymmetric distribution of PK is lost and then re-established). Depending on the tissue, only a subset of the respective proteins has been analysed (the *D. melanogaster* wing is the only tissue in which all proteins were analysed; all but DSH have been analysed in the eye). These illustrations represent the localizations patterns of PCP proteins at the proposed time of signalling. In the wing, asymmetry of Flamingo (FMI) has been reported earlier, but the relevance of this is unknown⁹². Note that in the mouse inner ear (as shown in c) vang-like 2 (VANGL2) and FZ3/FZ6 localize to the same side of the cells; it is not known whether other Fz family members localize with the DSH homologues DVL1 and DVL2 to the opposite side. During zebrafish gastrulation (as shown in e) Prickle (Pk), which is represented by green circles, is cytoplasmic during cell division but regains polarity after separation of the daughter cell. Only PK has been analysed in this context, but its localization depends on the presence of Strabismus (STBM).


Defect in the non-canonical Wnt pathway in mammals

Changes in the "haircut"







Non-canonical/PCP (Planar cell polarity) pathway: phenotypes in mouse




Stereocilia orientation in inner ear hair cells



Qian et al., 2007, Dev. Biol.




Ror2



1321-1325del(5)
IVS8+3+5del3ins19
1398-1399insA

W749X
2249delG
Y755X
Q760X




ror2

C182X
R189W
R184C
R205X
R366W
R396X


Q502X
1740-1774del35
N620K
W720X

Mutations in *Ror2* cause dominant brachydactyly type B (BDB) and recessive robinow syndrome (RRS)



BDB



Known Wnt5a knockout phenotypes




Yamaguchi et al., 1999




Qian et al, 2007

Non-canonical/PCP (Planar cell polarity) in mouse (and human) convergent extension




Konvergentní extenze – migrace buněk směrem ke středu těla – vede k prodlužování tělní osy

Důsledky narušené konvergentní extenze (CE)



Exencephaly





urulation.m c

Open neural tube




Hamblet et al., 2002, Development


Možnosti studia CE - Kellerovy explantáty (Xenopus)




control



XLRP5 MO





XLRP5 MO
+ mLRp5




Wnt-induced assymetry

W-RAMP = Wnt-mediated
receptor-actin-myosin polarity








Možnosti studia migrace – Transwell assays (mammalian cells)




Attractant

Možnosti studia migrace – attractant/repellent assays

Attractant/repellent →







g



Primary cilia vs. motile (secondary) cilia

- struktura 9+0
- nepohyblivé
- téměř všechny buňky (www.primary-cilium.co.uk)
- solitérní

- struktura 9+2
- pohyblivé
- epitely tracheje, vejcovodů, ependym...



The primary cilium

Purification of Wnt ligands

.....

Wnt proteins are lipid-modified and can act as stem cell growth factors


Karl Willert*, Jeffrey D. Brown*, Esther Danenberg*, Andrew W. Duncan †, Irving L. Weissman ‡, Tannishtha Reya †, John R. Yates III § & Roel Nusse*

NATURE | VOL 423 | 22 MAY 2003 |


Wnt-3a



no detergent



detergent added



Frizzled – crucial receptor of most (all?) Wnt pathways






Figure 1. Schematic view of the 7TM model of FZDs. The model indicates extra- and intracellularly interacting proteins, putative glycosylation and phosphorylation sites. The N-terminal CRD is the primary binding site for ligands. The pink stretch in the C terminus indicates the internal PDZ-interacting motif (KTxxxW), which is absolutely conserved in the different FZD isoforms and necessary for DVL binding and signaling. The blue stretch at the far C terminus indicates the presence of a classical, less well-conserved PDZ-ligand sequence present in a subset of FZDs.

Lrp5/6 – crucial co-receptor of the canonical Wnt pathway




Destruction complex

- A working model for the destruction complex. (1) Initially, the destruction complex contains Axin, GSK3, CK1 and APC (with the 15 aa and 20 aa repeat regions shown). The complex contains other components such as PP2A, which are not shown here. (2) β -Catenin enters the complex by binding Axin and potentially the APC 15 aa repeats. This positions the N-terminus of β -catenin near CK1 and GSK3. (3) CK1 phosphorylates β -catenin at Ser45. (4) GSK3 phosphorylates β -catenin at, successively, Thr41, Ser37 and Ser33. (5) The 20 aa repeats, particularly repeat 3, are phosphorylated by a CK1 (and possibly GSK3) which greatly increases their affinity for β -catenin. The binding of a phosphorylated 20 aa repeat to β -catenin displaces Axin from β -catenin. (6) β -TRCP1 binds the phosphorylated N-terminus of β -catenin, causing the ubiquitination of β -catenin by an E2 ligase. APC is then either desphosphorylated within the complex, allowing the ubiquitinated β -catenin to leave the complex, or the ubiquitinated β -catenin bound to APC leaves the complex and is separated from APC at the proteasome. The complex then returns to Step 1



Důsledky aktivace β -catenin v epidermis (po depilaci)



Lo Celso, C. L. et al. Development 2004;131:1787-1799