

APPENDIX 1

The steps in derivation of the probability $p(n,x)$ for the production of an avalanche of n electrons at the distance x from the cathode are as follows:

Let $N(x)$ is the number of electrons emitted from the cathode which pass the distance x' from the cathode without any ionizing collision. Then

$$\begin{aligned} dN(x') &= -\alpha \cdot N(x) dx' \\ N(x') &= N_0 \cdot \exp(-\alpha x') \end{aligned}$$

where $N_0 = N(x'=0)$. As a consequence of this

$$(1) \quad p(1,x') = \frac{N(x')}{N_0} = \exp(-\alpha x')$$

Let the probability that the avalanche contains $n-1$ at x' is

$$(2) \quad p(n-1,x')$$

The probability that one and only one of these electrons will ionize in the region between x' and $x'+dx'$ can be found from the binomial distribution considering that $W(k,1) = p(n-1, 1)$, and $y = \alpha \cdot dx'$. Thus

$$p(n-1,1) = (n-1) \cdot \alpha \cdot dx' (1-\alpha \cdot dx')^{n-2}$$

for $dx' \approx 0$

$$(3) \quad p(n-1,1) \cong (n-1) \cdot \alpha \cdot dx'$$

The number of electrons in the avalanche has now increased from $n-1$ to n . The probability that none of these electrons will ionize in the region between $x'+dx'(\cong x')$ and x is:

$$(4) \quad [p(1,x-x')]^n = [\exp\{-\alpha \cdot (x-x')\}]^n = \exp\{-n \cdot \alpha \cdot (x-x')\}$$

where $p(1,x-x')$ is the probability that a single electron will not ionize between x' and x . If we take the product of expressions (2),(3) and (4), and integrate over x' , for $n > 1$, we get

$$\begin{aligned} p(n,x) &= \int_0^x p(n-1,x') \cdot p(n-1,1) \cdot [p(1,x-x')]^n dx' \\ p(n,x) &= \int_0^x p(n-1,x') \cdot (n-1) \cdot \alpha \cdot \exp\{-n \cdot \alpha \cdot (x-x')\} dx' \end{aligned} \quad (5)$$

$$p(n,x) = \exp(-n \cdot \alpha \cdot x) \cdot (\exp\{\alpha \cdot x\} - 1)^{n-1} =$$

The solution of the equation (5) is:

$$= \left(\frac{1}{n}\right)^n \cdot \left(\frac{-}{n-1}\right)^{n-1} = -\frac{1}{n} \cdot \left(1 - \frac{1}{n}\right)^{n-1}$$