

# Matrix Analyses

"Populační ekologie živočichů"

Stano Pekár

# Net reproductive rate $(R_0)$

• average total number of offspring produced by a female in her lifetime

$$R_0 = \sum_{x=0}^n l_x m_x$$

# Average generation time (T)

- average age of females when they give birth
- not valid for populations with generation overlap

$$T = \frac{\sum_{x=0}^{n} x l_x m_x}{R_0}$$

# **Expectation of life**

▶ age specific expectation of life – average age that is expected for particular age class

▶ *o* .. oldest age

$$e_x = \frac{T_x}{l_x}$$
 where  $T_x = \sum_x^o L_x$   $L_x = \frac{l_x + l_{x+1}}{2}$ 

# **Growth rates**

Discrete time/generations

- estimate of  $\lambda$  (finite growth rate) from the life table:

$$\mathbf{A}\widetilde{\mathbf{N}}_{t} = \lambda \widetilde{\mathbf{N}}_{t}$$

where  $\mathbf{N}_t$  is vector at stable age distribution  $\lambda$  is dominant positive eigenvalue of  $\mathbf{A}$ 

 $\det(\mathbf{A} - \lambda \mathbf{I}) = 0$ 

$$- \text{ or } \lambda \approx \frac{R_0}{T}$$

<u>Continuous time</u> *r* can be estimated from λ *r*by approximation

or by Euler-Lotka method

$$r = \ln(\lambda)$$

$$r \approx \frac{\ln(R_0)}{T}$$

$$1 = \sum_{x}^{\omega} l_{x} m_{x} e^{-rx}$$

#### **Stable Class distribution (SCD)**

relative abundance of different life history age/stage/size categories
population approaches stable age distribution:

 $N_0: N_1: N_2: N_3: ...: N_s$  is stable

- once population reached SCD it grows exponentially
- $\mathbf{w}_1$  .. right eigenvector (vector of the dominant eigenvalue)
- provides stable age distribution
- scale  $\mathbf{w}_1$  by sum of individuals

$$\mathbf{A}\mathbf{w}_1 = \lambda_1 \mathbf{w}_1$$





## **Reproductive value** $(v_x)$

 measures relative reproductive potential and identifies age class that contributes most to the population growth
 such class is under highest selection force

- when population increases then early offspring contribute more to  $v_x$  than older ones
- ▶ is a function of fertility and survival

$$\mathbf{v}_1 \mathbf{A}' = \lambda_1 \mathbf{v}_1$$

▶ v<sub>1</sub> .. left eigenvector (vector of the dominant eigenvalue of transposed A)

-  $\mathbf{v}_1$  is proportional to the reproductive values and scaled to the first category

$$v_x = \frac{v_{1x}}{v_{11}}$$

 $x \neq 1$ 



#### Sensitivity (s)

• identifies which process (p, F, G) has largest effect on the population increase  $(\lambda_1)$ 

- examines change in  $\lambda_1$  given small change in processes  $(a_{ij})$
- sensitivity is larger for survival of early, and for fertility of older classes
- not used for postreproductive census with class 0

$$s_{ij} = \frac{v_{ij} w'_{ij}}{\langle \mathbf{v}, \mathbf{w} \rangle} \leftarrow \text{sum}$$

- sum of pairwise products

Elasticity (e)

- weighted measure of sensitivity
- measures relative contribution to the population increase
- impossible transitions = 0

$$e_{ij} = \frac{a_{ij}}{\lambda_1} s_{ij}$$

# **Conservation biology**

to adopt means for population promotion or control

# **Conservation/control procedure**

- 1. Construction of a life table
- 2. Estimation of the intrinsic rates
- 3. Sensitivity analysis helps to decide where
- conservation/control efforts should be focused
- 4. Development and application of management plan
- 5. Prediction of future