

## **Product Data Sheet**

## APC/Cy7 anti-mouse/human CD44

Catalog # / Size:	1115140 / 100 μg 1115135 / 25 μg
Clone:	IM7
Isotype:	Rat IgG2b, к
Immunogen:	Dexamethasone-induced myeloid leukemia M1 cells
<b>Reactivity:</b>	Human
Preparation:	The antibody was purified by affinity chromatography, and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.2





## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq$ 1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	Clone IM7 has been reported to recognize an epitope common to alloantigens and all isoforms of CD44 <sup>17,18</sup> that is located between amino acids 145 and 186 <sup>20</sup> . Additional reported applications (for the relevant formats) include: immunohistochemistry of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections <sup>6,7</sup> , complement-mediated cytotoxicity1, immunoprecipitation <sup>1,3</sup> , and <i>in vivo</i> inhibition of DTH <sup>4,5</sup> . The LEAF <sup>™</sup> purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 103014). For highly sensitive assays, we recommend Ultra-LEAF <sup>™</sup> purified antibody (Cat. No. 103046) with a lower endotoxin limit than standard LEAF <sup>™</sup> purified antibodies (Endotoxin <0.01 EU/microg).
Application References:	<ol> <li>Trowbridge IS, <i>et al.</i> 1982. <i>Immunogenetics</i> 15:299. (ICFC, IP, CMCD)</li> <li>Katoh S, <i>et al.</i> 1994. <i>J. Immunol.</i> 153:3440. (ELISA)</li> <li>Budd RC, <i>et al.</i> 1987. <i>J. Immunol.</i> 138:3120. (IP)</li> <li>Camp RL, <i>et al.</i> 1993. <i>J. Exp. Med.</i> 178:497. (Block)</li> <li>Weiss JM, <i>et al.</i> 1997. <i>J. Cell Biol.</i> 137:1137. (Block)</li> <li>Frank NY, <i>et al.</i> 2005. <i>Cancer Res.</i> 65:4320. (IHC) PubMed</li> <li>Cuff CA, <i>et al.</i> 2001. <i>J. Clin. Invest.</i> 108:1031. (IHC)</li> <li>Lee JW, <i>et al.</i> 2005. <i>J. Immunol.</i> 174:6967. PubMed</li> <li>Huabiao C, <i>et al.</i> 2005. <i>J. Immunol.</i> 175:591. PubMed</li> <li>Gui J, <i>et al.</i> 2007. <i>Int. Immunol.</i> 19:1201. PubMed</li> <li>Kenna TJ, <i>et al.</i> 2008. <i>Blood</i> 111:2436. PubMed</li> <li>Kenna TJ, <i>et al.</i> 2009. <i>Blood</i> PubMed</li> <li>Kmieciak M, <i>et al.</i> 2009. <i>J. Transl. Med.</i> 7:89. (FC) PubMed</li> <li>Chen YW, <i>et al.</i> 2010. <i>Mol. Cancer Ther.</i> 9:2879. PubMed</li> </ol>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

	<ol> <li>Zheng Z, <i>et al.</i> 1995. <i>J. Cell. Biol.</i> 130:485.</li> <li>Wiranowska M, <i>et al.</i> 2010. <i>Int. J. Cancer</i> 127:532.</li> <li>Hirokawa Y, <i>et al.</i> 2014. <i>Am J Physiol Gastrointerest Liver Physiol.</i> 306:547.</li> <li>PubMed</li> <li>Sandmaier BM, <i>et al.</i> 1998. <i>Blood</i> 91:3494.</li> <li>Nagaoka M, <i>et al.</i> 2014. <i>J Immunol.</i> 193:2812. PubMed</li> <li>Young KH, <i>et al.</i> 2014. <i>Cancer Immunol Res.</i> 2:1011. PubMed</li> </ol>
Description:	CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.

Antigen
1. Barclay AN, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
2. Haynes BF, *et al.* 1991. *Cancer Cells* 3:347.
3. Goldstein LA, *et al.* 1989. *Cell* 56:1063.
4. Mikecz K, *et al.*