

Diffusion

- Interstitial

- Fick's first law

$$J_B = -D_B \frac{\partial C_B}{\partial x}$$

$$D_B = D_0 e^{-\frac{Q}{RT}}$$

- Intrinsic diffusivity

- Fick's second law

$$\frac{\partial C_B}{\partial t} = D_B \frac{\partial^2 C_B}{\partial x^2}$$

- Substitutional

- Self diffusion, vacancy diffusion

- Lattice fixed frame of reference, drifting lattice

- Kirkendall effect: net flow of vacancy: $J_v = -(J_A + J_B)$

- Darken equation

$$\tilde{J}_B = -\tilde{D} \frac{\partial C_B}{\partial x}$$

$$\tilde{D} = x_A D_B + x_B D_A$$

course@calphad.org

3

Frames of reference

- Lattice-fixed frame of reference:

- Defined by the inert markers, or that there is no net flow of lattice sites.

$$\sum_{K=1}^n J_K = -J_{Va}$$

- Volume-fixed frame of reference:

- Defined in such a way that there is no net flow of volume.

$$\sum_{K=1}^n \tilde{J}_K V_K = 0$$

- DICTRA frame of reference

- Volume-fixed frame, however only the substitutional components are assumed to contribute to the volume

$$\sum_{K \in S} \tilde{J}_K = 0$$

$V_{inheritor} = \phi$
 $V_{subst} = \text{step in } \rho \text{ to } \text{isod}$

course@calphad.org

4

Frames of reference

The equation for the velocity of the markers in the diffusion zone relative to the ends of the diffusion couple, in the Kirkendall experiment, is:

$$v = \sum_{k=1}^n V_k J_k$$

Velocity of marker (circled around v)

← k₂ J₂ (arrow pointing to the sum)

This is the velocity with which the two frames move relative to each other.

We may use this equation in order to transfer the fluxes from the volume-fixed to the lattice-fixed frame, or vice versa.

$$J'_k = J_k - c_k \sum_{k=1}^n V_k J_k$$

lattice fixed (circled around J'_k)

Volume fix (circled around J_k)

not $\frac{1}{m^2}$ mol $\frac{1}{m^2}$ mol $\frac{1}{m^3}$ (handwritten notes above the equation)

15 (handwritten note below the sum)

course@calphad.org

za prvot poloze dle
že s₂ stálo udržen
rovnovážné
vážnosti

Atomistic treatment of diffusion

For crystalline phases it's usually believed that diffusion occurs through a vacancy exchange mechanism.

Assuming that there is a random distribution of vacant sites and that the number of vacancies is everywhere adjusted to equilibrium, it's possible to derive the following expression for the flux of k in a lattice-fixed frame of reference:

$$J_k^{LF} = -c_k y_{va} M_{kva} \nabla \mu_k$$

lattice fixed (circled around J_k^{LF})

→ $\frac{\partial \mu_k}{\partial t}$ (handwritten note with arrow pointing to the right)

where M_{kva} is some kinetic factor which gives the rate of exchange if there is a vacancy adjacent to a k atom.

o o o
o o o
o o o
o o o
o o o

rektor

Phenomenological equations

$$J_k^L = - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z} - L_{1T} \frac{\partial T}{\partial z} - L_{1P} \frac{\partial P}{\partial z} - L_{1\phi} \frac{\partial \phi}{\partial z}$$

They are called phenomenological since they stem from no model, but from the observed conditions of equilibrium.

If we choose to consider an isothermal, isobaric and isopotential system we have:

$$J_k^L = - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z} \quad \left(J_k^L = - L_{kk} \frac{\partial \mu_k}{\partial z} \right)$$

course@calphad.org

7

Mobility vs L Parameter

Assuming that the vacancy exchange mechanism is predominant, and by comparing to the expression derived earlier under this assumption, we may identify:

$$L_{kk} = c_k y_{va} M_{kva} \quad (1)$$

$$L_{ki} = c_i y_{va} M_{iva}$$

We have now established a relation between M and L.

course@calphad.org

8

Phenomenological equations

$$J_k^L = - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z} - L_{1T} \frac{\partial T}{\partial z} - L_{1P} \frac{\partial P}{\partial z} - L_{1\phi} \frac{\partial \phi}{\partial z}$$

They are called phenomenological since they stem from no model, but from the observed conditions of equilibrium.

If we choose to consider an isothermal, isobaric and isopotential system we have:

$$J_k^L = - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z} \quad \left(J_k^L = - L_{kk} \frac{\partial \mu_k}{\partial z} \right)$$

course@calphad.org

7

Mobility vs L Parameter

Assuming that the vacancy exchange mechanism is predominant, and by comparing to the expression derived earlier under this assumption, we may identify:

$$L_{kk} = c_k y_{va} M_{kva} \quad (1)$$

$$L_{ki} = c_i y_{va} M_{iva}$$

We have now established a relation between M and L.

course@calphad.org

8

by guid

$$= - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z} - c_k \sum_{k=1}^n V_k \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z} \rightarrow \rightarrow$$

$$J_k = - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial z}$$
$$L'_{ki} = - \sum_{j=1}^n (\delta_{jk} - c_k V_j) L_{ji} \quad (2)$$

•

11

$$J_k = - \sum_{i=1}^n L_{ki} \frac{\partial \mu_i}{\partial c_{\vec{A}}} \frac{\partial c_{\vec{A}}}{\partial z} \quad ?$$

Or equally if the diffusivities, D_{ki} , are introduced:

$$J_k = - \sum_{j=1}^n D_{kj} \frac{\partial c_j}{\partial z}$$

$$D_{kj} = \sum_{i=1}^n L'_{ki} \frac{\partial \mu_i}{\partial c_j}$$

$$\mathbb{D} = \mathbb{L} \oplus \Phi$$

(3)

(3a)

10

Independent set of driving forces

There is a relation between the n concentration gradients, and it's possible to eliminate one of them:

$$\frac{\partial c_n}{\partial z} = -\frac{\partial c_1}{\partial z} - \frac{\partial c_2}{\partial z} - \dots - \frac{\partial c_{n-1}}{\partial z}$$

and thus, the flux is now expressed: $J_k = -\sum_{j=1}^{n-1} (D_{kj} - D_{kn}) \frac{\partial c_j}{\partial z}$

we may identify: $D_{kj}^n = D_{kj} - D_{kn}$ (4) (4a)

and finally obtain: $J_k = -\sum_{j=1}^{n-1} D_{kj}^n \frac{\partial c_j}{\partial z}$

Time Dependence


$$\frac{\partial c_k}{\partial t} = -\text{div}(J_k)$$

course@calphad.org

11

very good

Summary of steps taken when transforming from M's to D's



course@calphad.org

12

Atomic mobility

$$M_i = \frac{M_i^0}{RT} \cdot e^{-\frac{Q_i}{RT}} = \frac{1}{RT} e^{\frac{-Q_i + RT \ln(M_i^0)}{RT}}$$

- Tracer diffusivity $D_i^* = RTM_i$

course@calphad.org

13


Moving Boundary Problems

- Sharp Interface
 - Zero thickness
 - Interface tracking
 - Interfacial energy
 - Local equilibrium and deviation from local equilibrium
 - Typically simple geometry
- Diffuse Interface (Day 2)
 - Finite interface width
 - Field variables across a boundary continuously
 - Gradient energy to represent the interfacial energy
 - Complex geometry

course@calphad.org

14

Moving Phase Boundary




Unknowns: Tie-line, specified by $n-2$ a_i or μ_i
Velocity of phase boundary, v

Equations: $n-1$ flux-balance equations, $v(c_k^\alpha - c_k^\gamma) = J_k^\alpha - J_k^\gamma$

Solved as: $v(c_k^\alpha - c_k^\gamma) - (J_k^\alpha - J_k^\gamma) = 0$

course@calphad.org

15



16


What is DICTRA?

- Software for diffusion simulations
 - One dimensional (planar,cylindrical,spherical)
 - Linked to Thermo-Calc for thermodynamics→Has all modules in Thermo-Calc plus a Dictra module.
 - Quantitative description of multicomponent diffusion
 - Simulations based on databases for thermodynamics and diffusion
 - Written in FORTRAN, runs on most platforms

course@calphad.org

17

Numerical Calculation of Diffusional Reaction



course@calphad.org

18

Mobility databases

MOB 2

The most general diffusion database.

Could be used for Steels, Nickel alloys, Aluminium alloys and more.

75 Elements: Ag, Al, Am, As, Au, B, Ba, Be, Bi, C, Ca, Cd, Co, Cr, Cs, Cu, Dy, Er, Fe, Ga, Gd, Ge, Hf, Hg, Ho, In, Ir, K, La, Li, Lu, Mg, Mn, Mo, N, Na, Nb, Nd, Ni, Np, Os, P, Pa, Pb, Pd, Pr, Pt, Pu, Rb, Re, Rh, Ru, S, Sb, Sc, Se, Si, Sm, Sn, Sr, Ta, Tb, Tc, Te, Th, Ti, Tl, Tm, U, V, W, Y, Yb, Zn and Zr

Phases with diffusion data: BCC_A2 CEMENTITE FCC_A1 FE4N HCP_A3 LIQUID

Assessed data for the following binary systems

BCC_A2: C-Fe, C-Cr, Cr-Fe, Cr-N, Cr-Ni, Fe-N, Fe-Ni
FCC_A1: Al-Cr, Al-Ni, C-Fe, C-Ni, Cr-Fe, Cr-Ni, Fe-N, Fe-Ni, Fe-Si
HCP_A3: C-Fe, Fe-N
FE4N: C-Fe, Fe-N

Assessed data for the following ternary systems

BCC_A2: C-Cr-Fe
FCC_A1: Al-Cr-Ni, C-Cr-Fe, C-Fe-Ni


Assessed data for the following higher order systems

BCC_A2: C-Cr-Fe-N-Ni
FCC_A1: C-Cr-Fe-Ni


course@calphad.org

19

System



Cell



course@calphad.org

20


Boundary conditions in DICTRA

- Boundary Conditions can be specified as functions of time, temperature and pressure.
- Different functions may be used in different time intervals.

Example of conditions are:

- Closed system (default)
- State variable expressions
- Flux conditions
- Mixed conditions

Application example:



course@calphad.org

21

ENTER_GEOMETRICAL_EXPONENT

Command that determines the geometry of the system. The program handles problems that can be reduced to 1-dimensional geometries. Default value is zero, i.e. planar geometry.

Exponent Geometry



0 Planar geometry.

This corresponds to an infinitely wide plate of a certain thickness.



1 Cylindrical geometry.

This corresponds to an infinitely long cylinder of a certain radius.




2 Spherical geometry.

A sphere with a certain radius.


course@calphad.org

22


Some different possible geometries



Growth or dissolution
of spherical precipitate



Growth of spherical film
(Grain-boundary film)



Planar growth




Growth of cylindrical
precipitate

course@calphad.org

23

Region




course@calphad.org


24

Region

Matrix phase




Lamella phase



course@calphad.org

25


Grid points



course@calphad.org

26

Microstructure vs Dictra




course@calphad.org

27

Multi-cell Approach

$\sum J_i = 0$
Diffusion potential = constant




course@calphad.org


28

Some DICTRA applications


Carbide dissolution




Coarsening



Carbon diffusion in a weld between two steels




Microsegregation during solidification



Lippard et al. *Metall. Mater. Trans. B* 1998

Carburizing



course@calphad.org

29

More DICTRA applications

- Homogenisation of alloys
- Carburization and decarburization
- Carburization of alloys
- Nitriding of steels
- Diffusion during sintering of cemented carbides
- Nitrocarburizing of steels
- Austenite/ferrite diffusional transformations in steels
- Growth or dissolution of individual particles
- Transient liquid phase bonding of alloys
- Calculation of TTT-diagrams
- Interdiffusion between coating/substrate
- Coarsening of a particle distribution
- Gradient sintering of cemented carbides
- Growth of pearlite in alloyed steels
- Sigma phase precipitation in stainless steel
- Post weld heat treatment of welds between dissimilar materials


course@calphad.org

30

Moving Phase Boundary

- Moving phase boundaries simulations may be setup in DICTRA in two different ways:

- Introducing two or more adjacent regions containing different phases



- Entering an inactive phase (formed when thermodynamically stable)




course@calphad.org

31

Possible Cell Calculations

- Simulation of dissolution of a size distribution of precipitates
- Simulation of diffusion at an immobile phase interface
- Coupled simulation of dissolution with precipitates of different phases



$$J_1^{Cell\#1} = J_1^{Cell\#2}$$

$$J_2^{Cell\#1} = J_2^{Cell\#2}$$

$$\mu_1^{Cell\#1} - \mu_n^{Cell\#1} = \mu_1^{Cell\#2} - \mu_n^{Cell\#2}$$

$$\mu_2^{Cell\#1} - \mu_n^{Cell\#1} = \mu_2^{Cell\#2} - \mu_n^{Cell\#2}$$

course@calphad.org

32

Cell calculations in DICTRA

- Makes it possible to couple several computational cells
- Conditions for cell boundaries:
 - ✓ Equal diffusion potentials Φ_i for the elements

$$\Phi_i = \mu_i - \mu_n \quad \text{for subst. elements}$$

$$\Phi_i = \mu_i \quad \text{for interstitial elements}$$

- ✓ Flux balances to conserve the mass of the elements

$$\frac{J_i^{cell\#1}}{n^{cell\#1}} = \frac{J_i^{cell\#2}}{n^{cell\#2}}$$


n , the number of identical cells

course@calphad.org

33

Disperse system model

- Assumptions:
 - Diffusion takes place in the matrix phase only.
 - Equilibrium holds locally on each grid point.
- Effective diffusivity
 - Diffusion matrix, D_{kj} , is multiplied with a labyrinth factor $f(\text{volfr})$, volfr being the volume fraction of the matrix phase.
 - The labyrinth factor is often taken as volfr^2 .



course@calphad.org

34

Examples

- Diffusion in one phase: Homogenization of an Fe-Ni alloy.
- Moving boundary: Dissolution of carbide in one cell and in three cells.
- Diffusion in dispersed system: Carburization of the Ni-25% Cr alloy.
- Exercise: γ to α transformation in the Fe-0.15C alloy

course@calphad.org

35

Simulation Scheme

1. Get Gibbs energy and mobility
2. One cell simulation
3. Set geometry, temperature, time
4. Set boundary conditions (default: closed system)
5. Enter regions
6. Enter grid in region
7. Enter phases in regions
8. Enter concentration profiles of phases

course@calphad.org

36

Dictra: Common Commands

LIST_MOBILITY_DATA
CREATE_NEW_CELL
SET_CONDITION
ENTER_REGION
ENTER_GRID_COORDINATES
ENTER_PHASE_IN_REGION
ENTER_COMPOSITIONS
ENTER_GEOMETRICAL_EXPONE
SET_SIMULATION_TIME
SIMULATE_REACTION
POST_PROCESSOR

course@calphad.org

37

Post: Common Commands

SET_DIAGRAM_AXIS
SET_INDEPENDENT_VARIABLE
SET_PLOT_CONDITION
MAKE_EXPERIMENTAL_DATAFILE
PLOT_DIAGRAM

course@calphad.org

38

Variables in the Post-Processor

State variables

T, P, Amount of components (e.g. $X(\text{Cr})$, $W(\text{Cr})$, $W(\text{Fcc}, \text{Cr})$, etc.)

Amount of a phase, e.g. $NP(\text{Fcc})$

Activity for a component: $AC(\text{Cr})$

Chemical Potential for a component: $MU(\text{Cr})$

Energetic properties: G, H, S, A

course@calphad.org

39

Variables in the Post-Processor

Integral variables (Integration in space)

Variables constructed in the following way:

First letter always I for integral variable, second letter specifies quantity

N (number of moles)

W (mass)

V (volume) and many more..

Third letter (optional) specifies normalizing quantity e.g. V (total volume).

course@calphad.org

40

Variables in the Post-Processor

Integral variables, *Examples*

IVV(CARBIDE)

Volume of region CARBIDE divided with total volume,
i.e. volume fraction of carbide

IW(CARBIDE,CEMENTITE,CR)

The mass of Cr in the cementite phase in region carbide

course@calphad.org

41

Variables in the Post-Processor

Auxiliary variables "Special variables" e.g.

POSITION-OF-INTERFACE

VELOCITY-OF-INTERFACE

LOGDT(phase,J)

LOGDC(phase,J,K,N)

User-defined functions

course@calphad.org

42


Plotting in DICTRA

- Two independent variables in a DICTRA-simulation, TIME and DISTANCE
- When plotting a diagram we need to specify one of them i.e. use it as plot condition. Command: SET-PLOT-CONDITION
- TIME can be specified as one or several values.
- When plotting integral quantities, time is automatically chosen as independent variable.

course@calphad.org

43

Exa1: Homogenization of an Fe-Ni alloy (Initially a linear concentration profile)





course@calphad.org

44

Input of T and c

- Temperature (T) can be entered as a function of time
- Many different functions can be used (+, -, *, **, SQRT(X), EXP(X), LOG(X), SIN(X))
- Concentration can be entered as a function of distance or read from a file
- Many functions e.g. error-functions (erf(x)) and heavy-side step functions (hs(x))





course@calphad.org

45

Cementite dissolution in a Fe-Cr-C alloy

Dissolution of cementite at 910C (or 1183K):



Initial particle radius is estimated to 0.5255 μm .

course@calphad.org

46

Cementite dissolution in a Fe-Cr-C alloy

The volume fraction of cementite and the composition in the cementite, is calculated at the normalizing temperature 735C (1008K).

The size of the γ region is calculated from:

$$\frac{R_{\text{cementite}}^3}{R_{\text{tot}}^3} = \frac{V_{\text{cementite}}}{V_{\text{tot}}} = V_{\text{cementite}}^f$$

$$\Rightarrow R_{\gamma} = R_{\text{tot}} - R_{\text{cementite}} = \frac{R_{\text{cementite}}}{\sqrt[3]{V_{\text{cementite}}^f}} - R_{\text{cementite}}$$

$$\left(V_{\text{cementite}}^f = \frac{n(\text{cem}, \text{Cr}) + n(\text{cem}, \text{Fe})}{n(\text{Cr}) + n(\text{Fe})} \right)$$


course@calphad.org

47

Cementite dissolution in a Fe-Cr-C alloy

Cell calculation with size-distribution

Dissolution of cementite at 910C (or 1183K):



Instead of assuming an average particles size as in previous example, we will perform the simulation for a particle distribution using three cells of different sizes.

course@calphad.org

48

Conditions at cell boundaries

$$\sum \frac{J_i}{n^{cell}} = 0$$

$$\mu_1^{Cell\#1} - \mu_n^{Cell\#1} = \mu_1^{Cell\#2} - \mu_n^{Cell\#2}$$


$$\mu_2^{Cell\#1} - \mu_n^{Cell\#1} = \mu_2^{Cell\#2} - \mu_n^{Cell\#2}$$

course@calphad.org

49

Exd1: Carburization of a Ni-25% Cr Alloy

T=1123 K




course@calphad.org

50

Set-rod glob t 0 973-time ; 190 y 783 ; * h

Exercise: γ to α transformation in Fe-0.15C with the following temperature variation with time




course@calphad.org

51

100 (bcc)

Suggestions



course@calphad.org

52