


HYBRIDIZATION AND HYBRID ZONES





Secondary contact

ring species:

Chorthippus parallelus

postglacial expansion



Ensatina eschscholtzii - *klauberi*



greenish warbler (*Troglodytes troglodytes*)
Larus argentus/*L. fuscus*

Hybridization:

25% species of vascular plants

10% species of animals

probably underestimation (only conspicuous species: ducks, birds of paradise, butterflies)

often result of environmental disturbance:

eg. „Darwin’s finches“ *Geospiza fuliginosa*, *G. fortis* and *G. scandens* after El Niño event



Geospiza fuliginosa




G. fortis



G. scandens

Possible outcomes of hybridization



Program NewHybrids:

A Model-Based Method for Identifying Species Hybrids Using Multilocus Genetic Data

E. C. Anderson¹ and E. A. Thompson

Department of Statistics, University of Washington, Seattle, Washington 98195

Manuscript received October 3, 2001

Accepted for publication December 24, 2001

ABSTRACT

We present a statistical method for identifying species hybrids using data on multiple, unlinked markers. The method does not require that allele frequencies be known in the parental species nor that separate, pure samples of the parental species be available. The method is suitable for both markers with fixed allelic differences between the species and markers without fixed differences. The probability model used is one in which parentals and various classes of hybrids (F_1 's, F_2 's, and various backcrosses) form a mixture from which the sample is drawn. Using the framework of Bayesian model-based clustering allows us to compute, by Markov chain Monte Carlo, the posterior probability that each individual belongs to each of the distinct hybrid classes. We demonstrate the method on allozyme data from two species of hybridizing trout, as well as on two simulated data sets.







FIGURE 1.—Six arrangements of founders on a pedigree of $n = 2$ generations. Each box represents a locus. The circles within each box represent the two genes possessed by the diploid organism at the locus. The founders are the individuals in the top row of each pedigree. Black gene copies are those originating from the species *A* population, and the white genes are from species *B*. Genes that are not determined to be either black or white by the pedigree and the founders in it are denoted by broken circles. The individual at the bottom of each pedigree belongs to a different hybrid class, determined by the arrangement of species among the founders. a–f represent six distinct *genealogical classes*. a–f also represent six distinct *genotype frequency classes*. There are, however, only five distinct *gene frequency classes*; the individuals at the bottoms of pedigrees c and f are both in the same gene frequency class.






(a) STRUCTURE $K = 2$



(b) NEWHYBRIDS



Hybrid zone (Barton a Hewitt 1985)

= area, where genetically different populations meet, mate and give rise at least some hybrid offspring

Hybrid zones may be classified as:

primary

secondary

tension, mosaic, staggered, „mottled“ ...

extrinsic selection (external environment)


intrinsic selection (prezygotic or postzygotic barriers)

Mosaic hybrid zone:

influence of environment


in fact a set of several hybrid zones

eg.: *Gryllus firmus* x *G. pennsylvanicus* (NE USA)
sandy x clayish soils




Iris fulva x *I. brevicaulis*:

I. fulva is limited to more forested sites




Bombina:



fire-bellied toad
B. bombina:

lowlands
mostly in water
larger water surfaces
thinner skin
territorial
530 Hz
longer development

mosaic HZ in Croatia,
not in Poland



yellow-bellied toad
B. variegata:

hills, highlands
terrestrial
mating in puddles
thick skin
nonterritorial
580 Hz
shorter development




Nick Barton


Most hybrid zones are
tension zones.

... i.e., they are maintained by balance between dispersal and selection (Barton & Hewitt, 1985)

Tension zone is when...




Tension zone is when...




⇒ zone widening

Tension zone is when...




⇒ zone narrowing

Tension zone is maintained by dynamic equilibrium
between *dispersal* and *selection*




tension zone moves
along the gradient of
population density


Tension zone is independent of external conditions (*intrinsic selection*)
⇒ its movement ends at a geographical barrier or in the area
of the lowest population density („*population/density trough*“)




„population trough“




tension zone moves
along population gradient






Theory of cline:

secondary contact:  coincident and concordant clines




Cline = gradient of trait(s) (eg. allele frequency or mean of quantitative trait) across spatially continuous habitat

clines




coincidence = same positions of zone centres



concordance = same cline shapes
(in practice usually same widths)


secondary contact

cline




Theory of cline:


neutral vs. selected loci




diffusion of neutral marker




diffusion of neutral marker



diffusion of neutral marker




diffusion of neutral marker




neutral vs. selected loci

with time, concordance is disappearing ...



... but (in tension zone) selection pushes clines for individual loci to each other
⇒ maintains coincidence

sometimes ...




locus 1

locus 2

... but clines still parallel

cline models (diffusion approximation etc.), linkage disequilibrium, evolutionary parameters



problem, how to analyse

Hybrid zone study

1. Sampling along linear or 2D transect, geographic coordinates of localities
2. Genetic (morphological, behavioural etc.) analysis
... problem of sample independence (F_{ST} , F_{IS} ... effective No. alleles)
3. Geographic clines
4. Estimation of dispersal, selection, and other parameters
5. Alternative approaches:
 - monotonic clines
 - 2D analysis
 - genomic clines
 - concordance analysis

Case study: house mouse hybrid zone



musculus



domesticus



Mouse colonization of Europe


Neolithic



Cucchi et al. (2005)

Mouse colonization of Europe


Bronze and Iron Age



Cucchi et al. (2005)

Mouse colonization of Europe


Late Neolithic



Cucchi et al. (2011)

Hybrid zone in Europe

musculus



time of origin unknown...




domesticus

Hybrid zone in Europe












Co tuto zónu ovlivňuje?



oceanic climate


continental climate

- | | |
|--|--|
|  Semi-arid |  Humid continental |
|  Subtropical dry summer |  Subarctic |
|  Humid subtropical |  Tundra |
|  Humid oceanic |  Highland |



climatic factors don't determine

Hybrid zone in Europe




Most hybrid zones are „tension zones“




Nick Barton





also the mouse hybrid zone?



hybrid zone course may be complex....


... moreover, usually we don't know a priori, or we extrapolate from global direction






Real local cline


**Cline interpolated
from widescale survey**



2D → 1D clines



Multiple genes:



„stepped“ model (symmetrical, asymmetrical)

linkage disequilibrium resulting from influx of parental allele combinations \Rightarrow
synergistic effect: strengthening of selection in zone centre \Rightarrow central step
 \times introgression tails reflect selection at individual loci

We can estimate some other key evolutionary parameters
from LD and cline parameters:

dispersal:

effective selection:

selection on marker loci:


selection on selected loci:

fitness of hybrids:


number of loci under selection:

model comparison: LRT (they are nested); d.f. = difference in number of parameters


likelihood profiles:




hybrid zone




hybrid index



selected areas





hybrid index




centromere


molecular markers


Problems – Y chromosome





Monotonic clines – consensus orientation










salient/invagination $\approx 330 \text{ km}^2$

Weird behaviour of the Y in the hybrid zone – summary:

1. *musculus* Y more successful than *domesticus* Y on its own genetic background
2. higer proportion of males relative to other areas

Either coincidence, or ...


... or genetic conflict between X and Y
and probably some autosomal genes as well





Gregor Mendel



segregation
law



Intragenomic conflict results in higher proportion of a genomic element in the next generation










Gregor Mendel



vychýlení segregacího (transmisního) poměru


= segregation distortion (SD)

= transmission ratio distortion (TRD)




X chromosome

Chr. X - 2D analysis Geneland




'Genomic clines'


A. Underdominance




C. Epistasis





B. Overdominance












D. Directional selection











Concordance analysis:






S.J.E. Baird



very steep cline




strong introgression in both directions



S.J.E. Baird










Concordance analysis:

introgression to the left




S.J.E. Baird

Concordance analysis:



S.J.E. Baird


Proximal marker on the X





non-introgressed localities

Recombination reduces size of introgressed block far of the zone centre

Using cline model for analysis of reinforcement – odour preference in the mouse hybrid zone



Using cline model for analysis of gene expression in the mouse hybrid zone – asymmetric model





Neanderthal DNA specialist Svante Pääbo examines the femur, found near Ust'-Ishim in western Siberia. Photograph by [unreadable]

Blocks of Neanderthal DNA found in modern humans can act like a biological clock, because they are fragmented more and more with each generation since interbreeding happened. The **blocks** of Neanderthal DNA in the Siberian man were on average three times longer than those seen in people alive today. Working backwards, the scientists calculate that Neanderthals contributed to the man's genetic ancestry somewhere between 7,000 and 13,000 years before he lived.

The findings, published in the journal Nature, suggest that humans and Neanderthals had reproductive sex around 50,000 to 60,000 years ago...


P




X




F1



F2




F3




⋮

Hybridization makes a cascade of blocks




advancing introgression into 'blue' genome



R. A. Fisher

Recombination brings together DNA of different origin and makes *junctions* (breakpoints) they divide genome into *blocks* (chunks, tracts, segments)




Romania, ~40 kya,
mating before
200–100 years


chromosome 12
of Neanderthal

Siberia, ~45 kya,
mating before
8000–5000


contemporary China,
54–49 kya



Chromosome 17



Chromozom X



cM 10 20 30 40 50 60 70

Why hybrid zones? Reproductive barriers and speciation!

Dobzhansky-Muller model



W. Bateson

T. Dobzhansky

H. Muller





Fig. 1 The Dobzhansky-Muller model for postzygotic isolation

„Arms races“ and secondary contact



genetic conflict: "classical" scenario



arms race
in ancestral
population



subpop. 1



incompatible!

secondary contact

continuing
arms race

subpop. 2



MAD = *mutually assured destruction*

"speciation genes"

genetic conflict: alternative scenario



arms race
in ancestral
population




subpop. 1

continuing
arms race

subpop. 2

*Never bring a knife
to a gunfight!*




secondary contact

the winner thrives on
“naive” genetic
background



“antispeciation
genes”

Why we don't see this more often?



Ticking time-bomb...



Cytonuclear disequilibria

- = non/random associations of nuclear and cytoplasmic (mitochondrial) alleles
- 3×2 table

	nuclear genotype:			
mtDNA:	<i>AA</i>	<i>Aa</i>	<i>aa</i>	total
<i>M</i>	u_1	v_1	w_1	x
<i>m</i>	u_2	v_2	w_2	y
total	u	v	w	1

No hybridization

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	+++	0	0
<i>m</i>	0	0	+++

Random mating, hybrid swarm

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	obs=exp	obs=exp	obs=exp
<i>m</i>	obs=exp	obs=exp	obs=exp

Hybridization without apparent introgression, crossing independent of sex

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	++	obs=exp	0
<i>m</i>	0	obs=exp	++

Hybridization without apparent introgression, crossing depends on sex

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	++	++	0
<i>m</i>	0	--	++

Hybrids mate more often with less discriminating species

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	obs=exp	++	--
<i>m</i>	obs=exp	--	++

Symmetrical introgression

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	++	obs=exp	--
<i>m</i>	--	obs=exp	++

Potential introgression, crossing dependent on sex

	nuclear genotype:		
mtDNA:	AA	Aa	aa
<i>M</i>	++	++	--
<i>m</i>	0	0	++