



Oddělení funkční genomiky a proteomiky
Národní centrum pro výzkum biomolekul
Přírodovědecká fakulta MU



Charakterizace proteinů hmotnostní spektrometrií

C7250

Část IV

Zbyněk Zdráhal

*Výzkumná skupina Proteomika, CEITEC-MU
Centrální laboratoř-Proteomika, CEITEC-MU
Funkční genomika a proteomika, NCBR PřF
zdrahal@sci.muni.cz*

Charakterizace Modifikací Proteinů



Proč?

počet druhů PTMs

> 400

počet PTMs

≈ 90 000 (experimentálne identifikovaných)

≈ 230 000 (predikce)

(SwissProt, per ≈ 530 000 proteinů)

G. A. Khoury et al., Sci. Rep. 1, 90; (2011); <http://selene.princeton.edu/PTMCuration>

...PTMs are known to act alone and in combination **to regulate nearly all aspects of protein function...**

...Post-translational modifications (PTMs) occur on **nearly all proteins**. Many domains within proteins are **modified on multiple amino acid sidechains** by diverse enzymes to create a myriad of possible protein species. **How these combinations of PTMs lead to distinct biological outcomes is only beginning to be understood...**


A. P. Lothrop, M. P. Torres, S. M. Fuchs, FEBS Letters. 587 (2013) 1247–1257

Protein p53


p53 exerts **irreplaceable anti-neoplastic functions** at homeostasis and thus is considered to be '**the guardian of the genome**'.

p53 is able to coordinate a regulatory network that supervises and responds to a variety of stress signals:

- DNA damage
- aberrant oncogenic activation
- telomere erosion
- ribosomal stress
- loss of cell-cell or cell-matrix adhesion
- hypoxia




Mutations of p53 or disruptions of p53 coordination,
to a lesser extent, can disturb the normal physiological balance,
if genome disarrangement reaches a critical value **it leads to cancer**



Overview of p53 PTMs

Nuclear export



Suppression of Nuclear export



Degradation



Stabilization



Antirepression



Transcriptional activation



Transcriptional suppression



General Enhancement of DNA binding



Promoter-specific Enhancement of DNA binding



- phosphorylation
- ▲ acetylation
- methylation
- ◆ mono-ubiquitination
- ◆◆ poly-ubiquitination
- SUMOylation
- O-GlcNAcylation
- ◆ Neddylation
- dimethylation
- ◆◆◆ ADP-ribosylation

SnapShot: Histone Modifications


He Huang,¹ Benjamin R. Sabari,² Benjamin A. Garcia,³ C. David Allis,² and Yingming Zhao¹


¹Ben May Department of Cancer Research, The University of Chicago, Chicago, IL 60637, USA

²Laboratory of Chromatin Biology and Epigenetics, The Rockefeller University, New York, NY 10021, USA


³Department of Biochemistry and Biophysics, University of Pennsylvania, Philadelphia, PA 19104, USA

Cell





Histone H3 most frequent PTMs



Možnosti MS při analýze modifikací

- Druh
- Lokalizace místa
- Obsazenost daného místa



MS
“screening”
*(paralelní charakterizace tisíců PTM míst
včetně dosud neznámých)*
detailní charakterizace jednotlivých
modifikací



Western blot
detekce druhu PTM
lokalizace jednotlivé modifikace

Specifické barvení gelů
detekce druhu PTM bez možnosti lokalizace
(fosfo, glyko proteiny)

Druhy modifikací:


- mutace (záměna AMK)
- chemické
- posttranslační

Užitečný přehled modifikací:


DeltaMass - <https://www.abrf.org/delta-mass>

„Chemické“ modifikace:

- záměrné modifikace (*karbamidometylace Cys, derivatizace, kvantifikační značky aj.*)
- nechtěné modifikace (*oxidace Met, deamidace N → D při zpracování vzorku, adukty farmak*)



Common Posttranslational Modifications



Common Posttranslational Modifications

| | | | | |
|---|------------------------|-----------|---------------------|-----------|
| Amines (K/N-terminus) | Methylation | +14.0269 | Formylation | +28.0104 |
| | Acetylation | +42.0373 | Lipoic acid | +188.3147 |
| | Farnesylation | +204.3556 | Myristoylation | +210.3598 |
| | Biotinylation | +226.2994 | Palmitoylation | +238.4136 |
| | Stearoylation | +266.4674 | Geranylgeranylation | +272.4741 |
| | | | | |
| Acids & amides (E/D/Q/N) | Pyroglutamic acid (Q) | -17.0306 | Deamidation (Q/N) | +0.9847 |
| | Carboxylation (E/D) | +44.0098 | | |
| Hydroxyl groups (S/T/Y) | Phosphorylation | +79.9799 | Sulphation | +80.0642 |
| Carbohydrates (S/T/N) | Pentoses | +132.1161 | Deoxyhexoses | +146.1430 |
| | Hexosamines | +161.1577 | Hexoses | +162.1424 |
| | N-acetylhexosamines | +203.1950 | Sialic acid | +291.2579 |




Další detaily např:


<http://themedicalbiochemistrypage.org/protein-modifications.php>



MS/MS
fragmentace peptidů
opakování

C7250




CID vs ETD*b, y* *c, z*C-terminus (*z-series*)N-terminus (*c-series*)

MS Charakterizace mutací

Potvrzení výměny AMK na peptidové úrovni

Protein: Zm-p60 (pozice 186)

wild type 179 - 203 **NWLTFNEPQTFTSFSYGTGVFAPGR** **2824**
E186Q **Q** **2823**
 peptide mass difference: **- 1 Da**



Identifikace výměny AMK na aminokyselinové úrovni

Protein ve dvou variantách v jednom spotu na 2-D gelu

Vstupní informace:

Změna hmotnosti tryptického peptidu: **–14 Da**

...

DEEELQKENVKNTASLTGKITLSVTQSKPETGEVIGVFESI**QPSD**TDLGAKVPKDVKIQG

...

MALDI-MS potvrzení změny hmotnosti daného peptidu (2 proteázy)
MALDI-PSD nejednoznačné výsledky

LC-MS/MS nalezena mutace D/E v pozici 210

...


DEEELQKENVKNTASLTGKITLSVTQSKPE**E**TGEVIGVFESI**QPSD**TDLGAKVPKDVKIQG



...

MS/MS peptidu LSVTQSKP~~X~~TGEVIGVFES, MW 2006.0 (1992.0)


„Modif“ - detail MS/MS spektra




Ion b₉, LSVTQSKP~~X~~

D / E


„Nemodif“ – detail MS/MS spektra



EIC chromatogram




EIC chromatogram



LC-MSMS

methylace, potvrzení D v pozici 210
po derivatizaci výměna H za CH₃ na karboxy skupině
tj. $\Delta = 14$ Da/skupinu


D, E a C-terminal
pouze



$$\Delta = 42 \text{ Da}$$

y_8 nederiv m/z 787.2 pro D
 y_8 methyl m/z 829.3 pro D

MS/MS peptidu VTQSKPXTGEVIG před a po methylaci



MS Charakterizace modifikací „chemické“

MALDI-MS spektrum digestů před a po modifikaci

(výřez spektra)


nemodifikovaný peptid

2060

162 Da


Peptid + X

2222



MALDI-MS spektrum – Oxidace Met


(výřez spektra)




Výsledek identifikace proteinu (Mascot)

1. gi|15803837 Mass: 13532 Score: **487** Queries matched: 5
50S ribosomal protein L14 [Escherichia coli O157:H7]

| <i>Observed</i> | <i>Mr(expt)</i> | <i>Mr(calc)</i> | <i>Delta</i> | <i>Miss</i> | <i>Score</i> | <i>Peptide</i> |
|-----------------|-----------------|-----------------|--------------|-------------|--------------|--|
| | | | | | | |
| 939.45 | 1876.89 | 1876.88 | 0.02 | 0 | (125) | MIQEQTMLNVADNSGAR |
| 947.44 | 1892.87 | 1892.87 | -0.01 | 0 | 159 | MIQEQT <u>M</u> LNVADNSGAR + Oxidation (M) |
| 947.45 | 1892.88 | 1892.87 | 0.01 | 0 | (147) | MIQEQT <u>M</u> LNVADNSGAR + Oxidation (M) |
| 955.45 | 1908.89 | 1908.87 | 0.03 | 0 | (118) | MIQEQT <u>M</u> LNVADNSGAR + 2 Oxidation (M) |
| | | | | | | |





Jen část týkající se modifikovaného peptidu
(Petra P. Vz. 3, 080821)



C7250

MS/MS spektrum nemodifikovaného peptidu - MIQEQTMLNVADNSGAR






$b_9 - \mathbf{M}_1\mathbf{I}\mathbf{Q}\mathbf{E}\mathbf{Q}\mathbf{T}\mathbf{M}_7\mathbf{L}\mathbf{N}$

$y_{10} - \mathbf{L}\mathbf{N}\mathbf{V}\mathbf{A}\mathbf{D}\mathbf{N}\mathbf{S}\mathbf{G}\mathbf{A}\mathbf{R}$


$y_{11} - \mathbf{M}_7\mathbf{L}\mathbf{N}\mathbf{V}\mathbf{A}\mathbf{D}\mathbf{N}\mathbf{S}\mathbf{G}\mathbf{A}\mathbf{R}$

Posuny v m/z vybraných fragmentů pro jednotlivé případy $M_{(ox)}$


| | <i>bez</i> | <i>1 Ox</i> | <i>1 Ox</i> | <i>2 Ox</i> |
|----------|----------------------|----------------------|-------------|-------------|
| | M₁ | M₇ | oba | |
| y_{10} | 0 | 0 | 0 | 0 |
| y_{11} | 0 | 0 | 16 | 16 |
| b_9 | 0 | 16 | 16 | 32 |



LC-MSMS spectrum (výřez spektra)
Potrzení modifikace na Cys ...VC^{*}AR




Histon H3, lokalizace acetylace (*in-vitro*) MALDI-MS digestu (detail spektra)



C7250

Histon H3, lokalizace acetylace K(14) MALDI-MS (detail spektra)

ARTKQTARKSTGG**K**A
 c_{14}
 c_{13}



MS Charakterizace modifikací posttranslační

Fosforylace

*Protein Phosphorylation is of Fundamental Importance in Biological Regulation
cca 10-30% of all proteins are phosphorylated*

| | |
|-----------|--------------------|
| ⊕ S, T, Y | 1800 : 200 : 1 ??? |
| ⊕ H | ??? |

Whereas phosphorylation of **serine, threonine or tyrosine** results in the formation of a **phosphoester linkage**, phosphorylation of **histidine** residues occurs **on nitrogen atoms**, producing a phosphoramidate bond. Phosphohistidines have a large standard free energy of hydrolysis making them **the most unstable** of any known phosphoamino acid.

Klumpp et al, Eur. J. Biochem. **269**, 1067-1071 (2002)

Phosphorylation sites db:

<http://www.phosphosite.org/homeAction.do>
<http://phospho.elm.eu.org>

Phospho.ELM version 9.0 (September 2010) contains 8,718 substrate proteins from different species covering 3,370 tyrosine, 31,754 serine and 7,449 threonine instances.

Fosforylace

9 aminokyselin, které mohou být fosforylovány

serin (Ser) > threonin (Thr) > tyrosin (Tyr)

histidin (His)



kys. asparagová (Asp), kys. glutamová (Glu)

lysin (Lys), arginin (Arg), cystein (Cys)



Fosfoproteom - Potížista

- potlačení signálu v MS
 - pouze malá část z celkového množství proteinů je fosforylovaná
(přednostní ionizace nemodifikovaných peptidů)
- většina signálních proteinů je v buňce v nízkých koncentracích
- protein se může vyskytovat v různých fosfo formách
- proteiny mohou být během zpracování defosforylovány fosfatázami přítomnými ve vzorku



Úprava vzorku

- inhibitory fosfatáz (co nejdříve), FASP (lyze v SDT)
- frakcionace fosfopeptidů (proteinů)
 - **TiO₂** (*resp.jiné oxidy kovů, MOAC – „metal oxide affinity chromatography“*)
 - **IMAC** (*„immobilized metal affinity chromatography“*)
 - **SCX** resp. **SAX** or **HILIC** (*„ion exchange or hydrophilic interaction chromatography“*)
 - **imunoprecipitace** pomocí specifické protilátky (zejména Y(fosfo)

I.L. Batalha, Trends in Biotechnology 30 (2), 100-110 (2012)

MS analýza

jiné typy fragmentací

CID

ETD (ECD)

electron transfer (capture) dissociation

HCD


higher-energy collision dissociation

EThcD

electron-transfer/higher-energy collision dissociation

Frese at al., J. Proteome Res., 12, 1520–1525 (2013)

Specifické barvení fosforylovaných proteinů, 2D GE



phosphoproteins
(Pro-Q Diamond , **blue**)
proteins
(SYPRO Ruby, **red**).



alternativa
Metabolické značení ^{32}P
měření radioaktivity

immunoblotting

phosphatase treatment
phosphoproteins display a basic shift in their pI after the dephosphorylation.
comparison 2D gels

Mobility Shift Detection of Phosphorylated Proteins

SDS-PAGE using an Phos-tag™ complex with two manganese(II) ions
 Mn^{2+} -Phos-tag™




- : Phosphorylated proteins (octa-, penta-, and di-phosphorylated, respectively)

+ : Dephosphorylated proteins (AP-treated proteins)

A commercially available β -casein contains partially dephosphorylated proteins.

Immobilized metal affinity chromatography (IMAC)


charging




specific binding

elute

IMAC enrichment of -Casein phosphopeptides (1 pmol of tryptic digest)




Kasein (1 µg) po digesci trypsinem




C7250

Kasein (1 µg) po digesci trypsinem




C7250


Kasein (1 µg) po digesci trypsinem



MALDI-MS spektrum peptidu bez a s fosforylací




Potvrzení fosforylace pomocí alkalické fosfatázy



ESI–MS (IT) spektrum peptidu bez a s fosforylací

positive mode




Fosforylace histonu H4 kinázou Aurora B

(K. Šedová)


- fosforylace
- proteolytické štěpení trypsinem
- frakcionace fosfopeptidů na TiO_2
- LC-MS/MS analýza – sken neutrální ztráty (ETD)

Schéma MS analýzy




C7250



LC-MS chromatogram



MS chromatogram vybraného iontu o m/z 559.0



C7250



Identifikace peptidu databázovým prohledáváním MS/MS Ion Search (MASCOT)

C7250

| | | | |
|------------|--------|----------|-----------|
| <i>m/z</i> | Charge | RT (min) | Expect Mr |
| 559.0 | 3+ | 22.063 | 1674.0 |

MASCOT

- gi|223582 Mass: 11230 Score: 74 Queries matched: 2
histone H4

| Observed | Mr(expt) | Mr(calc) | Delta | Miss | Score | Expect | Rank | Peptide | CID |
|----------|----------|----------|-------|------|-------|--------|------|--|-----|
| 558.98 | 1673.93 | 1673.86 | 0.076 | 2 | (23) | 7.8 | 1 | K.RK <u>T</u> VTAMDVVYALK.R + Phospho (ST) | CID |
| 558.98 | 1673.93 | 1673.86 | 0.076 | 2 | 75 | 5e-05 | 1 | K.RK <u>T</u> VTAMDVVYALK.R + Phospho (ST) | ETD |

Modifications: Optional: Phospho (ST)

Search Parameter: Charge=2+ and 3+, MS Tol.:0.500000 Da, MSMS Tol.:0.500000 Da, Trypsin
Mascot 2.2.03, NCBIInr NCBIInr_20081101.fasta

Biotools

| | Score (Biotools) | Score (Mascot) |
|--|---------------------|-------------------|
|--|---------------------|-------------------|

CID


| | | |
|-------------------------|-----|----|
| RK <u>T</u> VTAMDVVYALK | 520 | 23 |
| RKT <u>V</u> TAMDVVYALK | 518 | 22 |

ETD

| | | |
|-------------------------|------|----|
| RK <u>T</u> VTAMDVVYALK | 6149 | 75 |
| RKT <u>V</u> TAMDVVYALK | 774 | 47 |

RKTVTAMDVVYALK


C7250



na základě CID MS/MS dat nelze fosfokupinu lokalizovat

RK^TVTAMDVVYALK

C7250



na základě ETD MS/MS dat lze jednoznačně lokalizovat fosfokupinu na T(3)

T(3) x T(5)

C7250




TABLE 1 | Phosphopeptide/phosphoprotein enrichment methodologies.

| Enrichment method | Description | Advantage | Disadvantage | References |
|--|---|--|--|------------------------|
| Immunoaffinity enrichment | Use of antibodies directed against pTyr, pSer, pThr, and more recently against the surrounding consensus sequences for pSer/pThr. | Highly specific. | Low efficiency, high cost, use of different antibodies for different phosphorylation motifs. | Stokes et al., 2012 |
| Immobilized metal affinity chromatography (IMAC) | Negatively charged phosphate groups on the phosphorylated amino acids interact with positively charged metal ions such as Ni ²⁺ , Fe ³⁺ , Ga ³⁺ , Zr ⁴⁺ , and Ti ⁴⁺ that are chelated with silica or agarose through nitriloacetic acid or iminodiacetic acid. | Good for both phosphoproteins and phosphopeptides. When used with peptides, it can enrich mono- and multiple phosphorylated peptides. | Tends to bind strongly to monophosphorylated peptides, which makes it difficult for elution. Non-specific binding of acidic peptides can occur. | Fila and Honys, 2012 |
| Metal oxide affinity chromatography (MOAC) | Similar to IMAC, the phosphate groups on the amino acids interact with positively charged metal oxides, e.g., titanium or zirconium that acts as anchoring molecules to trap phosphopeptides through the formation of multi-dentate bonds. | Good for both phosphoproteins and phosphopeptides. When used with peptides, it can enrich mono- and multiple phosphorylated peptides. | Tends to binds strongly to multiple phosphorylated peptides, which makes it difficult for elution. Nonspecific binding of acidic peptides can occur. | Gates et al., 2010 |
| Phos-Tag chromatography, | Uses 1,3-bis[bis(pyridine-2-ylmethyl)amino]propan-2-olato zinc(II) complex as a selective phosphate binding tag in aqueous solution at neutral pH. | Increased sensitivity due to complete deprotonation of phosphoproteins/ phosphopeptides at neutral pH. Elution at the physiological pH allow for protein activity and functional analysis. | Mainly used to confirm the phosphorylation state in relatively pure proteins, but not with complex mixtures. | Kinoshita et al., 2006 |
| Prefractionation by strong cation exchange (SCX) and strong anion exchange (SAX) | In SCX, tryptic peptides often carry a charge of +2, except for phosphopeptides with a net charge of +1, making them elute early in the chromatography. SAX retains phosphopeptides, allowing separation based on the number of phosphorylated residues. | Used for fractionation of highly complex mixtures, it can be performed on-line with mass spectrometry. | Similar degree of unspecific binding as IMAC and MOAC. | Leitner et al., 2011 |
| Hydrophilic interaction liquid chromatography (HILIC) | Phosphopeptides with polar phosphate groups are strongly retained on the HILIC stationary phase resulting in separation from non-phosphorylated species. | Good for both phosphoproteins and phosphopeptides. When used with peptides, it can enrich mono- and multiple phosphorylated peptides. | Similar degree of unspecific binding as IMAC and MOAC. | (Yang et al., 2013) |
| Electrostatic repulsion hydrophilic interaction chromatography (ERLIC) | ERLIC is a variation of HILIC using electrostatic repulsion as an additional phase to adjust selectivity by varying pH or organic solvents. | Good for both phosphoproteins and phosphopeptides. When used with peptides, it can enrich mono- and multiple phosphorylated peptides. | Similar degree of unspecific binding as IMAC and MOAC. | Gan et al., 2008 |
| Hydroxyapatite chromatography | It takes advantage of the strong interaction between positively charged hydroxyapatite and phosphate ions. | Good for fractionating mono-, di-, tri-, and multi-phosphorylated peptides when using gradient of a phosphate buffer. | Developed with phosphoprotein standards, not tested with complex samples. | Mamone et al., 2010 |





Photo Copyright Ralf Langer

AIRLINERS.NET

Ubiquitinace

Ubiquitination is an enzymatic, protein post-translational modification (PTM) process in which the carboxylic acid of the terminal glycine from the **di-glycine motif** in the activated ubiquitin forms an amide bond to the epsilon amine of the lysine in the modified protein.

Protein ubiquitination regulates many cellular processes including transcription, endocytosis, cell cycle control, signal transduction, stress response, DNA repair as well as **proteasomal-mediated degradation**




Ubiquitin linkage types

| Crystal structure of Ub dimer | Physiological function | Examples for E3s and DUBs with preference for certain linkage types |
|-------------------------------|---|---|
| K6 | DNA damage response; Parkin-mediated mitophagy | E3: ERCA1, Parkin DUB: USP30, OTUD3 |
| K11 | Human cell cycle control | E3: APC/C DUB: Cezanne |
| K27 | Nuclear translocation; DNA damage response | E3: RNF168 DUB: unknown |
| K29 | Ub-fusion degradation; Wnt/ β -catenin signaling | E3: Smurfl, UBE3C DUB: TRABID |
| K33 | TCR signaling; post-Golgi trafficking; AMPK-related kinase signalling | E3: Cu3-KLHL20, AREL1 DUB: TRABID |
| K48 | Canonical signal for proteasomal degradation | E3: SCF, E6AP DUB: OTUB1 |
| K63 | Endocytosis; protein trafficking; innate immunity; NF- κ B signalling | E3: TRAF6 DUB: AMSH; OTUD1 |
| M1 | Innate immunity; NF- κ B signalling; angiogenesis; selective autophagy | E3: LUBAC DUB: OTULIN |


Ubikvitinace

(A) Homotypic and heterotypic Ub chains



The complex Ub code contains numerous variants of homotypic and heterotypic (mixed or branched) chains. Based on the eight possible linkages (M1, K6, K11, K27, K29, K33, K48, and K63) between two Ub moieties, **at least 92 different Ub chain types exist.**


Stolz A. et al., *Trends in Cell Biology*, 28 (1), 1-3 (2018)



Akutsu M. et al., *J. Cell Sci.*, 129, 875-880 (2016)

ubikvitin – protein 8.5 kDa (76 AMK)


MQIFV**K**TGTITLEVEPSDTIENV**KAKIQD**K****EGIPPDQQRLIFAG**K**
QLEDGRTLS**DYNIQ**K****ESTLHLV LRLRG**G**



— **K**


- lokalizace modifikovaných AMK
- určení vazeb v polyubikvitinu

heterogenita forem



The most studied polyubiquitin chains - lysine48-linked - target proteins for destruction

Strategie analýzy ubikvitinovaných míst



Chybná identifikace ubikvitinace nebo K(91) je ubikvitinylován ???

BBAP monoubiquitylates histone H4 at lysine 91 and selectively modulates the DNA damage response.["]

Yan Q., Dutt S., Xu R., Graves K., Juszczynski P., Manis J.P., Shipp M.A.
Mol. Cell 36:110-120 (2009)

Histon H4 (trypsin)

| | |
|-----|---|
| 1 | SGRGKGGKGL GKGGAKRHRK VL RDNIQGIT KPAIRRLARR GGVKRISGLI |
| 51 | YEETRGVLKV FLENVIRDAV TYTEHAKRKT VTAMDVVYAL KRQGRTLYGF |
| 101 | GG |

Mascot

79 –100 658.6059 2630.3944 2630.4003 -2.23 0 45 0.0013 1 R.KTVTAMDVVYALKRQGRTLYGF.G + UBI_dT (K)

All matches to this query

| Score | Mr(calc) | Delta | Sequence |
|-------|-----------|---------|-----------------------|
| | 2630.4003 | -0.0059 | |
| | 2630.4003 | -0.0059 | |
| 0.3 | 2630.3792 | 0.0152 | KSAPAPKKGSKKAVTKAQKKD |

Chybná identifikace ubikvitinace nebo K(91) je ubikvitinylován ???

K(1) není ubikvitinylován


KVTAMDVVYAL**K**RQGRTLYGF

| # | b | b^{++} | b^* | b^{*++} | b^0 | b^{0++} | Seq. | y | y^{++} | y^* | y^{*++} | y^0 | y^{0++} | # |
|----|------------------|-----------|-----------|-----------|-----------|-----------|----------|------------------|------------------|-----------|-----------|-----------|-----------|-----------|
| 1 | 129.1022 | 65.0548 | 112.0757 | 56.5415 | | | K | | | | | | | 22 |
| 2 | 230.1499 | 115.5786 | 213.1234 | 107.0653 | 212.1394 | 106.5733 | T | 2503.3126 | 1252.1599 | 2486.2860 | 1243.6467 | 2485.3020 | 1243.1547 | 21 |
| 3 | 329.2183 | 165.1128 | 312.1918 | 156.5995 | 311.2078 | 156.1075 | V | 2402.2649 | 1201.6361 | 2385.2384 | 1193.1228 | 2384.2543 | 1192.6308 | 20 |
| 4 | 430.2660 | 215.6366 | 413.2395 | 207.1234 | 412.2554 | 206.6314 | T | 2303.1965 | 1152.1019 | 2286.1700 | 1143.5886 | 2285.1859 | 1143.0966 | 19 |
| 5 | 501.3031 | 251.1552 | 484.2766 | 242.6419 | 483.2926 | 242.1499 | A | 2202.1488 | 1101.5780 | 2185.1223 | 1093.0648 | 2184.1383 | 1092.5728 | 18 |
| 6 | 632.3436 | 316.6754 | 615.3171 | 308.1622 | 614.3330 | 307.6702 | M | 2131.1117 | 1066.0595 | 2114.0852 | 1057.5462 | 2113.1011 | 1057.0542 | 17 |
| 7 | 747.3706 | 374.1889 | 730.3440 | 365.6756 | 729.3600 | 365.1836 | D | 2000.0712 | 1000.5392 | 1983.0447 | 992.0260 | 1982.0607 | 991.5340 | 16 |
| 8 | 846.4390 | 423.7231 | 829.4124 | 415.2098 | 828.4284 | 414.7178 | V | 1885.0443 | 943.0258 | 1868.0177 | 934.5125 | 1867.0337 | 934.0205 | 15 |
| 9 | 945.5074 | 473.2573 | 928.4808 | 464.7441 | 927.4968 | 464.2520 | V | 1785.9759 | 893.4916 | 1768.9493 | 884.9783 | 1767.9653 | 884.4863 | 14 |
| 10 | 1108.5707 | 554.7890 | 1091.5442 | 546.2757 | 1090.5601 | 545.7837 | Y | 1686.9075 | 843.9574 | 1669.8809 | 835.4441 | 1668.8969 | 834.9521 | 13 |
| 11 | 1179.6078 | 590.3075 | 1162.5813 | 581.7943 | 1161.5973 | 581.3023 | A | 1523.8441 | 762.4257 | 1506.8176 | 753.9124 | 1505.8336 | 753.4204 | 12 |
| 12 | 1292.6919 | 646.8496 | 1275.6653 | 638.3363 | 1274.6813 | 637.8443 | L | 1452.8070 | 726.9071 | 1435.7805 | 718.3939 | 1434.7964 | 717.9019 | 11 |
| 13 | 1534.8298 | 767.9185 | 1517.8032 | 759.4053 | 1516.8192 | 758.9132 | K | 1339.7229 | 670.3651 | 1322.6964 | 661.8518 | 1321.7124 | 661.3598 | 10 |
| 14 | 1690.9309 | 845.9691 | 1673.9043 | 837.4558 | 1672.9203 | 836.9638 | R | 1097.5851 | 549.2962 | 1080.5585 | 540.7829 | 1079.5745 | 540.2909 | 9 |
| 15 | 1818.9895 | 909.9984 | 1801.9629 | 901.4851 | 1800.9789 | 900.9931 | Q | 941.4839 | 471.2456 | 924.4574 | 462.7323 | 923.4734 | 462.2403 | 8 |
| 16 | 1876.0109 | 938.5091 | 1858.9844 | 929.9958 | 1858.0004 | 929.5038 | G | 813.4254 | 407.2163 | 796.3988 | 398.7030 | 795.4148 | 398.2110 | 7 |
| 17 | 2032.1120 | 1016.5597 | 2015.0855 | 1008.0464 | 2014.1015 | 1007.5544 | R | 756.4039 | 378.7056 | 739.3774 | 370.1923 | 738.3933 | 369.7003 | 6 |
| 18 | 2133.1597 | 1067.0835 | 2116.1332 | 1058.5702 | 2115.1492 | 1058.0782 | T | 600.3028 | 300.6550 | | | 582.2922 | 291.6498 | 5 |
| 19 | 2246.2438 | 1123.6255 | 2229.2172 | 1115.1123 | 2228.2332 | 1114.6202 | L | 499.2551 | 250.1312 | | | | | 4 |
| 20 | 2409.3071 | 1205.1572 | 2392.2806 | 1196.6439 | 2391.2965 | 1196.1519 | Y | 386.1710 | 193.5892 | | | | | 3 |
| 21 | 2466.3286 | 1233.6679 | 2449.3020 | 1225.1547 | 2448.3180 | 1224.6626 | G | 223.1077 | 112.0575 | | | | | 2 |
| 22 | | | | | | | F | 166.0863 | 83.5468 | | | | | 1 |

Nelze rozhodnout zda je ubi na K nebo GG na C-terminu

Charakterizace ubikvitinací


Schéma experimentu pomocí imunoprecipitace

B


Ubikvitinace

Obsazenost místa, semikvantifikace

poměr obsazenosti jednotlivých Ubi míst
vzorek vs kontrola



četnost jednotlivých míst řetězení ubikvitinu
vzorek vs kontrola



Charakterizace ubikvitinací

Schéma experimentu pomocí imunoprecipitace II





schéma experimentu


two human cell lines
Hep2
Jurkat

proteasomal inhibitors treatment
BOR - bortezomib
AP15 - b-AP15




Charakterizace ubikvitinací

Schéma experimentu pomocí imunoprecipitace II



Numbers of ubiquitination sites and ubiquitinated proteins identified in the two cell lines ($n = 3$ independent biological replicates).

Numbers on the top of bars indicate identified ubiquitination sites; numbers within bars indicate identified proteins.



Overlap of ubiquitination sites and ubiquitinated proteins between Hep2 and Jurkat cells.

Numbers indicate the number of identified ubiquitination sites (left) or proteins (right).

Ubiquitin-like proteins

Table 1. Ubl's and Their E1 and E2 in Human and Budding Yeast

| family | proteins in <i>H. sapiens</i> | | | | proteins in <i>S. cerevisiae</i> | | |
|--------|---|-----------|--------------------|-------|----------------------------------|-------|--|
| | Ubl | E1 | E2 | Ubl | E1 | E2 | |
| SUMO | SUMO1, SUMO2, SUMO3, SUMO4 ^a | UBA2/SAE1 | UBC9 | Smt3 | Uba2/Aos1 | Ubc9 | |
| NEDD8 | NEDD8 | UBA3/NAE1 | UBC12, UBE2F | Rub1 | Uba3/Ula1 | Ubc12 | |
| ATG8 | LC3A, LC3B, LC3B2, LC3C, GABARAP, GABARAPL1, GATE-16 ^a | ATG7 | ATG3 | Atg8 | Atg7 | Atg3 | |
| ATG12 | Atg12 | ATG7 | ATG10 | Atg12 | Atg7 | Atg10 | |
| URM1 | URM1 | UBA4 | — | Urm1 | Uba4 | — | |
| UFM1 | UFM1 | UBA5 | UFC1 | — | — | — | |
| FAT10 | FAT10 | UBA6 | UBE2Z ^b | — | — | — | |
| ISG15 | ISG15 | UBA7 | UBCH8 ^b | — | — | — | |

^aSUMO5 and GABARAPL3 were not included in this table as they are likely pseudogenes. ^bUBE2Z and UBCH8 can also work with ubiquitin.



Photo Copyright YOSHIAKI HOSHINA


AIRLINERS.NET

Glykosylace

one of the most common post-translational modifications of proteins in eukaryotic cells.

involved in a wide range of biological functions such as receptor binding, cell signaling, immune recognition, inflammation, and pathogenicity.


Základní typy glykanů:



- **N-linked**
- **O-linked**
- **GPI anchors**
 - C-linked
 - glykace


Variation in the degrees of saturation at available glycosylation sites results in heterogeneity in the mass and charge of glycoproteins

Signal Supression



N - linked

O - linked



N-linked glycosylations


glycans are attached to the protein backbone via an amide bond to an asparagine during protein synthesis (in endoplasmic reticulum)

N-X-S(T)


X nesmí být P

subtypes:

- High-mannose
- Hybrid
- Complex



N-linked:High-mannose subtype




β -D-Mannose



β -D-*N*-Acetylglucosamine

N-linked: Hybrid subtype




β -D-Galactose



α -N-Acetylneurameric acid (Sialic Acid)

N-linked: Complex subtype





α -L-Fucose

O-linked glycosylations


glycans are linked via the hydroxyl group of serine or threonine

examples:



β -D-N-Acetylgalactosamine

GPI (glycosylphosphatidylinositol) anchors anchors are linked via C-terminus, membrane bound proteins



Charakterizace glykoproteinů

- specifická detekce glykosylovaných proteinů
- identifikace proteinů
- určení glykosylačního místa
- určení struktury glykanu

Specifická detekce glykosylovaných proteinů

Pro-Q Emerald 300 - glyko only



další techniky detekce:

*kolorimetrická detekce
fluorescenční detekce*

specifické obohacení:

*afinitní chromatografie
(lektinové matrice.
m-Aminophenylboronic Acid)*

identifikace proteinu:

*Peptide mapping
MS/MS Ion search*

Sypro Ruby - all

Deglykosylace

chemická:

Hydrazinolysis

Hydrazine hydrolysis has been found to be effective in the complete release of unreduced O- and N-linked oligosaccharides.

Alkaline β -Elimination - jen O-linked s vyjímkami

Trifluoromethanesulfonic Acid - destrukce glykanu


enzymatická:

PNGase F N-linked, vše pryč, pokud ne

PNGase A N-linked, vše pryč,

Endoglycosidase H N-linked, štípe až za prvním

Endoglycosidase F1, F2, F3 N-linked, štípou specificky ke struktuře



O-Glycosidase O-linked. vše pryč


β -galactosidase štípe před Gal

...

Určení místa glykosylace, resp. struktury glykanů


- **glykosylace „jen“ na S nebo T** (O-linked)
NXS(T) (N-linked)
*lze vtipovat potencionální glyko místa
určité strukturní typy u glykanů (high-mannose....)*
- **kombinace MS a MS/MS technik**
- **separace glykoproteinů resp. glykopeptidů**
- **vhodná deglykosylační strategie**
- **derivativizace glykanů**

MALDI-MS spectrum of glycosylated and non-glycosylated protein size of glycan part




1D GE of protein before and after deglycosylation

confirmation of glycosylation




MALDI-MS spectrum of deglycosylated protein
confirmation of glycosylation




MALDI-MS celých proteinů

C7250




MALDI-MS tryptických digestů

C7250




Detail spekter digestů proteinu před a po deglykosylaci

C7250



Detail spekter digestů proteinu před a po deglykosylaci

C7250



Shrnutí výsledků

tryptický peptid **2796 Da** ...PHIFDYSGS... ,
kde D vzniká z N po deglykosylaci PNGasou A

původní sekvence je tedy ...PHIFNYSGS... (hmotnost 2795 Da)

Peptid potvrzen také LC-MS/MS analýzou (v glykosylovaném vzorku digestu nebyl nalezen)

**Hmotnost glykanu 1170 Da odpovídá
xylose+fucose+3*mannose+2*N-acetylglukosamin**


Nebyl dále potvrzen MS/MS technikami

...missing parts have potential N-glycosylation sites...


MLRNVCVPVLILLIIGATAQDPTDV**G**EAFANVEWSVAELKRV
LVMGVPRDCGELFLSGQ**NHS**GVYNIYPYKDSLLPVS
AYCDMETDGGGWTVFQRRGQFGNPVYYFYKKWA
DYAHGFGDPAKEYWLGNNVLHALTSDKAMSLRIE
KNHSLETLTAEYSVFK**V**ASEEYFKINVGGYIGSK
GSDAFSIANGSMFTASDQDHDTYTNNCAVEFKG
AWYTSCHGSNLNGLNLNGEHPsyADGIEWSAR
GGSTGLYYYSYPNVEMKVRDAHFISRVAADGRAS

from lecture of Petr Man


... glycopeptide...



MS/MS from 1202.2 - - glycopeptide, type of glycan identified




MS/MS from 971.7 - peptide with one HexNAc - site of glycosylation identified



MLRNVCPVLILLIIGATA QDPTDV GEAFANVEWSVAELKRV
LVMGVPR DCGELFLSGQNHSGVYNIYPYKDSLLPVS
AYCDMETDGGGWTVFQRRGQFGNPVYYFYKKWA
DYAHGFGDPAKEYWLGNVLHALTSDKAMSLRIE
KNHSLETLTAEYSVFK VASEEEYFKINVGGYIGSK
GSDAFSIANGSMFTASDQDHDTYTNNCAVEFKG
AWYTSCHGSNLNGLNLNGEHPSYADGIEW SAR
GGSTGLYYY SYPN VEMKVRDAHFISR VADGRAS

from lecture of Petr Man

Kombinace deglykosylačních enzymů



Postupná deglykosylace různými enzymy (MALDI-MS)





Fig. 2. MALDI-MS analysis of fraction T9 (A), after the digestion of fraction T9 with sialidase S (B), followed by β 1-4 galactosidase (C) and then by β 1-2-N-acetylglucosaminidase digestion (D). MALDI-MS analysis of fraction T9 after treatment with α 1-6-fucosidase and α -mannosidase (E).


Glycan profiling and structural analysis of glycans

C7250


Lattová E. et al., J. Proteome Res., 15 (8), 2777-2786 (2016)



NSCLC - Bronchoalveolar Carcinoma



Bronchoalveolar Adenocarcinoma



Large Cell Carcinoma

MALDI-TOF-MS spectra of N-glycans after desialylation

● Man; ○ Gal; ■ GlcNAc; ▲ Fuc

Glycoproteomics of stem cells

example of LC-MS/MS based global study





Figure 1 | Glycoproteomics. **a**, Glycoproteomic workflow combining proteomics platforms with a new algorithm for the identification of intact glycopeptides from complex biological samples. **b**, MS/MS raw data pre-processing by charge deconvolution and precursor mass correction allows the automated identification of glycopeptides, on the basis of low-abundant peptide fragment ions in the lower mass-range of glycopeptide MS/MS spectra (a.m.u., atomic mass unit). **c**, Coverage of experimentally identified glycoproteins (orange circle; Supplementary Table 10) among all predicted mouse glycoproteins (Uniprot, white) and the glycoproteins expressed (on the basis of RNA-seq, grey) in mESC. **d**, Transcript abundance of all glycoproteins expressed (RNA-seq) in our mESC clone (grey), compared with the glycoproteome experimentally detected (orange). Data are representative of two independent mESC glycoproteomics experiments with similar results.

MSFragger-Glyco

example of software development



A to je konec



Bolesa 360
360.bolesa.com

joke.mpp.com