

11. Molecular Biotechnology in Agriculture

Bi7430 Molecular Biotechnology

Outline

- Definition of green biotechnology
- □ Genetic engineering of plants
- □ Genetic engineering of animals
- Biopharming
- **GMO** benefits and controversies

Green (agricultural) biotechnology

- **green biotechnology applied to agricultural processes**
- environmentally-friendly solutions as alternative
 - to traditional agriculture, horticulture, and animal breeding
- modification of plants and animals increasing value in agriculture
 - traditional agriculture selective crossbreeding and hybridization
 - modern molecular biotechnology transgenesis (rDNA)
- **transgenic organism** altered by addition of exogenous DNA
- **transgene** DNA that is introduced

Genetic engineering of plants

- □ > 150 different plant species in 50 countries worldwide
- DNA sequence of A. thaliana (2000), rice (2005), cotton (2006), corn (2009), potato (2011), tomato (2012), etc.
- transgenic plants engineered to

overcome biotic and abiotic stress

- o pesticides (herbicides)
- o pests and diseases (insects, viruses, bacteria, fungi)
- o environmental stress (salt, temperature, cold and drought)

improved crop quality


- o improved nutritional quality
- o enhance taste, appearance and fragrance
- o increase shelf-life

biopharming


- o plants as bioreactors for production of useful compounds
 - (e.g., therapeutics, vaccines, antibodies)
- phytoremediation

Genetic engineering of plants

- plant transgenesis procedure
 - construction of vector/plasmid (restriction digests, ligation)
 - 2. propagation in E.coli
 - 3. transformation
 - 4. culture and selection



totipotency - entire plant generated from a single, non-reproductive cell











Methods of plant transformation

direct methods

- protoplast polyethylene glycol (PEG) method
 - first technique for plant transgenesis
 - PEG induces reversible permeabilization of the plasma membrane


protoplast electroporation

• intensive electrical field leads to pores on plasma membrane

silicon carbide fibers

o fibers punch holes through plant cells during vortexing

protoplast microinjection




Methods of plant transformation

direct methods


particle bombardment

- most common technique for direct transformation
- "particle gun" or "gene gun"
- DNA precipitated onto tungsten or gold particles
- particles shot into the plant tissue/cells










Methods of plant transformation

indirect methods (vectored)

Agrobacterium-mediated transformation

- A. tumefaciens plant pathogenic bacteria causes Crown gall (tumors)
- tumor inducing (Ti) plasmid
- T-DNA transferred and integrated into plant cell



Markers and selection

- transformation frequency is low (less than 3%)
- without selective advantage transformed cells overgrown by non-transformed

selection markers

- antibiotics resistance
 (Kanamycin, Geneticin)
- herbicides resistance (Phosphinothricin)

reporter genes

- GUS (β-glucuronidase)
- GFP (green fluorescent protein)
- LUC (luciferase)





pest and disease resistance

- **toxin gene** from *Bacillus thuringiensis*
 - o Bt-corn resistant to European corn borer
 - Bt-cotton resistant to cotton bollworm
 - Bt-peanut resistant to cornstalk borer
- Papaya ringspot virus resistance inserting gene from pathogen into crop affords the crop plant resistance







herbicide resistance

- herbicide target modification
- herbicide target overproduction
- herbicide detoxification (enzymatic)

EXAMPLES

- sulfonylurea resistance
 blocking the enzyme for synthesis Val, Leu, isoLeu
 mutated gene transferred from resistant tabaco
- bromoxynil resistance
 - transgene encoding enzyme bromoxynil nitrilase
- glyphosinate resistance
 - bacterial transgene protein inactivating herbicide







- resistance to environmental stress
- marginal land or climate change induced drought
- crucial ways of securing the world's food supply


drought tolerance

- gene from Xerophyta viscosa unique protein in cell membrane
- gene for production of protective waxy cuticle on leaves
- gene for expression of trehalose (stabilization of biomolecules)

salt tolerance

- gene for enhanced glycinebetaine production





improved crop quality

- higher nutrition value
 - golden rice (beta carotene genes)
 120 million children suffers from vitamin A deficiency healthy vision and prevents night blindness
 - black tomato (anthocyanin antioxidant gene)
 prevent heart disease, diabetes and cancer

improve shelf life

delayed fruit ripening (FlavrSavr tomato)
 antisense gene blocking pectinase

improved appearance

- delphinine gene from pansy cloned to rose
- biopharming











Genetic engineering of animals

selective breeding

- time consuming and costly
- Iimited number of properties available
- difficult to introduce new genetic traits / lines

transgenic animals

- fast generation lines carrying desired properties
 - o increased growth
 - o improved disease resistance
 - improved nutritional quality
 - o increased wool quality
- model animals for human disease research
- biopharming production of useful molecules
- biosensors for environmental pollution








direct microinjection (pronucleus method)

- injection of desired DNA to male pronucleus
- most popular, commercial available
- success range from 10 to 30%
- transfer of large genes possible
- no theoretical limit for gene construct size
- random insertion of the transgene

(affecting other genes and expression patterns)






retrovirus mediated gene transfer

- retroviruses used as vectors (gene therapy)
- virus gene is replaced by transgene
- replication defective virus infect host cells (e.g., ES cells, embryo cells)
- efficient mechanism of transgene integration
- transfer of genes < 8 kb only possible</p>
- random insertion of the transgene








embryonic stem cell method

- transfection of gene construct into *in vitro* culture of embryonic stem (ES) cells
- ES recombinant cells incorporated into embryo at blastocyst stage
- 1 in a million incorporated at desired position
- ES cell lines not available in farm animals
- random insertion of the transgene








engineered nucleases, "molecular scissors"


- site-specific double stranded breaks
- Zinc finger nucleases (ZFNs)
- transcription-activator like effector nucleases (TALEN)
- RNA-guided DNA endonuclease (CRISPR-Cas9)




CRISPR-Cas9

- synthetic guide RNA (gRNA)
- delivering Cas9 nuclease complexed with gRNA into a cell
- *in vivo* (nucleus), stem cells, fertilized egg
- can target several genes at once







Application of transgenic animals

disease-resistant livestock

- *in vivo* immunization overexpress genes encoding monoclonal antibodies
- eliminate production of host cell components interacting with infectious agent

improving milk quality

- increase casein contents let to increase cheese production
- decrease lactose content by overexpress lactase
- abolish lacto globulin expression (for milk allergic consumer)

improving animal production traits

- transgenic fish enhanced growth 3-5 times (growth hormone)
- transgenic pig production of omega-3-fatty acids (roundworm gene)
- transgenic poultry lower cholesterol and fat in eggs

biopharming

use of plants or animals for the production of useful molecules

industrial products

- proteins (enzymes)
- fats and oils
- polymers and waxes

pharmaceuticals

- recombinant human proteins
- therapeutic proteins and pharmaceuticals
- vaccines and antibodies

industrial products from plants

- cheap and easy to produce
- free of animal viruses
- risk of food supply contamination
- environmental contamination
- **EXAMPLES** (transgenic corn, Sigma):
 - trypsin
 - o traditionally isolated from bovine pancreas
 - first large scale transgenic plant product
 - o worldwide market = US\$120 million
 - avidin
 - o medical diagnostics
 - β-glucuronidase
 - o visual marker in research labs





edible vaccines from plants

- no purification required
- no hazards associated with injections
- may be grown locally where needed
- no transportation costs
- no need for refrigeration or special storage
- **EXAMPLES**:
 - HIV-suppressing protein in spinach
 - rabies virus G protein in tomato
 - vaccine for rotavirus or hepatitis in **potato**







plant-made antibodies

- plantibodies monoclonal antibodies produced in plants
- free from potential contamination of mammalian viruses
- plants used include tobacco, corn, potatoes, soya and rice
- EXAMPLES: cancer, herpes simplex virus

plant-made pharmaceuticals

- therapeutic proteins and intermediates
- EXAMPLES: proteins to treat cystic fibrosis, HIV, hypertension

production of pharmaceuticals in milk

- easy to purify few other proteins in milk
- dairy cattle produce 10,000 liters of milk/year (35 g protein/liter)
- only few transgenic cows can meet worldwide demand
- risk of food supply contamination
- **EXAMPLES**:
 - COW: human serum albumin, human lactoferrin
 - SHEEP: alpha-1-antitrypsin
 - GOAT: human antithrombin III (FDA approved), tissue plasminogen activator, malaria antigen

production of materials in milk

BioSteel from spider silk (Nexia Biotech)



GMO benefits

crops

- increased stress tolerance
- improved resistance to disease, pests and herbicides
- increased nutrients, yields, enhanced taste and quality

animals

- improved animal health, resistance, productivity and feed efficiency
- better yields of meat, eggs, and milk

environment

- more efficient processing
- conservation of soil, water, and energy
- better natural waste management

society

- increased food security for growing populations
- climate change induced drought

GMO controversies

safety

- human health toxicity, allergens, antibiotic resistance, unknown effects
- environment unintended transfer through cross-pollination, unknown effects on other organisms, loss of biodiversity

ethics


- tampering with nature by mixing genes among species / cloning
- violation of natural organism's intrinsic values
- stress for animals

access and intellectual property

- domination of world food production by few companies
- increasing dependence on industrialized nations by developing countries

GMO future

- GMO crop first commercialized in 1996
- **17.3** million farmers grew biotech crops on 170 million hectares
- 90% of new users are small resource-poor farmers in developing countries
- EU research on risk of GMOs over the past two decades unable to detect any risks that have not yet been known from conventional agriculture*



* EU Commission (2012): A Decade of EU-funded GMO Impacts Research



Reading

- U.S. Agency for International Development, Agricultural Biotechnology Support Project II, and the Program for Biosafety Systems
- How are Biotech Crops & Foods Assessed for Safety?, Developing a Biosafety System (BRIEF #5 and #6)

